

Plant Gene Register

Complete Nucleotide Sequence of Potato Tuber Acid Invertase cDNA

Dingbo Zhou, Autar Mattoo, Ning Li, Hidemasa Imaseki, and Theophanes Solomos*

Department of Horticulture, University of Maryland, College Park, Maryland 20742 (D.Z., T.S.); Plant Molecular Biology Laboratory, United States Department of Agriculture, Beltsville, Maryland 20705 (A.M., N.L.); and National Institute for Basic Biology at Okazaki, Japan (H.I.)

Acid invertase (EC 3.2.1.26), which hydrolyzes Suc to Glc and Fru, plays an important role in regulating sugar concentrations in many plant storage organs. Its activity is inversely related to Suc concentration but positively related to Glc and Fru concentrations in many plant tissues. Acid invertase activity increases severalfold during storage of potato (*Solanum tuberosum* L.) tubers at low temperatures (Pressey, 1969; Sasaki et al., 1971). Further, the ratio of the increments of Glc to Fru during sweetening of potato tubers at low temperatures is close to 1, indicating that Suc is hydrolyzed via invertase (Zhou, 1994). On the basis of the lack of correlation between levels of sugars and acid invertase activity in potato tubers after 3 months of storage at low temperature, Pressey (1969) suggested that invertase activity was regulated by an invertase-invertase inhibitor system. However, Richardson et al. (1990) demonstrated that during the initial storage of potatoes at low temperature, changes in total acid invertase activity assayed after destroying the endogenous invertase inhibitor reflected the sugar changes more closely than did the basal activity assayed with the inhibitor present. This raises questions concerning both the *in vivo* role of the invertase inhibitor and the contribution of acid invertase to the accumulation of reducing sugars in potato tubers stored at low temperature.

To study the regulatory mechanism by which acid invertase is regulated, a cDNA library of Uni-Zap XR (Stratagene) made from RNA isolated from cold-stored potato tubers was constructed. Six cDNA clones were isolated by screening the library with radiolabeled mung bean acid invertase cDNA probe (Arai et al., 1992) prepared by PCR. The sizes of these clones are between about 2.0 and 2.4 kb. The longest cDNA, pPAI11, was sequenced. It is 2336 bp long and contains one open reading frame extending from an ATG start codon at position 9 to a TAA stop codon at position 1926, encoding a polypeptide of 639 amino acid residues. The features of this cDNA are summarized in Table I.

Sequence analysis shows that pPAI11 is 60 to 95% identical to carrot, mung bean, and tomato acid invertase (Sturm and Chrispeels, 1990; Arai et al., 1992; Klann et al., 1992; Elliott et al., 1993; Sato et al., 1993). The deduced amino acid sequence is 97.9% similar and 95.6% identical to a tomato

Table I. Characteristics of acid invertase cDNA from potatoes

Organism:	<i>Solanum tuberosum</i> L. cv Russet Burbank.
Genome Location:	Nuclear genome.
Gene Copy Number:	Single copy.
Gene Product:	Acid invertase (β -fructosidase, EC 3.2.1.26).
Source:	A cDNA library in Uni-Zap XR constructed with poly(A) ⁺ RNA isolated from potatoes stored at low temperature.
Sequencing Technique:	Dideoxy nucleotide chain-termination method was used to completely sequence both strands.
Method of Identification:	Screening with mung bean acid invertase cDNA. Nucleotide and amino acid sequence comparison with other plant acid invertases.
Expression and Regulation:	Northern blot detected the 2.3-kb acid invertase transcript in potato tubers stored at 1°C, but not in potatoes stored at 10°C.
Features of the Predicted Amino Acid Sequence:	The open reading frame encodes 639 amino acid residues. The deduced amino acid sequence contains a putative signal sequence with a predicted cleavage site at residue 50 (Heijne, 1986; Elliott et al., 1993). The β -fructosidase motif is found in bacterial, yeast, and other plant invertases (Sturm and Chrispeels, 1990). There are six potential N-linked glycosylation sites (Asn-X-Ser/Thr):

fruit vacuolar acid invertase cDNA (Klann et al., 1992; Elliott et al., 1993). Southern blot analysis indicated that the potato acid invertase gene is probably present in one copy per haploid genome. Northern blot analysis showed that the 2.3-kb acid invertase transcript was detected in potato tubers stored at 1°C, suggesting that increases in acid invertase activity in potato stored at low temperatures is regulated at the transcription level.

NOTE ADDED IN PROOF

After the submission of this paper we became aware of the cloning of an apoplastic invertase from potato (P. Hedley, G.C. Marchray,

* Corresponding author; fax 1-301-314-9308.

H.V. Davies, L. Burch, and R. Waugh, unpublished results). The amino acid sequence of this invertase is 61.1% similar to the acid invertase reported here.

ACKNOWLEDGMENTS

We thank Drs. M. Tucker and M. Swegle for their technical advice. We thank Mr. M. Reinsel and Mr. P. Kalaitzis for their assistance in synthesizing oligonucleotides. We would also like to thank the Maryland Biotechnology Institute for its support of the computer facility.

Received March 21, 1994; accepted April 4, 1994.

Copyright Clearance Center: 0032-0889/94/106/0397/02.

The GenBank accession number for the sequence reported in this article is L29099.

LITERATURE CITED

- Arai M, Mori H, Imaseki H** (1992) Cloning and sequencing of cDNA for an intracellular acid invertase from etiolated hypocotyls of mung bean and expression of the gene during growth of seedlings. *Plant Cell Physiol* **33**: 245-255
- Elliott JK, Butler WO, Dickinson CD, Konno Y, Vedvick TS, Fitzmaurice L, Mirkov TE** (1993) Isolation and characterization of fruit vacuolar invertase genes from two tomato species and temporal differences in mRNA levels during fruit ripening. *Plant Mol Biol* **21**: 515-524
- Heijne GV** (1986) A new method for predicting signal sequence cleavage sites. *Nucleic Acids Res* **14**: 4683-4690
- Klann E, Yelle S, Bennett AB** (1992) Tomato fruit acid invertase complementary DNA nucleotide and deduced amino acid sequences. *Plant Physiol* **99**: 351-353
- Pressey R** (1969) Role of invertase in accumulation of sugars in cold-stored potatoes. *Am Potato J* **46**: 291-297
- Richardson DL, Davies HV, Ross HA, Mackay GR** (1990) Invertase activity and its relation to hexose accumulation in potato tubers. *J Exp Bot* **41**: 95-99
- Sasaki T, Tadokoro K, Suzuki S** (1971) Multiple forms of invertase of potato tubers stored at low temperature. *Phytochemistry* **10**: 2047-2050
- Sato T, Iwatsubo T, Takahashi M, Nakagawa H, Ogura N, Mori H** (1993) Intercellular localization of acid invertase in tomato fruit and molecular cloning of a cDNA for the enzyme. *Plant Cell Physiol* **34**: 263-269
- Sturm A, Chrispeels MJ** (1990) cDNA cloning of carrot extracellular β -fructosidase and its expression in response to wounding and bacterial infection. *Plant Cell* **2**: 1107-1119
- Zhou D** (1994) Biochemical aspects of low oxygen storage of potato tubers at low temperature and molecular cloning of potato acid invertase. PhD thesis. University of Maryland, College Park