

# Genetic and Transgenic Evidence That Phytochromes A and B Act to Modulate the Gravitropic Orientation of *Arabidopsis thaliana* Hypocotyls

Paul R. H. Robson\* and Harry Smith

Department of Botany, University of Leicester, Leicester, LE1 7RH, United Kingdom

---

Hypocotyls of *Arabidopsis thaliana* exhibit negative gravitropism in the dark, growing against the gravity vector. The direction of growth is randomized in red light (R). In single mutants lacking either phytochrome A or B randomization of hypocotyl orientation in R is retained. However, a double mutant lacks this response, indicating that either phytochrome A or B is capable of inducing randomization and phytochrome A and B are the only phytochromes involved in this process. The induction of randomization was confirmed using lines that express to different levels *PHYA* and *PHYB* cDNAs. Overexpression of *PHYA* cDNAs induced randomization of hypocotyl orientation in the dark. Dark randomization was also seen in the *phyB-1* mutant but not in two other *phyB* alleles, suggesting that dark randomization in the *phyB-1* line may be due to a second mutation. When germination was induced by gibberellin, rather than exposure to brief white light, randomization in the dark associated with phytochrome A overproduction was not observed but was retained in the *phyB-1* mutant. Overexpression of *PHYB* cDNAs induced a light-dependent randomization of hypocotyl orientation that responded to R:far-red light ratio. We conclude that the default situation in *Arabidopsis* hypocotyls is, therefore, negative gravitropism, and either phytochrome A or phytochrome B can mediate randomization.

---

The phytochromes are a family of photoreceptors having multiple functions in the regulation of plant processes in response to the light environment (Smith and Whitelam, 1990; Smith, 1994a). Elucidation of these functions has been greatly helped by the selection of mutants deficient in the action of a particular phytochrome. *phyA* functions primarily in regulating the FR high-irradiance response, whereby continuous FR inhibits hypocotyl elongation as a function of fluence rate (Nagatani et al., 1993; Parks and Quail, 1993; Whitelam et al., 1993). Other processes in which a role for *phyA* has been shown include the regulation of germination (Shinomura et al., 1994) and the promotion of flowering by end-of-day FR treatment (Johnson et al., 1994). *phyB* functions in a number of different processes, including seed germination; gravitropism; cotyledon unfolding; hypocotyl, stem, and petiole elongation; the transition from vegetative to reproductive growth; and responses of light-grown seedlings to the relative amounts of R and FR (Smith and Whitelam, 1990; Goto et al., 1991; Whitelam and Smith, 1991; Liscum and Hangarter, 1993;

Reed et al., 1993; Robson et al., 1993; Neff and Van Volkenburgh 1994; Shinomura et al., 1994). In the light-grown plant, many of the processes mediated by *phyB* contribute to the shade-avoidance syndrome, in which plants detect FR reflected from neighbors, resulting in growth changes appropriate to the competition for light (Smith, 1994a, 1995).

*Arabidopsis* hypocotyls exhibit negative gravitropism when grown in the dark, growing in the opposite direction to the gravity vector; however, the direction of growth is highly randomized when seedlings are exposed to continuous R. Liscum and Hangarter (1993) showed that the *phyB-1* mutant of *Arabidopsis* exhibits randomized growth under both continuous R and dark growth conditions and concluded that the photocontrol of negative gravitropism acts through *phyB*. The *hy2* mutant expresses wild-type levels of phytochrome apoproteins but lacks photoreversible holoproteins due to a presumed deficiency in the biosynthesis of the phytochrome chromophore (Parks and Quail, 1991). This mutant was shown to exhibit negative gravitropism in both the dark and R (Liscum and Hangarter, 1993). Phytochrome molecules exist in one of two forms: one that absorbs R, Pr, and one that absorbs FR, Pfr. The two forms interconvert on absorption of appropriate wavelengths of light. It has been demonstrated at least for *phyA* that the apoprotein is synthesized in the R-absorbing form, PrA (Parks et al., 1987). It is assumed that phytochromes synthesized in the *hy2* mutant are present in a form similar to Pr but lack a chromophore and cannot photoconvert to Pfr. These observations allowed the conclusion that it is PrB that induces negative gravitropism and that in WT it is the depletion of PrB by photoconversion to PfrB, rather than the production of PfrB per se, that attenuates hypocotyl gravitropism resulting in highly randomized growth (Liscum and Hangarter, 1993).

Liscum and Hangarter (1993) did not investigate the gravitropic responses of mutants simultaneously deficient in both *phyA* and *phyB*. In this paper we show that *phyA/phyB* double mutants completely lack the R-mediated randomization of growth orientation. In addition to null mutants, transgenic lines overexpressing *PHYA* or *PHYB*

---

Abbreviations: FR, far-red light; LED, light-emitting diode; *phyA*, phytochrome A; *phyB*, phytochrome B; R, red light; VLFR, very low fluence response; W, white light; WT, wild-type. Nomenclature for phytochrome genes and products is per Quail et al. (1994).

---

\* Corresponding author; e-mail pxr@le.ac.uk; fax 44-116-252-2791.

cDNAs have proved useful in determining the functions of individual phytochromes (Boylan and Quail, 1991; Wagner et al., 1991; Smith, 1994b). In this paper we also examine the effect of overexpressing *PHYA* and *PHYB* cDNAs on the gravitropic response. If PrB induces negative gravitropism, as proposed by Liscum and Hangarter (1993), then high levels of phyB expressed by the transgene should maintain negative gravitropism under light treatments that result in randomization of growth in WT because the cellular concentration of PrB would be high under such light conditions. We also used lines expressing introduced *PHYA* cDNAs to probe the role of phyA in the induction of gravitropic randomization. The use of several lines expressing cDNAs at different levels allows a comparison of response measured with level of phytochrome expression and also excludes the possibility that an observed phenotype is due to insertional inactivation by the transgene.

## MATERIALS AND METHODS

### Plant Material and Growth Conditions

Seeds of *Arabidopsis thaliana* (L.) Heynh. were obtained from the WT and five homozygous lines derived from the ecotype Nossen. The transgenic lines designated Z15, ABO, and D2 had been transformed with an *Arabidopsis phyB* cDNA fused to the cauliflower mosaic virus 35S promoter. Immunoblot quantification indicated that the *PHYB* transgenic lines contained phyB at 15-fold (Z15), 30-fold (ABO), and 60-fold (D2) the light-grown wild-type level of phyB (D. Wagner and P. H. Quail, personal communication). The *PHYA* transgenic lines, designated 13K7 and 21K15, had been transformed with oat *PHYA* fused to the cauliflower mosaic virus 35S promoter. Quantification was less precise in the *PHYA* lines, but the sequence of expression level was WT < 13K7 < 21K15 (M.T. Boylan and P.H. Quail, personal communication). Phytochrome mutants in the Landsberg *erecta* ecotype were *phyA-2*, *phyB-1*, and *phyB-7*; the double mutant was *phyA-2/phyB-1*. Phytochrome mutants in the RLD ecotype were *phyA-101* and *phyB-6*; the double mutant was *phyA-101/phyB-6*. Seeds were incubated for 4 d at 4°C on BG11 (Stanier et al. 1971) agar medium in Petri plates and then exposed to fluorescent white light for 30 min to induce uniform germination. The plates were vertically oriented to allow the seedlings to grow along the surface of the agar and were incubated in darkness at 25°C for 1 d before transfer to one of the treatments described below. After 3 d plates were photographed, and the orientation of the hypocotyl was measured. When the 30-min W pretreatment was replaced by GA pretreatment, seeds were soaked in a solution of 500 µg/mL GA in the dark for 1 h, washed twice with water before immediate plating onto vertically oriented Petri plates, and transferred to dark growth conditions.

### Light Sources

R was obtained from LEDs with an emission maximum at 665 nm and a half-bandwidth of 27 nm and provided a fluence rate of 22 to 24 µmol m<sup>-2</sup> s<sup>-1</sup> at the seed surface; R was given either continuously or as two treatments of 2 h per 24-h period. A range of R:FR ratios was obtained using

a mixed array of both R and FR LEDs (Quantum Devices, Barneveld, WI). The R LEDs were as above. The emission maximum of the FR LEDs was 730 nm and a half-bandwidth of 29 nm. The mixed array provided a fluence rate of 20 to 22 µmol m<sup>-2</sup> s<sup>-1</sup> at wavelengths between 400 and 800 nm and was given as two 1-h periods per 24-h period. FR was obtained from the mixed array with the R LEDs switched off, and the resulting FR light was filtered through black Plexiglas. The resulting light was provided continuously and at a fluence rate of 11 to 13 µmol m<sup>-2</sup> s<sup>-1</sup>. Fluence rates at the level of the seedlings were measured with an Analytical Spectral Devices personal spectrometer model PS2 000A (Analytical Spectral Devices, Boulder, CO). The phytochrome photoequilibrium set up by the R:FR source was measured as a bioassay using a sample of isolated oat phyA in an Aminco dual-wavelength spectrophotometer model DW2A (American Instrument Co., Silver Spring, MD).

### Measurement of Growth Orientation and Statistical Methods

After treatment the plates were photographed, and the image was projected and traced along with a horizontal reference onto paper, from which angle measurements were taken. The angle of orientation was expressed according to the method of Liscum and Hangarter (1993), who found that in darkness and in R values were normally distributed around 0° (vertical). Angles to the left of vertical were assigned a negative value; those to the right were assigned a positive value. The SD was used as a measure of randomization around the vertical (0°).

## RESULTS AND DISCUSSION

### The Effects of phyA and phyB Deficiency on Gravitropism

Results similar to those obtained by Liscum and Hangarter (1993) for single null mutants are shown in Tables I and II. Lack of phyA had little effect compared to WT, with strong negative gravitropism under dark growth condi-

**Table I.** Hypocotyl orientation of phytochrome mutant and wild-type *Arabidopsis* seedlings in ecotype Landsberg *erecta* grown in the dark or exposed to continuous R

Seedlings and data were handled as described in "Materials and Methods." The orientation of hypocotyl angle was measured as the SD around the vertical 0°. Values represent the pooled SD for a minimum of 150 seedlings from at least three replicate experiments. High SDs indicate randomization of hypocotyl growth.

Genotype	Treatment	SD
Landsberg <i>erecta</i>	Dark	20.0
Landsberg <i>erecta</i>	Continuous R	58.3
<i>phyA-2</i>	Dark	16.6
<i>phyA-2</i>	Continuous R	56.3
<i>phyB-1</i>	Dark	79.4
<i>phyB-1</i>	Continuous R	92.8
<i>phyB-7</i>	Dark	22.1
<i>phyB-7</i>	Continuous R	76.6
<i>phyA-2/phyB-1</i>	Dark	23.4
<i>phyA-2/phyB-1</i>	Continuous R	27.1

**Table II.** Hypocotyl orientation of phytochrome mutant and wild-type *Arabidopsis* seedlings in ecotype RLD grown in the dark or exposed to continuous R

Seedlings and data were handled as described in "Materials and Methods." The orientation of hypocotyl angle was measured as the SD around the vertical 0°. Values represent the pooled SD for a minimum of 150 seedlings from at least three replicate experiments. High SDs indicate randomization of hypocotyl growth.

Genotype	Treatment	SD
RLD	Dark	24.4
RLD	Continuous R	83.4
<i>phyA-101</i>	Dark	9.9
<i>phyA-101</i>	Continuous R	78.8
<i>phyB-6</i>	Dark	22.1
<i>phyB-6</i>	Continuous R	58.1
<i>phyA-101/phyB-6</i>	Dark	7.9
<i>phyA-101/phyB-6</i>	Continuous R	13.22

tions and randomization under continuous R. In *phyB*-deficient mutants, randomization of hypocotyl angle under continuous R was seen. Under dark growth conditions there was a difference between the Landsberg *erecta* and the RLD ecotypes, with dark randomization occurring in the *phyB-1* allele of the former but not the *phyB-6* allele of the latter. Table I shows that the *phyB-7* allele of Landsberg *erecta* is consistent with the *phyB-6* mutant, being negatively gravitropic in the dark. We conclude, therefore, that randomization in the dark is caused by a second mutation in the *phyB-1* mutant and that a mutant that is deficient in *phyB* alone will show negative gravitropism in the dark and randomization in continuous R.


In the absence of both phytochromes (i.e. in the *phyA/phyB* double mutant) no randomization of growth was seen under continuous R or in dark growth conditions. The behavior of the double mutant is crucially important for interpretation. Because there is no randomization by continuous R, it is clear that no other phytochrome than *phyA* or *phyB* is required for this response. This means that the interpretation reasonably reached by Liscum and Hangarter (1993) concerning the data available to them, namely that PrB imposes negative gravitropism in the WT, is untenable. If PrB were necessary for negative gravitropism, then the growth orientation of the *phyA/phyB* double mutants would be randomized. We conclude, therefore, that either *phyA* or *phyB* is capable of randomizing growth orientation and that negative gravitropism is the default situation in the absence of both phytochromes. To probe the relative importance of *phyA* and *phyB* in hypocotyl growth orientation, we investigated the responses of transgenic seedlings that produce increased levels of either *phyA* or *phyB*.

### The Effects of *phyB* on Gravitropism

*Arabidopsis* lines that overexpress *PHYB* cDNAs were used to test two alternative hypotheses. First, according to the hypothesis of Liscum and Hangarter (1993), in which PrB is proposed to impose negative gravitropism, increasing levels of *phyB* would be predicted to lead to negative gravitropism under radiation conditions that induce ran-

domization in WT. In other words, with high levels of *phyB*, higher levels of PrB would be present at all photo-equilibria, indicating that R would lead to lower levels of randomization than are achieved in WT plants. The alternative hypothesis, suggested by the results of the single and double null mutants in Tables I and II, is that PrB induces randomization. In this scenario, higher levels of *phyB* might be expected to lead to increases in randomization of hypocotyl angle upon treatment with R, because of a higher cellular concentration of PrB.

Figure 1 shows the effect of two pulses of R of 2 h duration per 24-h period on the distribution of hypocotyl angle in three transgenic lines expressing different levels of *phyB* (Z15, ABO, D2) and their isogenic WT (Nossen). This low-level R treatment was chosen because we had previously determined that it provided intermediate, nonsaturating responses. Contrary to the result predicted by the Liscum and Hangarter hypothesis, the values for hypocotyl angle in seedlings expressing lower levels of *phyB* (Nossen and Z15) are clustered around the vertical, displaying negative gravitropism, whereas seedlings expressing higher




**Figure 1.** Frequency histogram showing hypocotyl orientation of WT and transgenic lines expressing different levels of *phyB* after exposure to two periods of 2 h of R per 24-h period, between which seedlings were maintained in darkness. Seedlings and data were handled as described in "Materials and Methods." Hypocotyl orientation of lines expressing low levels of *phyB*, WT (A) and Z15 (B), was centered near vertical 0°, whereas lines expressing higher levels of *phyB*, ABO (C) and D2 (D), hypocotyl orientation were greatly randomized.

levels of *phyB* (ABO and D2) exhibit greater randomization of hypocotyl angle, or loss of negative gravitropism. This indicates that high levels of *phyB* do not induce negative gravitropism through PrB. Table III shows the sds calculated from the angles of hypocotyls grown in the dark, under continuous R, and under continuous FR. The four strains do not appear to be different from one another in any growth regime, indicating that overexpression of a *phyB* cDNA does not affect the wild-type response, i.e. negative gravitropism in the dark, and that continuous R was sufficient to cause complete randomization in all strains. Under continuous FR the level of randomization was greater than that seen in the dark but less than that seen in continuous R, suggesting that PfrB is responsible for randomizing hypocotyl orientation.

To test whether randomization is a function of Pfr concentration, we studied the effect of different phytochrome photoequilibria, set up by varying the R:FR ratio of the incident light, on the sds of hypocotyl growth around 0° (vertical) (Fig. 2). Under the highest R:FR ratio the sds of hypocotyl angle are as follows: in Nossen, 23.3; in Z15, 33.2; in ABO, 71.6; and in D2, 65.6. Under the lowest R:FR ratios the sds decrease to 13.8, 17.8, 35.9, and 36.4, respectively. These data indicate that decreasing the phytochrome photoequilibrium by decreasing the R:FR ratio causes a decrease in the sds of hypocotyl orientation. The decrease is more pronounced in lines with high levels of *phyB* (ABO and D2); however, all lines are responsive to the R:FR ratio. Higher levels of expression of the introduced *PHYB* gene result in larger sds, or greater randomization, under all R:FR ratios.

These results indicate that the removal of PrB is not the mechanism responsible for the loss of negative gravitropism in R but, conversely, that PfrB induces randomization. This randomization, when *phyB* is present at high



**Figure 2.** The sds of hypocotyl orientation around the vertical of WT and transgenic lines expressing different levels of *phyB* under different R:FR ratios. Light was provided as two pulses of 1 h per 24-h period, between which seedlings were maintained in darkness. The phytochrome photoequilibrium was measured as described in "Materials and Methods." Seedlings and data were handled as described in "Materials and Methods." The sds of hypocotyl orientation of lines expressing low levels of *phyB*, WT (●) and Z15 (■), were low, indicating that orientation was clustered around the vertical. The sds of hypocotyl orientation of lines expressing high levels of *phyB*, ABO (▲) and D2 (▼), were high, indicating randomization and lack of negative gravitropism. All strains appear to respond to the R:FR ratio, showing decreased sds of hypocotyl angles under low R:FR ratio compared to high R:FR ratio.

levels, is sufficient to overcome the default mechanism that imposes negative gravitropism in the absence of both *phyA* and *phyB*.

### The Effects of *phyA* on Gravitropism

Comparison of the results for the *phyB* single mutants and the *phyA/phyB* double mutants in Tables I and II indicates that *phyA* is also capable of inducing randomization. In the RLD ecotype the sd of hypocotyl angles in the *phyB* mutant increases from 22.1 to 58.1, comparing dark and continuous R growth conditions. Even in the presence of the second-site mutation in the *phyB-1* mutant of Landsberg *erecta*, an increase in randomization is seen in R from 79.4 to 92.8, which must be due to the action of *phyA*.

Table IV shows the sd of hypocotyl orientation of WT and transgenic *Arabidopsis* expressing increased levels of *phyA*. Overexpression of *PHYA* cDNAs leads to randomization of hypocotyl orientation. Unlike overexpression of *PHYB* cDNAs, this randomization is seen in the dark and is largely unaffected by growth under continuous R or FR, although higher levels of *phyA* are associated with higher degrees of randomization under all growth conditions. Thus, as the level of *phyA* relative to that of *phyB* increases, there is an increased tendency to randomization in the dark, or under continuous FR, so that in the highest expressers (21K15), randomization in the dark is as great as that under continuous R. All treatments involved a pre-

**Table III.** Effect of dark, continuous R, and continuous FR on growth orientation of wild-type and transgenic lines expressing different levels of *phyB*

Seedlings and data were handled as described in "Materials and Methods." The orientation of hypocotyl angle was measured as the sd around the vertical 0°. Values represent the pooled sd for a minimum of 150 seedlings from at least three replicate experiments. High sds indicate randomization of hypocotyl growth. The genotypes are as described in "Materials and Methods." The approximate relative levels of expressed *phyB* are: Nossen, 1; Z15, 15; ABO, 30; and D2, 60.

Genotype	Treatment	SD
Nossen	Dark	24.5
Nossen	Continuous R	52.3
Nossen	Continuous FR	25.4
Z15	Dark	18.57
Z15	Continuous R	65.6
Z15	Continuous FR	22.1
ABO	Dark	19.3
ABO	Continuous R	71.9
ABO	Continuous FR	40.3
D2	Dark	19.8
D2	Continuous R	55.8
D2	Continuous FR	48.2

**Table IV.** Effect of dark, continuous R, and continuous FR on growth orientation of wild-type and transgenic lines expressing different levels of *phyA*

Seedlings and data were handled as described in "Materials and Methods." The orientation of hypocotyl angle was measured as the SD around the vertical 0°. Values represent the pooled standard deviation for a minimum of 150 seedlings from at least three replicate experiments. High SDs indicate randomization of hypocotyl growth. The genotypes are as described in "Materials and Methods." The levels of expressed *phyA* are Nossen < 13K7 < 21K15.

Genotype	Treatment	SD
Nossen	Dark	24.5
Nossen	Continuous R	52.3
Nossen	Continuous FR	25.4
13K7	Dark	44.5
13K7	Continuous R	61.6
13K7	Continuous FR	51
21K15	Dark	67.4
21K15	Continuous R	62.8
21K15	Continuous FR	57.1

treatment with W to stimulate germination. When this was replaced by treatment with GA (Table V), dark randomization was not seen, suggesting that the randomization observed in the dark treatments was due to a *phyA*-mediated VLFR induced by the W pretreatment. This conclusion is supported by studies of the mutants, particularly the RLD ecotype in which the absence of *phyA* in the single and double mutants is associated with less randomization in dark growth conditions (Table II) and may reflect the absence of a *phyA*-mediated VLFR stimulated by the W pretreatment. Dark randomization in lines overexpressing *PHYA* cDNAs is not analogous to that seen in the *phyB-1* mutant, which when pretreated with GA retains dark randomization (Table V). These data suggest that the predominant activity associated with the action of *phyA* in hypocotyl gravitropism is randomization induced by a VLFR.

When one interprets data from lines overexpressing cDNAs, an account must be taken of the possibility that an aberrant phenotype being displayed is due to ectopic expression of the cDNA. At present there are no examples of problems due to ectopic expression involving phytochrome genes, and there are several experiments that suggest that phytochromes behave normally when overexpressed. First, *phyA* expressed by a transgene in *Arabidopsis* maintains the ability to display a FR high-irradiance response and is light labile (Boylan and Quail, 1991). Second, *phyB* expressed by a transgene has been shown to be spectrally active in transgenic *Arabidopsis*, undergoing R:FR-dependent conformational changes and to be stable in the light (Wagner et al., 1991). Third, the dominant negative suppression of *Arabidopsis* photoresponses by mutant sequences of *PHYA* genes was not due to aberrant expression patterns for either the introduced or endogenous *Arabidopsis* phytochromes (Boylan et al., 1994). Other possible mechanisms, such as co-suppression and competition for a limiting amount of chromophore, were also effectively discounted.

In this study overexpressed *phyB* remains responsive to R:FR ratio in regulating the gravitropic response of hypocotyls. Such a response to R:FR ratio is characteristic of the

action of *phyB*. Overexpression of the *PHYB* cDNA does not cause negative gravitropism in the dark, which means that overexpressed PrB is not affecting the response seen in the WT. Additionally, Z15 behaves similarly to WT in all experiments despite the fact that the *PHYB* cDNA will be expressed ectopically in this line. Consequently, the effects of overexpressing the *PHYB* cDNA in the lines ABO and D2 are most likely due to the level of expression rather than its ectopic nature.

### Interpretation

The data presented here clearly demonstrate that either *phyA* or *phyB* is capable of causing randomization of hypocotyl growth orientation. The data in Tables I and II concerning mutants demonstrate that in the absence of *phyA* R causes randomization through the action of *phyB*, which would be classically interpreted as being due to the conversion of PrB to PfrB. In the *phyA* mutants of both ecotypes the degree of randomization in continuous R is similar to that in the corresponding WT, indicating that *phyB* is able to substitute for the action of *phyA*. It is interesting that the converse is not true in the RLD ecotype, in which less randomization is seen in continuous R in the absence of *phyB*, suggesting that *phyA* cannot fully substitute for *phyB*.

Overproduction of *phyB* leads to increased sensitivity to R in the randomization of growth orientation; furthermore, growth orientation is a function of the photoequilibrium established by exposure to mixtures of continuous R and FR. Liscum and Hangarter (1993) reached the conclusion, logically correct on the basis of the data available to them, that the Pr form of *phyB* imposed negative gravitropism and that it was the removal of PrB by light, rather than the production of PfrB, that led to randomization. This conclusion was strongly dependent on the finding that the *hy2* mutant, which presumably contains only phytochrome apoproteins, is negatively gravitropic in both light and dark. According to the Liscum and Hangarter hypothesis, it is proposed that the PhyB apoprotein has the action putatively assigned to PrB in imposing negative gravitropism. Our data show that negative gravitropism occurs in the absence of *phyB* and therefore does not require PrB. Moreover, the activity associated with the action of *phyB* as shown by the mutants and overexpressing lines is randomization of hypocotyl orientation.

**Table V.** Hypocotyl orientation of *phyB-1*, *phyA* overexpresser, and wild type when induced to germinate using GA and grown in complete darkness

Seedlings and data were handled as described in "Materials and Methods." The orientation of hypocotyl angle was measured as the SD around the vertical 0°. Values represent the pooled SD for a minimum of 150 seedlings from at least three replicate experiments. High SDs indicate randomization of hypocotyl growth.

Genotype	SD
Landsberg <i>erecta</i>	14.8
<i>phyB-1</i>	65.3
Nossen	10.2
21K15	11.3

The highly randomized growth seen in the *phyB-1* allele in Landsberg *erecta* is not consistent with at least two other *phyB* alleles; *phyB-6* in RLD and *phyB-7* in Landsberg *erecta*. Consequently, it seems likely that *phyB-1* contains a second-site mutation that affects negative gravitropism. When the *phyB-1* mutant was germinated via a GA pretreatment in the complete absence of light, randomization persisted, confirming that this action was not associated with the action of phytochrome.

Because of the presence of a second-site mutation in the *phyB-1* mutant that results in high levels of randomization, it was important to determine the action of phyA in the absence of this mutation. In the RLD ecotype, phyA is clearly able to induce randomization of hypocotyl orientation in continuous R in the absence of phyB. In the phyA overproducers, dark randomization is seen, which is greater in the higher level expressers but is absent when the W pretreatment is replaced by GA pretreatment, indicating that "dark randomization" is actually a VLFR.

In conclusion, therefore, we propose that negative gravitropism is the default situation in *Arabidopsis* hypocotyls and that either PfrA or PfrB, produced by photoconversion of PrA or PrB, can cause randomization. No other phytochrome is apparently capable of this action. In this respect, the induction of randomization is a normal, phytochrome-mediated response, which, like most others in *Arabidopsis*, requires continuous irradiation for maximum effect. There is no requirement to invoke action of the Pr form of either phyA or phyB in this response. The dark randomization of hypocotyl orientation observed in dark treatments in the *phyB-1* mutant and in lines overexpressing *PHYA* cDNAs were induced by different mechanisms. In the former, randomization is probably due to a second-site mutation; in the latter, randomization is due to a phyA-mediated VLFR.

It is now possible to speculate that the function of phyA and phyB in the developing seedling with respect to gravitropism is to switch off negative gravitropism in the presence of light, thereby allowing other phototropic stimuli to determine the orientation of growth. The randomized state may simply occur because of the lack of phototropic stimuli to replace gravitropism, i.e. as a consequence of using monochromatic light, rather than represent a desired response induced in a developing seedling.

#### ACKNOWLEDGMENTS

The authors are grateful to Anne Irvine for excellent technical assistance. We thank Peter Quail, Margaret Boylan, and Doris Wagner for the RLD mutants and transgenic lines and Garry Whitelam for the *phyA* and *phyA/phyB* mutants in Landsberg *erecta*. We thank Professor Eberhard Schäfer and Dr. Roger Hangarter for helpful discussions concerning the possibility of a second-site mutation in the Landsberg *erecta phyB-1* strain of *A. thaliana*.

Received April 13, 1995; accepted October 6, 1995.

Copyright Clearance Center: 0032-0889/96/110/0211/06.

#### LITERATURE CITED

Boylan M, Douglas N, Quail PH (1994) Dominant negative suppression of *Arabidopsis* photoreponses by mutant phytochrome

A sequences identifies spatially discrete regulatory domains in the photoreceptor. *Plant Cell* 6: 449-460

Boylan MT, Quail PH (1991) Phytochrome A overexpression inhibits hypocotyl elongation in transgenic *Arabidopsis*. *Proc Natl Acad Sci USA* 88: 10806-10810

Goto N, Kumagai T, Koornneef M (1991) Flowering responses to light-breaks in photomorphogenic mutants of *Arabidopsis thaliana*, a long-day plant. *Physiol Plant* 83: 209-215

Johnson E, Bradley M, Harberd NP, Whitelam GC (1994) Photoresponses of light-grown phyA mutants of *Arabidopsis*. Phytochrome A is required for the perception of daylength extensions. *Plant Physiol* 105: 141-149

Liscum E, Hangarter RP (1993) Genetic evidence that the red-absorbing form of phytochrome B modulates gravitropism in *Arabidopsis thaliana*. *Plant Physiol* 103: 15-19

Nagatani A, Reed JW, Chory J (1993) Isolation and initial characterization of *Arabidopsis* mutants that are deficient in phytochrome A. *Plant Physiol* 102: 269-277

Neff MM, Van Volkenburgh E (1994) Light-stimulated cotyledon expansion in *Arabidopsis* seedlings. *Plant Physiol* 104: 1027-1032

Parks BM, Jones AM, Adamse P, Koornneef M, Kendrick RE, Quail PH (1987) The *aurea* mutant of tomato is deficient in spectrophotometrically and immunologically detectable phytochrome. *Plant Mol Biol* 9: 97-107

Parks BM, Quail PH (1991) Phytochrome-deficient *hy1* and *hy2* long hypocotyl mutants of *Arabidopsis* are defective in phytochrome chromophore biosynthesis. *Plant Cell* 3: 1177-1186

Parks BM, Quail PH (1993) *hy8*, a new class of *Arabidopsis* long hypocotyl mutants deficient in functional phytochrome A. *Plant Cell* 5: 39-48

Quail PH, Briggs WR, Chory J, Hangarter RP, Harberd NP, Kendrick RE, Koornneef M, Parks B, Sharrock RA, Schafer E, Thompson WE, Whitelam GC (1994) Spotlight on phytochrome nomenclature. *Plant Cell* 6: 468-471

Reed JW, Nagpal P, Poole DS, Furuya M, Chory J (1993) Mutations in the gene for the red/far-red light receptor phytochrome B alter cell elongation and physiological responses throughout *Arabidopsis* development. *Plant Cell* 5: 147-157

Robson PRH, Whitelam GC, Smith H (1993) Selected components of the shade-avoidance syndrome are displayed in a normal manner in mutants of *Arabidopsis thaliana* and *Brassica rapa* deficient in phytochrome B. *Plant Physiol* 102: 1179-1184

Shinomura T, Nagatani A, Chory J, Furuya M (1994) The induction of seed germination in *Arabidopsis thaliana* is regulated principally by phytochrome B and secondarily by phytochrome A. *Plant Physiol* 104: 363-371

Smith H (1994a) Sensing the light environment: the functions of the phytochrome family. In RE Kendrick, GHM Kronenberg, eds, *Photomorphogenesis in Plants*, Ed 2. Kluwer Academic, Dordrecht, The Netherlands, pp 377-416

Smith H (1994b) Phytochrome transgenics: functional, ecological and biotechnological applications. *Semin Cell Biol* 5: 315-325

Smith H (1995) Physiological and ecological function within the phytochrome family. *Annu Rev Plant Physiol Plant Mol Biol* 46: 289-315

Smith H, Whitelam GC (1990) Phytochrome, a family of photoreceptors with multiple physiological roles. *Plant Cell Environ* 13: 695-707

Stanier RY, Kunisawa R, Mandel M, Cohen-Bazire G (1971) Purification and properties of unicellular blue-green algae (order Chroococcales). *Bacteriol Rev* 35: 171-205

Wagner D, Tepperman JM, Quail PH (1991) Overexpression of phytochrome B induces a short hypocotyl phenotype in transgenic *Arabidopsis*. *Plant Cell* 3: 1275-1288

Whitelam GC, Johnson E, Peng J, Carol P, Anderson ML, Cowl JS, Harberd NP (1993) Phytochrome A null mutants of *Arabidopsis* display a wild-type phenotype in white light. *Plant Cell* 5: 757-768

Whitelam GC, Smith H (1991) Retention of phytochrome-mediated shade avoidance responses in phytochrome-deficient mutants of *Arabidopsis* cucumber and tomato. *J Plant Physiol* 139: 119-125