

Second Messengers Mediate Increases in Cytosolic Calcium in Tobacco Protoplasts

Igor D. Volotovski, Sergei G. Sokolovsky, Olga V. Molchan, and Marc R. Knight^{1*}

Institute of Photobiology, Academy of Sciences of Belarus, Academicheskaja Street 27, 220072 Minsk, Belarus (I.D.V., S.G.S., O.V.M.); and Department of Plant Sciences, University of Oxford, South Parks Road, Oxford OX1 3RB, United Kingdom (M.R.K.)

Addition of membrane-permeable cyclic GMP (cGMP) and cyclic AMP (cAMP) were shown to cause elevation of cytosolic Ca²⁺ concentration ([Ca²⁺]_{cyt}) in tobacco (*Nicotiana plumbaginifolia*) protoplasts. Under the same conditions these cyclic nucleotides were shown to provoke a physiological swelling response in the protoplasts. Nonmembrane-permeable cAMP and cGMP were unable to trigger a detectable [Ca²⁺]_{cyt} response. Cyclic-nucleotide-mediated elevations in [Ca²⁺]_{cyt} involved both internal and external Ca²⁺ stores. Both cAMP- and cGMP-mediated [Ca²⁺]_{cyt} elevations could be inhibited by the Ca²⁺-channel blocker verapamil. Addition of inhibitors of phosphodiesterases (isobutylmethylxanthine and zaprinast) and the adenylate cyclase agonist forskolin to the protoplasts (predicted to elevate in vivo cyclic-nucleotide concentrations) caused elevations in [Ca²⁺]_{cyt}. Addition of the adenylate cyclase inhibitor 2',5'-dideoxyadenosine before forskolin significantly inhibited the forskolin-induced [Ca²⁺]_{cyt} elevation. Taken together, these data suggest that a potential communication point for cross-talk between signal transduction pathways using cyclic nucleotides in plants is at the level of Ca²⁺ signaling.

Environmental and hormonal signals regulate various physiological processes in plants via signal transduction pathways. Essential components of signal transduction pathways include second messengers. Vital roles for the second messengers Ca²⁺, cAMP, cGMP, IP₃, and 1,2-diacylglycerol were first discovered in animal systems (Berridge and Irvine, 1984; Nishizuka, 1984). The last decade has been marked by substantial progress in elucidating the mechanisms of intracellular signaling in plants (Gilroy and Trewavas, 1994). Attempts to draw comparisons with mammalian systems have been successful in showing that Ca²⁺-calmodulin, cGMP (Neuhaus et al., 1993), and phosphoinositide signal mechanisms (Alexandre et al., 1990; Allen et al., 1995) also exist in plants, although cAMP involvement in transduction pathways has remained controversial (Assmann, 1995; Bolwell, 1995).

It is known that a number of chemical and physical stimuli mediate their effects via transient increases in the concentration of intracellular free Ca²⁺ (Gilroy and Trewavas, 1994). Ca²⁺ influx into the cytosol can occur because of the opening of Ca²⁺ channels in plasma membrane, with

Ca²⁺ entering the cell down a concentration gradient. Receptor activation, at least in mammalian cells, can trigger the phosphoinositide cascade, which leads to the production of IP₃ and 1,2-diacylglycerol (Berridge and Irvine, 1984). IP₃ can then provoke the release from the internal Ca²⁺ stores of plants cells by opening intracellular Ca²⁺ channels (Alexandre et al., 1990; Gilroy et al., 1990; Allen et al., 1995). Increases in [Ca²⁺]_{cyt} produced either by influx or release from internal Ca²⁺ stores stimulate the phosphorylation of proteins within the cell (Poovaiah and Reddy, 1990). An increase in [Ca²⁺]_{cyt} detected by Ca²⁺ microelectrodes, Ca²⁺-sensitive fluorescence probes, and extrinsic and intrinsic aequorin (Cobbold and Rink, 1987; Chae et al., 1990; Knight et al., 1991; Shacklock et al., 1992), followed by the alteration of the activities of intercellular targets, enzymes, genes, pumps, and channels, have all been demonstrated in plants (Fallon et al., 1993; Neuhaus et al., 1993; Bowler et al., 1994a). In some circumstances activated plant cells can display repeated Ca²⁺ spikes or oscillations (McAinsh et al., 1995) and spreading Ca²⁺ waves (Shacklock et al., 1992; Campbell et al., 1996). Such elevations in [Ca²⁺]_{cyt} have been shown to be involved in several signaling pathways triggered, for example, by light via phytochrome (Neuhaus et al., 1993; Bowler et al., 1994a, 1994b), by phytohormones (Felle, 1988; Gehring et al., 1990), and even by mechanical signals and cold shock (Knight et al., 1991, 1992; Haley et al., 1995; Campbell et al., 1996). The multifunctional role of Ca²⁺ suggests that it participates in many signaling pathways in the plant cell. A simple but effective method for measuring changes in [Ca²⁺]_{cyt} in whole plants has been developed (Knight et al., 1991). This involves the expression of the Ca²⁺-activated photoprotein aequorin in transgenic plants (Knight et al., 1993; Knight and Knight, 1995). This approach can be adapted for use in protoplasts that provide a superlative system for calibration of the Ca²⁺ signal (Haley et al., 1995; Knight et al., 1996).

The data presented in this paper show that increases in intracellular concentrations of the second messengers cGMP and cAMP (Tepper et al., 1995) provoke increases in

Abbreviations: 8-Br-cAMP, 8-brom-cyclic AMP; 8-Br-cGMP, 8-brom-cyclic GMP; [Ca²⁺]_{cyt}, cytosolic Ca²⁺ concentration; db-cAMP, 2'-O-dibutyryl AMP; db-cGMP, 2'-O-dibutyryl GMP; DDOA, 2',5'-dideoxyadenosine; IBMX, 3-isobutyl-1-methylxanthine; IP₃, inositol 1,4,5-trisphosphate.

¹ M.R.K. is a Royal Society University Research Fellow.

* Corresponding author; e-mail marc.knight@plants.ox.ac.uk; fax 44-1865-275023.

$[Ca^{2+}]_{cyt}$ in plant cells. This suggests that a potential communication point for cross-talk between signal transduction pathways using these second messengers (Bowler et al., 1994b) is the release of intracellular and extracellular Ca^{2+} stores.

MATERIALS AND METHODS

Plant Material and Protoplast Isolation

Tobacco (*Nicotiana plumbaginifolia*) genetically transformed to express apoaequorin under the control of the constitutive 35S promoter of cauliflower mosaic virus was used in this study (transgenic line MAQ2.4; Knight et al., 1991). Protoplasts were isolated from the leaves of 8-week-old plants grown in white light ($150 \mu\text{mol m}^{-2} \text{s}^{-1}$) for 15 h/d at 20°C according to standard methods by Shillito and Potrykus (1985), with slight modifications. The epidermal layer was stripped from the leaves and they were incubated for 3.5 to 5.5 h with regular shaking (30 min) in 20 mM Mes (Sigma) buffer solution adjusted to pH 5.5 and containing 2% (w/v) cellulase Onozuka R-10 (Serva, Heidelberg, Germany), 400 mM mannitol (Chemapol, Bratislava, Slovakia), 100 mM Gly (Merck, Darmstadt, Germany), and 5 mM $CaCl_2$. After incubation, the protoplast suspension was filtered through a nylon filter with a $100\text{-}\mu\text{m}$ pore size and the protoplasts were washed several times in 20 mM HEPES (Sigma), pH 7.0, 500 mM mannitol, 2 mM $MgCl_2$, and 0.1 mM EGTA (Sigma), followed by centrifugation at $200g$ for 3 min. After isolation, the protoplast concentration was adjusted to 2×10^5 cells mL^{-1} in the same solution that was used for washing. Protoplasts were placed in this medium for all in vivo Ca^{2+} measurements. Protoplast viability was tested according to the procedure of Rudenok et al. (1973) with the exclusion dye methylene blue (0.3 mM, Sigma).

Aequorin Reconstitution and $[Ca^{2+}]_{cyt}$ Monitoring

For in vivo aequorin reconstitution (Knight and Knight, 1995), the protoplasts were incubated for 4 h at room temperature with $2 \mu\text{M}$ coelentraxine (Molecular Probes, Eugene, OR) diluted in methanol (Merck). After washing, chemiluminescence measurements were performed with a digital chemiluminometer (model PCHL-01, Biopribor, Moscow, Russia) equipped with a PEU-84 photomultiplier and chart recorder. The total sample volume in the luminometer cuvettes was 0.5 mL. All additions (usually not more than 0.2 mL) were added to the cuvette using a micropipette with a thin plastic tube. At the end of each measurement protoplasts were discharged by mixing with an equal volume of 100 mM $CaCl_2/0.1\%$ (v/v) Triton X-100. The peak value of the stimuli-induced $[Ca^{2+}]_{cyt}$ transient was calculated as described by Cobbold and Rink (1987), with some modifications, to take into account the specific isoform used and the experimental temperature (Knight et al., 1996).




Figure 1. Influence of db-cGMP (A) and db-cAMP (B) on Ca^{2+} -dependent chemiluminescence of aequorin in protoplasts isolated from transgenic tobacco leaves (traces from chart recorder). Applications of stimuli are shown by arrows. Aliquots ($200 \mu\text{L}$) of cyclic mononucleotides diluted in buffer to a final concentration of $10 \mu\text{M}$, were added to the suspension (0.5 mL) of protoplasts (2×10^5 cells mL^{-1}) in 0.5 M mannitol, 20 mM HEPES, pH 7.0, 2 mM $MgCl_2$, and 0.1 mM EGTA. Before the addition of cyclic mononucleotides, $CaCl_2$ solution ($200 \mu\text{L}$) was added to give a final $CaCl_2$ concentration of 1 mM (shown by arrows). C, Peak $[Ca^{2+}]_{cyt}$ values after the addition of cyclic mononucleotides and Ca^{2+} to protoplasts were calculated after chemiluminescence discharge under the action of 0.1% Triton X-100/ 100 mM $CaCl_2$, as described previously (Cobbold and Rink, 1987; Knight and Knight, 1995). Experiments were performed seven times; the error bars on histograms indicate SD.




Figure 2. Influence of db-cGMP (A), db-cAMP (B), 8-Br-cGMP (C), and 8-Br-cAMP (D) on Ca²⁺-dependent chemiluminescence of aequorin in protoplasts isolated from transgenic tobacco leaves and suspended in Ca²⁺-deficient medium (traces from chart recorder). The conditions of the experiments were the same as in Figure 1. E, Peak $[Ca^{2+}]_{cyt}$ values after the addition of cyclic trinucleotides and Ca²⁺ to protoplasts were calculated after chemiluminescence discharge under the action of 0.1% Triton X-100/100 mM CaCl₂ as described previously (Cobbold and Rink, 1987; Knight and Knight, 1995). Experiments were performed seven times; the error bars on histograms indicate sd.

Protoplast Swelling

Protoplast swelling was used as a simple indicator of plant cell physiological response (Kim et al., 1986; Bossen et al., 1988; Shacklock et al., 1992). After cAMP/cGMP addition, protoplasts (5×10^5 in 0.5 mL) were kept in the dark for different time intervals at 22°C. Then, protoplasts were

placed on a hemocytometer and photographed. The diameters of 100 protoplasts for each incubation time were determined and the mean volume was evaluated, making the assumption that the cells were spherical. All measurements of the changes in protoplast volume were made in the same medium that was used for Ca²⁺ measurements.





Figure 3. Dose dependence of $[Ca^{2+}]_{cyt}$ elevation in protoplasts to db-cGMP (●) and db-cAMP (○) concentration without any external Ca²⁺.

RESULTS

All *in vivo* $[Ca^{2+}]_{cyt}$ measurement experiments were repeated at least three times. The traces presented were taken from the replicates and represent luminescence over time. The size of the luminescence peak and the luminescence kinetics are governed not only by the rate and concentration of $[Ca^{2+}]_{cyt}$, but also by the absolute amount of aequorin present in the cells. Higher or lower amounts of aequorin will give larger or smaller peaks in response to the same changes in Ca²⁺. If the consumption of aequorin becomes significant, then a smaller-than-expected peak is obtained. All of these factors are taken into account when the peak $[Ca^{2+}]_{cyt}$ values are calibrated. Thus, these calibrated $[Ca^{2+}]_{cyt}$ values may be compared between treatments in the same experiment. For the reasons outlined above, however, no qualitative comparisons can be made

Figure 4. Effect of IBMX (A), zaprinast (B), forskolin (C), and DDOA plus forskolin (D) on $[Ca^{2+}]_{cyt}$ in tobacco protoplasts as measured by aequorin chemiluminescence. The compounds were added to the suspension of tobacco protoplasts at the indicated concentrations. The composition of the buffer with no Ca^{2+} was as described for Figure 1. E, The histograms show the peak $[Ca^{2+}]_{cyt}$ values obtained for each treatment. The error bars on the histograms indicate SD.



between experiments or treatments based on magnitude and kinetics of luminescence alone. Addition of 2 mM extracellular $CaCl_2$ to the protoplasts (Fig. 1, A and B) provoked increases of $[Ca^{2+}]_{cyt}$ of about 0.2 to 0.3 μM , indicating successful *in vivo* reconstitution of aequorin. Regardless of whether this extracellular Ca^{2+} was added before or after second messengers, its addition produced maximum $[Ca^{2+}]_{cyt}$ values that were quite similar (data not shown).

The effects of db-cGMP and db-cAMP on $[Ca^{2+}]_{cyt}$ levels of tobacco protoplasts were tested. As can be seen in Figure 1, A and B, the addition of either of these second messengers at 10 μM provoked large transient increases in $[Ca^{2+}]_{cyt}$. db-cGMP caused a transient $[Ca^{2+}]_{cyt}$ elevation of around 0.8 μM , lasting for approximately 40 s (Fig. 1, A and C). db-cAMP caused a transient $[Ca^{2+}]_{cyt}$ elevation of around 0.7 μM , lasting for approximately 100 s (Fig. 1C). Figure 2, C and D, demonstrates the effect of 8-Br-cGMP

and 8-Br-cAMP. These experiments were performed on protoplasts suspended in Ca^{2+} -deficient medium to compare the results obtained in Figure 1 (in medium containing Ca^{2+}) and hence gauge the involvement of external/internal Ca^{2+} stores in the $[Ca^{2+}]_{cyt}$ responses to cAMP/cGMP. Both types of cyclic-nucleotide derivatives caused similar transient $[Ca^{2+}]_{cyt}$ elevations, unequivocally indicating that the mononucleotide parts of the molecules were responsible. The transients also lasted about 40 and 100 s, respectively. Both db-cGMP and db-cAMP were able to produce a $[Ca^{2+}]_{cyt}$ elevation in Ca^{2+} -free medium (Fig. 2, A and B), strongly suggesting Ca^{2+} release from internal stores of ion. There was substantial variability in the calibrate cyclic-nucleotide-mediated $[Ca^{2+}]_{cyt}$ peak values between experiments performed under the same conditions (compare Figs. 2 and 3). Also, the apparent relative potency of the cAMP and cGMP varied (compare Figs. 2 and 3). This is most likely because of physiological differences between protoplasts from different preparations leading to differences in signaling. Practically, this means that the relative quantitative contributions of intracellular/extracellular stores cannot be gauged. Qualitatively, it is clear from our results that both types of Ca^{2+} stores are involved, although the 8-Br- derivatives were less potent than the db- derivatives at concentrations that mediated a $[Ca^{2+}]_{cyt}$ increase (Fig. 2). This difference might be attributable to nuances in uptake efficiencies of the two derivative forms of cAMP/cGMP.

Figure 3 shows a dose-response curve for cyclic-nucleotide concentration plotted against peak $[Ca^{2+}]_{cyt}$ values as a result of intracellular Ca^{2+} release (this experiment was performed in the absence of external Ca^{2+}). The concentration dependence reached a plateau somewhere between 1 and 10 μM for both cyclic mononucleotides (Fig. 3). The ability of cyclic nucleotides to cause $[Ca^{2+}]_{cyt}$ elevations was further demonstrated by the use of membrane-permeable modulators of anabolic and catabolic pathways




Figure 5. Dose dependence of $[Ca^{2+}]_{cyt}$ elevation in tobacco protoplasts for forskolin concentration under the same conditions as described for Figure 1.




Figure 6. Effects of buffer alone and water-soluble cAMP and cGMP on $[Ca^{2+}]_{cyt}$ levels in tobacco protoplasts. Media for cAMP, cGMP, and buffer plus ethanol (200- μ L solutions) were the same as for Figure 1. Peak $[Ca^{2+}]_{cyt}$ values after the addition of cyclic mononucleotides and Ca^{2+} to protoplasts were calculated after chemiluminescence discharge under the action of 0.1% Triton X-100/100 mM $CaCl_2$ as described previously (Cobbold and Rink, 1987; Knight and Knight, 1995).

of cyclic-mononucleotide metabolism. $[Ca^{2+}]_{cyt}$ elevations were observed with the addition of the cyclic-mononucleotide phosphodiesterase inhibitor IBMX (Fig. 4A), the cGMP phosphodiesterase inhibitor zaprinast (Fig. 4B), and the adenylyl cyclase activator forskolin (Fig. 4C). When the adenylyl cyclase inhibitor DDOA was added before forskolin, the forskolin-mediated $[Ca^{2+}]_{cyt}$ increase was inhibited (Fig. 4D). It seems that Ca^{2+} -sensitive chemiluminescence of cytosolic aequorin can be induced by the changes in the endogenous cyclic-mononucleotide concentrations, stimulating their formation or inhibiting their hydrolysis.

Figure 5 shows the correlation between the concentration of forskolin added and the $[Ca^{2+}]_{cyt}$ responses obtained. The most pronounced effect on $[Ca^{2+}]_{cyt}$ was observed at 20 μ M, a concentration that is consistent with the concentration range of optimal activity of forskolin in animal cells (Siamon et al., 1981). Higher concentrations of forskolin seem to have inhibitory effects (Fig. 5), most likely because of cell toxicity at these concentrations.

As described previously (Haley et al., 1995), additions to protoplasts in solution may result in the so-called "touch" effect, a consequence of mechanical induction of Ca^{2+} release from intracellular Ca^{2+} stores. To minimize the touch effect, Ca^{2+} solution was added to protoplast suspensions very carefully. This was verified by testing the effect of adding membrane-impermeable cAMP and cGMP to pro-

toplasts as well as by the addition of buffer alone, including ethanol (as a pharmaceutical diluent), as controls (Fig. 6). The addition of cAMP, cGMP, or buffer alone immediately produced very rapid and small transients of cytosolic free Ca^{2+} attributable to the touch response. As can be seen in Figures 1 and 2, a rapid transient is also seen immediately upon addition of db-cGMP or db-cAMP, which can be attributed to the same phenomenon. This is clearly separate in time and magnitude from the very large and prolonged responses seen with these membrane-permeable second messengers.

Figure 7 shows the effect of the Ca^{2+} -channel blocker verapamil on cyclic-nucleotide-mediated $[Ca^{2+}]_{cyt}$ increases. Verapamil inhibited the effects of db-cAMP and db-cGMP, producing lower elevations of $[Ca^{2+}]_{cyt}$. The inhibiting action of verapamil on db-cAMP/db-GMP-induced $[Ca^{2+}]_{cyt}$ elevation was more pronounced for cGMP than for cAMP (Fig. 7C).

To understand the signaling role of Ca^{2+} in plants in relation to the involvement of cyclic nucleotides as the messengers, the physiological action of protoplast swelling was investigated. Figure 8 shows that the time course of protoplast swelling was caused by db-cAMP and db-cGMP. According to our results, a 15-min incubation was sufficient for the curves to reach a plateau corresponding to about a 10% and 15% increase in volume for db-cGMP and db-cAMP, respectively.




Figure 7. Inhibition of db-cGMP-induced (A) and db-cAMP-induced (B) $[Ca^{2+}]_{cyt}$ elevation by verapamil in Ca^{2+} -deficient medium. C, Percentage of inhibition by verapamil. Experiments were performed three times; the error bars on histograms indicate SD.

DISCUSSION

The data presented in this paper show that second messengers such as cGMP and cAMP can provoke $[Ca^{2+}]_{cyt}$ elevations in tobacco protoplasts. In the case of cAMP and cGMP, the targets of action are likely to be intracellular, because even though membrane-permeable analogs at concentrations as low as 0.1 μ M provoked a large $[Ca^{2+}]_{cyt}$ response (Fig. 1, A and B), membrane-impermeable cAMP and cGMP did not produce any significant effect (Fig. 6). An intracellular site of action is also supported by the fact that a significant $[Ca^{2+}]_{cyt}$ response was induced by db-

cGMP and db-cAMP in the absence of external Ca^{2+} . Protoplasts in EGTA (zero external Ca^{2+}) still showed $[Ca^{2+}]_{cyt}$ responses to these second messengers, implicating the release of Ca^{2+} from internal stores as a mechanism for this response. Although variability was seen between protoplast preparations, cyclic-nucleotide-mediated $[Ca^{2+}]_{cyt}$ increases were generally higher in the presence of external Ca^{2+} . This suggests that cyclic mononucleotides may also trigger changes in the Ca^{2+} permeability of the plasma membrane and thus cause a Ca^{2+} influx. The results obtained with verapamil (Fig. 7) indicate that a significant proportion of the $[Ca^{2+}]_{cyt}$ responses occurs via the activation of verapamil-sensitive Ca^{2+} channels.

Similar $[Ca^{2+}]_{cyt}$ elevations were also mediated by the addition of 8-Br-cAMP and 8-Br-cGMP, which also penetrate membranes very easily (Fig. 2, C and D). The $[Ca^{2+}]_{cyt}$ responses stimulated by these two types of derivative of membrane-permeable cyclic nucleotides were relatively large, much greater in magnitude, and much more prolonged than the touch $[Ca^{2+}]_{cyt}$ response caused by the mechanical stimulation resulting from the addition of solutions to the protoplasts (Fig. 6).

$[Ca^{2+}]_{cyt}$ elevation in protoplast cytoplasm was also induced by manipulating endogenous cyclic-mononucleotide levels. The addition of various modulators controlling cyclic-nucleotide metabolism to protoplasts produced significant changes in $[Ca^{2+}]_{cyt}$. As can be seen in Figures 4 and 5, compounds that should increase endogenous cyclic-nucleotide concentration (forskolin, IBMX, and zaprinast) produced increases in $[Ca^{2+}]_{cyt}$. The interesting result that is consistent with the above explanation was obtained when DDOA was used before forskolin. Forskolin stimulates adenylate cyclase within the cell and, consequently, increases the concentration of endogenous cAMP. It seems likely that




Figure 8. Time course of tobacco protoplast swelling after treatment with db-cGMP and db-cAMP in complete darkness at 22°C, expressed as percent volume change ($\Delta V, \%$). Protoplasts were suspended in the media described for Figure 1. Cyclic mononucleotides were added at a concentration of 10 μ M and incubation times varied as shown. The volume changes were determined as described in "Materials and Methods". The percent volume change is relative to the control without cyclic mononucleotides. Each point represents the mean of at least three separate experiments; the error bars indicate SE.

DDOA competitively inhibits the adenylate cyclase and prevents its stimulation by forskolin (Fig. 4, C and D).

In addition to [Ca²⁺]_{cyt} elevation induced by db-cAMP and db-cGMP, the physiological response of protoplast swelling was observed at the same concentrations of cyclic mononucleotides (Fig. 8). These data are consistent with those obtained by other authors (Kim et al., 1986; Bossen et al., 1988; Gilroy and Trewavas, 1994). It has also been shown that protoplast swelling is under the control of red light absorbed by phytochrome (Bossen et al., 1988; Chung et al., 1988; Shacklock et al., 1992). It has been shown that the release of caged Ca²⁺ can cause protoplast swelling in the absence of light (Shacklock et al., 1992). Therefore, it is possible that cyclic-nucleotide-induced increases in [Ca²⁺]_{cyt} mediate protoplast swelling. Furthermore, it is possible that the phenomenon described in this paper is relevant to the photophysiology of phytochrome action, i.e. signal transmission from the photoreceptor to target elements within the cell (Barnes et al., 1997), where cyclic mononucleotides have been shown to be downstream signaling molecules (Bowler et al., 1994b).

A key area that needs to be understood concerns the mechanism by which these second messengers bring about Ca²⁺ release from intracellular Ca²⁺ stores into cytoplasm. It could be indirect via effects on protein kinases (Stone and Walker, 1995), or possibly by direct action on ion channels (Ward et al., 1995). The first possibility is very important because the activities of many plant cell proteins are modulated by phosphorylation (Ranjewa and Boudet, 1987). The activities of cGMP- and cAMP-dependent protein kinases have been detected in animals (Krebs, 1985). There is also some evidence for cyclic-mononucleotide-regulated kinases in plant cells (Komatsu and Hirano, 1993). However, unlike the situation in mammalian cells, the involvement of cAMP still needs to be unequivocally elucidated in plants (Bolwell, 1995). Generally, the problem is that the low levels of cAMP present in plant cells means that cAMP-dependent processes that are detected experimentally are viewed with suspicion. However, analogs and modulators of cAMP metabolism have been shown to have physiological effects in plant cells (Assmann, 1995). The data presented here suggest the possibility of a role for cAMP in the control of Ca²⁺ release from intracellular stores in plant cells. The second potential targets for cyclic mononucleotides, at least in animals, are ion channels, which can be directly controlled by cyclic mononucleotides, e.g. in visual and olfactory cells in animals (Fesenko et al., 1985; Delgado et al., 1991). Cyclic mononucleotides in these cases have been shown to be bound directly to specific binding sites located at the components of so-called cGMP- and cAMP-dependent ion channels.

Environmental, chemical, and hormonal stimuli that are capable of inducing changes in [Ca²⁺]_{cyt} are characterized by individual Ca²⁺ signatures with different amplitudes, kinetics, and spatial distribution, which, it has been suggested, allow cells to distinguish between stimuli (Gilroy and Trewavas, 1994; Haley et al., 1995; Campbell et al., 1996). Thus, stimulus-induced changes in [Ca²⁺]_{cyt} may be transient, sustained, or oscillatory (Knight et al., 1991; McAinsh et al., 1995; Campbell et al., 1996). Typically, these

[Ca²⁺]_{cyt} responses can last from a few seconds to several hours. It is generally accepted that changes in [Ca²⁺]_{cyt} are closely associated with transduction of the biological action of auxin, ABA, cytokinins, gibberellic acid, fungal elicitors, and light (Gilroy and Trewavas, 1994; Bush, 1995). The involvement of second messengers other than Ca²⁺ in these pathways has not been demonstrated, although cGMP involvement in phytochrome-mediated *chs* gene expression has been shown (Bowler et al., 1994a, 1994b). cGMP was shown to substitute for the action of red light in the phytochrome-dependent regulatory control of anthocyanin synthesis (Neuhaus et al., 1993; Bowler et al., 1994a). It is clearly possible that cyclic nucleotides (Bolwell, 1995) and metabolites from phosphoinositide cycles are involved in many other plant signal transduction pathways (Alexandre et al., 1991; Allen et al., 1995). The major challenge for the future will be to determine whether cyclic-nucleotide-mediated [Ca²⁺]_{cyt} increases are actually used in bona fide signal transduction pathways in plants.

ACKNOWLEDGMENTS

We thank Prof. A.A. Milutin (Sakcharov's College, Minsk), who donated coelentraxine, and Dr. V.B. Gavrillov (Institute of Photobiology, Academy of Sciences of Belarus, Minsk) for the use of the chemiluminometer.

Received January 5, 1998; accepted April 17, 1998.

Copyright Clearance Center: 0032-0889/98/117/1023/08.

LITERATURE CITED

- Alexandre J, Lassalles JP, Kado RT (1990) Opening of Ca²⁺ channels in isolated red beet root vacuole membrane by inositol 1,4,5-trisphosphate. *Nature* **343**: 567-570
- Allen GJ, Muir SR, Sanders D (1995) Release of Ca²⁺ from individual plant vacuoles by both InsP₃ and cyclic ADP-ribose. *Science* **268**: 735-737
- Assmann SM (1995) Cyclic AMP as a second messenger in higher plants. *Plant Physiol* **108**: 885-889
- Barnes SA, McGrath RB, Chua N-H (1997) Light signal transduction in plants. *Trends Cell Biol* **7**: 21-26
- Berridge MJ, Irvine RF (1984) Inositol trisphosphate, a novel second messenger in cellular signal transduction. *Nature* **312**: 315-321
- Bolwell GP (1995) Cyclic AMP, the reluctant messenger in plants. *Trends Biochem Sci* **12**: 492-495
- Bossen ME, Dassen HHA, Kendrick RE, Vredenberg WJ (1988) The role of calcium ions in phytochrome-controlled swelling of etiolated wheat (*Triticum aestivum* L.) protoplasts. *Planta* **174**: 94-100
- Bowler C, Neuhaus G, Yamagata H, Chua N-H (1994a) Cyclic GMP and calcium mediate phytochrome phototransduction. *Cell* **77**: 73-81
- Bowler C, Yamagata H, Neuhaus G, Chua N-H (1994b) Phytochrome signal transduction pathways are regulated by reciprocal control mechanisms. *Genes Dev* **8**: 2188-2202
- Bush DS (1995) Calcium regulation in plant cells and its role in signalling. *Annu Rev Plant Physiol Plant Mol Biol* **46**: 95-122
- Campbell AK, Trewavas AJ, Knight MR (1996) Calcium imaging shows differential sensitivity to cooling and communication in luminous transgenic plants. *Cell Calcium* **19**: 211-218
- Chae Q, Park HJ, Hong SD (1990) Loading of quin2 into the oat protoplasts and measurement of cytosolic calcium ion concentration changes by phytochrome action. *Biochim Biophys Acta* **1051**: 115-122
- Chung CH, Lim S-V, Song P-S, Berlin JD, Gooding JR (1988) Swelling of etiolated oat protoplasts induced by phytochrome,

- cAMP, and gibberellic acid: a kinetic study. *Plant Cell Physiol* **29**: 855–860
- Cobbold PH, Rink TJ** (1987) Fluorescence and bioluminescence measurements of cytoplasmic free calcium. *Biochem J* **248**: 313–328
- Delgado R, Hidalgo P, Diaz F, Latorre R, Labarca P** (1991) A cyclic AMP-activated K⁺ channel in *Drosophila* larva muscle is persistently activated in dunce. *Proc Natl Acad Sci USA* **88**: 557–560
- Fallon KM, Shacklock PS, Trewavas AJ** (1993) Detection *in vivo* of very rapid red light-induced calcium-sensitive protein phosphorylation in etiolated wheat (*Triticum aestivum*) leaf protoplasts. *Plant Physiol* **101**: 1039–1045
- Felle H** (1988) Auxin causes oscillations of cytosolic free calcium and pH in *Zea mays* coleoptiles. *Planta* **174**: 495–499
- Fesenko EE, Kolesnikov SS, Lyubarsky AL** (1985) Induction by cyclic GMP of cationic conductance in plasma membrane of retinal rod outer segment. *Nature* **313**: 310–313
- Gehring CA, Irving HR, Parish RW** (1990) Effect of abscisic acid on cytosolic calcium and pH in plant cell. *Proc Natl Acad Sci USA* **87**: 9645–9649
- Gilroy S, Read ND, Trewavas AJ** (1990) Elevation of cytosolic calcium by caged inositol trisphosphate initiates stomatal closure. *Nature* **346**: 769–771
- Gilroy S, Trewavas AJ** (1994) A decade of plant signals. *BioEssays* **16**: 677–682
- Haley A, Russel AJ, Wood N, Allan AC, Knight MR, Campbell AK, Trewavas AJ** (1995) Effects of mechanical signalling on plant cell cytosolic calcium. *Proc Natl Acad Sci USA* **92**: 4124–4128
- Kim Y-S, Moon DK, Goodin JR, Sdong P-S** (1986) Swelling of etiolated protoplasts induced by cAMP and red light. *Plant Cell Physiol* **27**: 193–197
- Knight H, Knight MR** (1995) Recombinant aequorin methods for intracellular calcium measurement in plants. *Methods Cell Biol* **49**: 201–216
- Knight H, Trewavas AJ, Knight MR** (1996) Cold calcium signalling in Arabidopsis involves two cellular pools and a change in calcium signature after acclimation. *Plant Cell* **8**: 489–503
- Knight MR, Campbell AK, Smith SM, Trewavas AJ** (1991) Transgenic plant aequorin reports the effects of touch and cold-shock and elicitors on cytoplasmic calcium. *Nature* **352**: 524–526
- Knight MR, Read ND, Campbell AK, Trewavas AJ** (1993) Imaging calcium dynamics in living plants using semi-synthetic recombinant aequorin. *J Cell Biol* **121**: 83–90
- Knight MR, Smith SM, Trewavas AJ** (1992) Wind-induced plant motion immediately increases cytosolic calcium. *Proc Natl Acad Sci USA* **89**: 4967–4971
- Komatsu S, Hirano H** (1993) Protein kinase activity and phosphorylation in rice (*Oryza sativa* L.) leaf. *Plant Sci* **94**: 127–137
- Krebs EG** (1985) The phosphorylation of proteins: a major mechanism for biological regulation. *Biochem Soc Trans* **13**: 813–820
- McAinsh MR, Webb AR, Taylor JE, Hetherington AM** (1995) Stimulus-induced oscillations in guard cell cytosolic free calcium. *Plant Cell* **7**: 1207–1219
- Neuhaus G, Bowler C, Kern R, Chua N-H** (1993) Calcium/calmodulin-dependent and -independent phytochrome signal transduction pathways. *Cell* **73**: 937–952
- Nishizuka Y** (1984) Turnover of inositol phospholipids and signal transduction. *Science* **225**: 1365–1370
- Poovaliah BW, Reddy ASN** (1990) Turnover of inositol phospholipids and calcium-dependent protein phosphorylation in signal transduction. In DJ Morré, WF Boss, FA Loewus, eds, *Inositol Metabolism in Plants*. Wiley-Liss, New York, pp 335–349
- Ranjewa R, Boudet AM** (1987) Phosphorylation of proteins in plants: regulatory effects and potential involvement in stimulus/response coupling. *Annu Rev Plant Physiol* **38**: 73–93
- Rudenok AN, Konev SV** (1973) On the phenomenon of cell protection against heat damage (in Russian). *Dokl Acad Nauk SSSR* **208**: 812–816
- Shacklock PS, Read ND, Trewavas AJ** (1992) Cytosolic free calcium mediates red light-induced photomorphogenesis. *Nature* **358**: 753–755
- Shillito R, Potrykus I** (1985) Protoplasts: isolation, culture, plant regeneration. *Methods Enzymol* **118**: 549–578
- Siamon KB, Padgett W, Daly DW** (1981) Forskolin: unique diterpene activation of adenylate cyclase in membrane and in intact cells. *Proc Natl Acad Sci USA* **78**: 3363–3367
- Stone JM, Walker JC** (1995) Plant protein kinase families and signal transduction. *Plant Physiol* **108**: 451–457
- Tepper CG, Jayadev S, Liu B, Bielawska A, Wolff R, Yonehara S, Hannun YA, Seldin MF** (1995) Role for ceramide as an endogenous mediator of Fas-induced cytotoxicity. *Proc Natl Acad Sci USA* **92**: 8443–8447
- Ward JM, Pei Z-M, Schroeder JI** (1995) Roles of ion channels in initiation of signal transduction in higher plants. *Plant Cell* **7**: 833–844