

An Endoplasmic Reticulum-Bound $\text{Ca}^{2+}/\text{Mn}^{2+}$ Pump, ECA1, Supports Plant Growth and Confers Tolerance to Mn^{2+} Stress¹

Zhongyi Wu, Feng Liang², Bimei Hong, Jeff C. Young³, Michael R. Sussman, Jeffrey F. Harper, and Heven Sze*

Department of Cell Biology and Molecular Genetics, University of Maryland, College Park, Maryland 20742 (Z.W., F.L., H.S.); Department of Cell Biology, Plant Division, The Scripps Research Institute, La Jolla, California 92037 (B.H., J.F.H.); and Biotechnology Center, 425 Henry Mall, University of Wisconsin, Madison, Wisconsin 53706 (J.C.Y., M.R.S.)

Plants can grow in soils containing highly variable amounts of mineral nutrients, like Ca^{2+} and Mn^{2+} , though the mechanisms of adaptation are poorly understood. Here, we report the first genetic study to determine in vivo functions of a Ca^{2+} pump in plants. Homozygous mutants of *Arabidopsis* harboring a T-DNA disruption in ECA1 showed a 4-fold reduction in endoplasmic reticulum-type calcium pump activity. Surprisingly, the phenotype of mutant plants was indistinguishable from wild type when grown on standard nutrient medium containing 1.5 mM Ca^{2+} and 50 μM Mn^{2+} . However, mutants grew poorly on medium with low Ca^{2+} (0.2 mM) or high Mn^{2+} (0.5 mM). On high Mn^{2+} , the mutants failed to elongate their root hairs, suggesting impairment in tip growth processes. Expression of the wild-type gene (CAMV35S::ECA1) reversed these conditional phenotypes. The activity of ECA1 was examined by expression in a yeast (*Saccharomyces cerevisiae*) mutant, K616, which harbors a deletion of its endogenous calcium pumps. In vitro assays demonstrated that Ca^{2+} , Mn^{2+} , and Zn^{2+} stimulated formation of a phosphoenzyme intermediate, consistent with the translocation of these ions by the pump. ECA1 provided increased tolerance of yeast mutant to toxic levels of Mn^{2+} (1 mM) and Zn^{2+} (3 mM), consistent with removal of these ions from the cytoplasm. These results show that despite the potential redundancy of multiple Ca^{2+} pumps and $\text{Ca}^{2+}/\text{H}^{+}$ antiporters in *Arabidopsis*, pumping of Ca^{2+} and Mn^{2+} by ECA1 into the endoplasmic reticulum is required to support plant growth under conditions of Ca^{2+} deficiency or Mn^{2+} toxicity.

Plants have a remarkable ability to adapt and grow in soils containing widely different levels of mineral nutrients. To complete a plant's life cycle, macronutrients, like Ca^{2+} and Mg^{2+} , ideally are required at 100- to 300-fold higher levels than that of micronutrients, such as Mn^{2+} and Zn^{2+} . Growth is inhibited when tissue content of a nutrient either drops below a critical concentration (deficiency zone) or increases above an adequate level (toxic zone). Nevertheless, plants demonstrate differences in their abilities to tolerate nutrient deficiency or toxicity (Horst, 1988; Marschner, 1995). For most nutrients, including Ca^{2+} and Mn^{2+} , the molecular bases underlying these differences are poorly understood.

Ca^{2+} and Mn^{2+} are essential for distinct functions in plants. The majority of calcium is associated with

the middle lamella of cell walls, where it plays roles in support and growth. Inside cells, most Ca^{2+} is sequestered within organelles, like the vacuole and the endoplasmic reticulum (ER). The transient release of Ca^{2+} into the cytosol generates calcium waves or oscillations that carry specific information for transduction of hormonal and environmental stimuli (Sanders et al., 1999). Calcium also mediates many developmental processes, including polarized growth, mitosis, and cytokinesis (Hepler and Wayne, 1985). Mn^{2+} is essential for photosynthesis because it is part of the oxygen-evolving complex in photosystem II. This cation is also important for several redox processes and is an activator of enzymes involved in oxidation reduction, decarboxylation, and biosynthesis (Marschner, 1995).

At high concentrations, both Ca^{2+} and Mn^{2+} are potentially toxic and their cytosolic levels are tightly regulated in the range of 0.1 to 0.2 μM (Hepler and Wayne, 1985; Quiquampoix et al., 1993; Sanders et al., 1999). Both cations enter plant cells down an electrochemical gradient (Clarkson, 1988; Sanders et al., 1999). Cytosolic Ca^{2+} is maintained at low levels by ATP-driven pumps and $\text{Ca}^{2+}/\text{H}^{+}$ antiporters located at membranes, including the plasma membrane, vacuole, and ER (Sze et al., 2000). Mn^{2+} is accumulated mostly in the vacuole and chloroplast

¹ This work was supported by the Department of Energy (grant nos. DE-FG02-95ER202200 to H.S. and DE-FG03-94ER20152 to J.F.H.), by the National Science Foundation (grant no. DBI 0077378 to J.F.H.), and by Syngenta (to J.F.H.).

² Present address: Invitrogen Corp., Carlsbad, CA 92008.

³ Present address: Biology Department, Western Washington University, Bellingham, WA 98225.

* Corresponding author; e-mail hs29@umail.umd.edu; fax 301-314-9081.

Article, publication date, and citation information can be found at www.plantphysiol.org/cgi/doi/10.1104/pp.004440.

(McCain and Markley, 1989); however, less is known about active transporters of Mn^{2+} .

Arabidopsis contains 15 putative Ca-ATPases, as predicted from the completed genome sequence (Axelsen and Palmgren, 2001), though the in vivo function of each pump is unknown. ECA1, previously shown to be a Ca^{2+} pump, is specifically blocked by cyclopiazonic acid (CPA; Liang and Sze, 1998), an inhibitor of animal sarcoplasmic/endoplasmic reticulum Ca-ATPase pump. ECA1 is one of four members of the ER-type calcium ATPase subfamily in Arabidopsis (Sze et al., 2000). ACA2 localized to the ER represents another subfamily characterized as autoinhibited calcium ATPase. These Ca^{2+} pumps are inactivated by interactions with an inhibitory region at the N-terminal region and stimulated by calmodulin interaction with the inhibitory region (Hwang et al., 2000). ACA4 and ACA8 are similar types of calmodulin-regulated Ca-ATPases localized at the vacuolar (Geisler et al., 2000) and at the plasma membrane (Bonza et al., 2000), respectively.

Here, we show that the ER-localized Ca^{2+} pump, ECA1, has a dual role in both Ca^{2+} and Mn^{2+} homeostasis. We provide biochemical evidence that ECA1 provides approximately 70% of the total ER-type calcium pump activity in Arabidopsis. Surprisingly, a plant with a T-DNA disruption (*ecal-1*) of this major transport system has a wild-type phenotype when grown under standard nutrient conditions. Nevertheless, under conditions of calcium deprivation, growth of mutant plants is impaired, demonstrating that ECA1 provides an important function in calcium nutrition. We further provide biochemical and genetic evidence that ECA1 behaves

as an Mn^{2+} pump, and that it confers tolerance to toxic levels of Mn^{2+} . These studies provide the first genetic evidence for the in vivo function of a calcium pump in plants.

RESULTS

ECA1 Can Transport Divalent Cations Other Than Ca^{2+}

Specific ions that are transported by P-type ATPases have been shown to stimulate the formation of a phosphorylated intermediate as part of the reaction cycle (MacLennan et al., 1997). To test which ions are potentially transported by ECA1, microsomes isolated from pECA1-transformed K616 yeast (*Saccharomyces cerevisiae*) were incubated with $[\gamma\text{-}^{32}\text{P}]\text{ATP}$ and 5 μM various cations. Very little phosphoprotein was formed in the absence of any divalent cation (+EGTA). Phosphorylation was enhanced by Ca^{2+} , Mn^{2+} , Zn^{2+} , and perhaps by Ni^{2+} , but not with Cd^{2+} (Fig. 1, A–C). The steady-state ^{32}P phosphoenzyme was decreased rapidly by excess unlabeled ATP, indicating the enzyme turns over rapidly. Ca^{2+} did not stimulate any phosphoprotein formation in control membranes isolated from vector-transformed yeast (Liang et al., 1997). The results are consistent with the idea that ECA1 could transport several divalent cations, including Ca^{2+} , Mn^{2+} , Zn^{2+} , and Ni^{2+} .

To confirm that Mn^{2+} is recognized by the Ca^{2+} -binding site(s) on ECA1, Mn^{2+} was tested for its ability to inhibit Ca^{2+} transport in a concentration-dependent manner. Microsomal vesicles were isolated from yeast strains microexpressing ECA1, and ATP-driven $^{45}\text{Ca}^{2+}$ uptake was measured in the presence




Figure 1. ECA1 behaves like a transporter with multi-cation specificity. A through C, Ca^{2+} , Mn^{2+} , and Zn^{2+} stimulated phosphoprotein formation in membranes of ECA1 transformants. ^{32}P ATP (300 nM) was added to a 2-ml reaction mixture containing 25 mM HEPES-1,3-bis(tris[hydroxymethyl]methylamino) propane (BTP; pH 7.0), 100 mM KCl, and 80 μg of vesicle with 0.5 mM EGTA alone (C, ■ and □) or in the presence of 5 μM divalent cations. The final free concentration of the cations was estimated from total cation added in the presence of 0.5 mM EGTA using the Maxchelator program. Aliquots were sampled, and unlabeled ATP was added at 120 s (arrow) to a final concentration of 1 mM. Black and white symbols represent ^{32}P phosphoprotein level before and after addition of ATP, respectively. The data are from the average of two experiments. A, Ca^{2+} , ●; Mn^{2+} , ▲; Cd^{2+} , ■. B, Zn, ●. C, Ni^{2+} , ▲. D, ^{45}Ca transport into membrane vesicles of ECA1 transformants is blocked by Mn. ATP-dependent ^{45}Ca (approximately 0.6 μM at 0.3 $\mu\text{Ci mL}^{-1}$) uptake at 5 min was measured without EGTA in the presence of Mn^{2+} as indicated. Pump activity shown is the difference in uptake by vesicles of mutants transformed with ECA1 and that of vector alone. Average of two experiments.

of an estimated Ca^{2+} concentration of approximately $0.6 \mu\text{M}$. Increasing Mn concentration from 10 nM to $100 \mu\text{M}$ inhibited $^{45}\text{Ca}^{2+}$ transport into vesicles isolated from yeast expressing ECA1. Mn^{2+} had little or no effect on Ca^{2+} binding/uptake in vesicles isolated from yeast harboring the empty vector (not shown). The Mn^{2+} concentration required to inhibit ECA1-dependent Ca^{2+} uptake by 50% was estimated as $0.5 \mu\text{M}$ Mn^{2+} (Fig. 1D), suggesting ECA1 had a high affinity for Mn^{2+} that is similar to that of Ca^{2+} .

Evidence that ECA1 can transport Mn^{2+} and Zn^{2+} in vivo was obtained by showing that ECA1 restored growth of yeast mutant K616 on media containing high Mn^{2+} (1 mM) or Zn^{2+} (3 mM ; Fig. 2). The K616 strain (*pmr1 pmc1 cnb1*) is defective in both a Golgi PMR1 and in a vacuolar PMC1 Ca^{2+} pump (Cunningham and Fink, 1994; Liang et al., 1997), and PMR1 is thought to remove Mn^{2+} from the cytosol and prevent toxicity (Lapinskas et al., 1995). ECA1 can provide the same activity, although a constitutively activated AtACA2-2 cannot (Fig. 2). Moreover, ECA1 also reversed the toxic effects of Zn^{2+} . We had shown before that ECA1 is a Ca^{2+} pump that may also transport Mn^{2+} (Liang et al., 1997; Liang and Sze, 1998). Together, these results indicate that ECA1 behaves as a multi-cation ATPase that transports Ca^{2+} as well as Mn^{2+} and Zn^{2+} .

Identification of a Mutant Plant Carrying a T-DNA Disruption in the ECA1 Gene

To investigate the in vivo functions of ECA1, we obtained a mutant plant, *eca1-1*. A T-DNA insertion associated with the *ECA1* gene (*eca1-1*) was previously detected using a PCR screening strategy ap-

plied to a population of T-DNA-transformed Arabidopsis plants (Krysan et al., 1996). To isolate the individual mutant plant and to determine the exact location of the T-DNA insert, both right and left *ECA1*/T-DNA borders were PCR amplified and sequenced. Both borders contained sequence from the coding region of ECA1, indicating that the T-DNA insertion was not associated with a major deletion or rearrangement of flanking sequences. Sequence analysis indicated that the insertion went into the middle of the last transmembrane domain (TM10; Fig. 3A).

A plant line homozygous for *eca1-1* was identified by PCR analysis after two backcrosses to the parental Wassilewskija (WS)-F ecotype. This backcrossing removed a second unlinked T-DNA insert. A single progeny was identified that had no wild-type *ECA1*. To test for the wild-type gene, a genomic DNA sample was subjected to a PCR amplification reaction using two primers ($5'$ + $3'$ primer pair) corresponding to the $5'$ and $3'$ ends of *ECA1*. In controls using a wild-type DNA template, the $5'$ + $3'$ primer pair amplified an approximately 4.3-kb fragment (Fig. 3B). However, with a DNA template isolated from a homozygous *eca1-1* mutant line, no PCR product was observed, presumably due to the presence of the very large T-DNA insert that prevented the formation of a detectable product. The absence of a wild-type *ECA1* PCR product indicated that the T-DNA insertional mutant was homozygous. As a control to show that the DNA sample was a suitable PCR template, a parallel reaction was conducted in which the $3'$ primer was replaced with T_L , a primer based on the left border of the T-DNA insert (i.e. $5'$ + T_L primer pair). In this control, a PCR product was produced showing the expected size (4.3 kb) for a T-DNA insertion near the $3'$ end of the gene.




Figure 2. ECA1, but not ACA2, expression restored growth of K616 mutant on Mn^{2+} -supplemented medium. Wild-type (W303) or K616 (*pmr1 pmc1 cnb1*) yeast cells were transformed with control vector (p426 Gal1) alone. K616 cells were transformed with pECA1 or pACA2-2. Each transformant was diluted with complete synthetic medium (SC)-uracil (URA)/Gal medium to a density at A_{600} of 1.0, 0.1, 0.01, or 0.001. Then, $10 \mu\text{L}$ of each dilution was dotted on SC-URA/Gal media (control), or with 1 mM MnCl_2 or 3 mM ZnCl_2 , and incubated for 3 d at 30°C .




Figure 3. Position of the T-DNA insert in *eca1-1* mutant. A, T-DNA inserts in the 10th transmembrane (TM) domain of ECA1 protein. Position 1,025 marks the last amino acid of ECA1 in *eca1-1* mutant corresponding to wild-type ECA1. B, PCR analysis showing the absence of a wild-type ECA1 gene in a homozygous *eca1-1* mutant plant. Genomic DNA samples from a wild-type plant (Wt) or *eca1-1* homozygous plant (Mutant) were used as templates in a PCR reaction using a primer pair that can amplify the entire *ECA1* coding sequence (5' + 3'), or a primer pair that amplifies the sequence between the T-DNA left border and the 5' end of *ECA1* (5' + T_L). A picture of a UV-illuminated ethidium bromide-stained gel is shown.

Growth of the *eca1-1* Mutant Is Normal under Standard Conditions But Retarded by Low Ca²⁺

Compared with wild-type plants, there is no obvious phenotype in *eca1-1* homozygous mutants grown on normal 0.5× Murashige and Skoog medium when Ca²⁺ and Mn²⁺ are 1.5 mM and 50 μM, respectively (Fig. 4A, -Mn). Plants also grew similarly on Gamborg's B5 medium containing 0.5% (w/v) MES at pH 5.7, and in various soils under growth room and greenhouse conditions (not shown). *eca1-1* mutant plants appeared to display normal germination, growth rates, morphology, seed set, and gravitropism relative to wild-type plants grown in parallel. They were similar in their fresh weight and their chlorophyll content (Fig. 4, B and D).

To test the effect on growth of reduced Ca²⁺, seeds were germinated on standard 0.5× Murashige and Skoog and 5-d-old seedlings were transferred to a

modified Murashige and Skoog medium with reduced Ca²⁺. After 10 d, mutants and wild-type plants grown in medium containing ≥0.5 mM Ca²⁺ were similar in size although growth in all cases was severely impaired when Ca²⁺ dropped to 0.1 mM or less (not shown). At 0.2 to 0.4 mM Ca²⁺, the mutant consistently showed Ca²⁺ deficiency symptoms not exhibited by wild-type plants. These included small plant size, short roots, small yellowish leaves, and lack of bolts (Fig. 5). The differential traits of mutant and control plants on medium with 0.2 mM Ca²⁺ were consistently observed in three independent experiments.

Growth of *eca1-1* Mutant Is Inhibited by High Mn²⁺

When grown on high (0.5 mM) Mn²⁺, both mutant and wild-type plants had elevated Mn²⁺ content (Ta-




Figure 4. Growth of the *eca1-1* mutant is inhibited by high Mn^{2+} and is rescued by an ECA1 transgene. Seeds of wild-type plant (ECA1), mutant (*eca1-1*), and mutants expressing wild-type ECA1 (35S-#6 and 35S-#7) were germinated on a 0.8% (w/v) agarose plate with 0.5 \times Murashige and Skoog. Five-day-old seedlings were transferred to 0.5 \times Murashige and Skoog medium alone (control, -Mn) or supplemented with 0.5 mM Mn^{2+} (+Mn). A, Plant morphology and size 12 d after transfer. B, Fresh weight of 25 plants 10 d after transfer ($\pm\text{SE}$, $n = 4$). C, Root hairs of plant 12 d after transfer. D, Chlorophyll content from seedlings 10 d after transfer. Average ($\pm\text{SE}$) from four extractions. Bar = 1 mm.

ble I). However, only mutant plants were severely stunted. This phenotype is easily detected after 4 to 5 d, and becomes dramatic after 10 to 12 d (Fig. 4A). The fresh weight of *eca1-1* mutant after 12 d was reduced by 66% relative to wild-type plants (Fig. 4B). Both root elongation and leaf expansion were inhibited. Two weeks later, the mutants appeared to stop growing; however, wild-type plants were pale green and continued to increase in size. At 1 mM Mn^{2+} or higher, both mutant and wild-type plants showed Mn^{2+} toxicity symptoms, although chlorosis appeared earlier (in 2–3 d) in the mutants.

Leaves of mutants growing at 0.5 mM Mn^{2+} were chlorotic and twisted, and young leaves were deformed and stuck together. Wild-type plants growing in media containing 0.5 mM Mn^{2+} showed a 26% drop in chlorophyll content relative to plants cultivated in control medium (Fig. 4D). However, the chlorophyll content of *eca1-1* mutant grown in 0.5 mM Mn^{2+} decreased by 74% relative to plants in normal 0.5 \times Murashige and Skoog.

Inhibition of Root Hair Elongation in the *eca1-1* Mutant

The morphological change in root growth was striking. Figure 4C shows that *eca1-1* mutant roots grown on normal 0.5 \times Murashige and Skoog medium had root hairs similar in length to that of wild-type plants. However, the *eca1-1* mutant, growing on medium supplemented with 0.5 mM Mn^{2+} , had only stubs or very short root hairs. These results indicated that the ability to initiate root hairs was not blocked; however, tip growth and root hair elongation were inhibited.

Reduction of ECA1 Protein and Ca^{2+} Pumping in the *eca1-1* Mutant

Immunostaining showed that ECA1 protein is relatively abundant in the root and the flower, although it is expressed in all major organs of the wild-type plant (Fig. 6A). The expression of a mutant protein in




Figure 5. *eca1-1* ($-/-$) (homozygous) mutant grew poorly on low Ca^{2+} . Seeds were germinated on a 0.8% (w/v) agarose plate with $0.5\times$ Murashige and Skoog salts at pH 5.8. Seedlings (5 d old) of similar size were transferred to medium containing reduced Ca^{2+} (0.2 mM) and grown for 10 d at 21°C . Plants are representative of mutant (*eca1-1*) and wild type (ECA1) from three independent experiments.

a homozygous *eca1-1* plant line was analyzed using three antibodies raised against the C- and N-terminal ends and middle hydrophilic portion of wild-type ECA1, respectively. All antisera detect a 5- to 10-fold reduced signal in a membrane extract from the homozygous mutant, so only results using anti-ECA1(M) are shown (Fig. 6B). Because the mutant *eca1-1p* is expected to have a truncated C-terminal end [hence no anti-ECA1(C)-detectable epitope], the residual signal revealed by anti-ECA1(C) most likely corresponds to a cross-reaction with other related pumps. Anti-ECA1(C) very likely recognizes AtECA4 (Arabidopsis Genome Initiative [AGI] no. At1g07670), which shares 99% protein identity with ECA1 (AGI no. At1g07810), and has only three differences in the C-terminal 27 amino acids used as an antigen to make the anti-ECA1(C) (Liang et al., 1997).

We isolated vesicles from 5-d-old seedlings grown on $0.5\times$ Murashige and Skoog to test Ca^{2+} pump

activity. Total pump activity in the mutant was decreased by 22% relative to that in wild type. However, CPA-sensitive activity was reduced by more than 70% of that in wild type (Fig. 6C), indicating that activity from ECA1-like pumps was specifically impaired. CPA is a specific and potent blocker of ECA1 activity (Liang and Sze, 1998), but not of ACA2 activity (Hwang et al., 2000). Similar results were obtained when Ca^{2+} content inside vesicles was enhanced by oxalate. Thus, a decrease in divalent cation pumping into endoluminal compartments leads to reduced growth and to chlorosis of plants under various nutrient stress conditions.

eca1-1 Is Complemented by the Wild-Type ECA1 Transgene

We tested whether these phenotypic changes were caused by loss of ECA1 function alone. The full

Table 1. Ion content of wild-type and *eca1-1* mutant plants

Five-day seedlings of wild-type plant (ECA1) and mutant (*eca1-1*) were grown in liquid medium with $0.5\times$ Murashige and Skoog alone ($-Mn$) or supplemented with 0.5 mM Mn^{2+} ($+Mn$) for 2 weeks. Ion content of whole plants was analyzed by ICP emission spectrometry. Average of two to three experiments.

| Medium | Plant | Ca | Mn | Mg | Fe | Zn |
|-----------------------------|---------------|-------|-------|-------|-----|-----|
| $\mu\text{g/g dry wt}^{-1}$ | | | | | | |
| $-Mn$ | Wild | 7,796 | 338 | 2,952 | 323 | 247 |
| | <i>eca1-1</i> | 6,913 | 295 | 2,644 | 253 | 200 |
| $+Mn$ | Wild | 7,352 | 2,045 | 2,617 | 272 | 303 |
| | <i>eca1-1</i> | 6,170 | 1,830 | 2,203 | 275 | 202 |




Figure 6. Reduction of ECA1 protein and Ca^{2+} pumping in the *eca1-1* mutant. A, ECA1-like protein is found in all organs of wild-type plants. Total protein extracts (10- μg samples) from roots (R), leaves (L), flowers (F), and siliques (S) were subjected to SDS-PAGE (8% [w/v] gel), transferred to nitrocellulose, and probed with pre-immune or anti-ECA1(C) serum (1:1,000 [v/v]). Secondary antibody (1:5,000 [v/v]) from donkey anti-rabbit IgG was conjugated with horseradish peroxidase, and activity was detected using enhanced chemiluminescence (Amersham-Pharmacia Biotech, Uppsala). The arrow marks the expected position of ECA1 (116 kD). B, ECA1 protein content in mutant and transgenic plants. Microsome (25 μg of protein) from 1-week-old seedlings of wild-type (ECA1), mutant (*eca1-1*), and transgenic mutants expressing ECA1 (35S#6 and 35S#7) were separated by SDS-PAGE (10% [w/v] acrylamide), transferred, and probed with rabbit anti-ECA1 (M; 1:1,000 [v/v]). Secondary antibodies (1:5,000 [v/v]) were linked to alkaline phosphatase. Arrow marks expected size (116 kD) of ECA1. C, Ca^{2+} transport. Assay mixture consisted of 1 mM ATP and 10 μM Ca in the absence of EGTA, and membrane from 1-week-old seedlings of wild type (ECA1) or mutant (*eca1-1*). Pump activity shown is the difference between uptake at 30 min with and without Mg^{2+} . When added, CPA was 100 nmol mg^{-1} protein. ΔCPA , Pump activity sensitive to CPA. Total, Activity without CPA. Average of two experiments.

length ECA1 cDNA under the control of the cauliflower mosaic virus 35S promoter was introduced into the *eca1-1* mutant. Five independent transgenic

lines expressing 35S::ECA1 were chosen for complementation analyses. All five complementation lines showed wild-type phenotype when grown on high (0.5 mM) Mn^{2+} -supplemented medium, and two examples are shown in Figure 4A (right). The fresh weight of lines 35S::ECA1-#6 and 35S::ECA1-#7 was 93% and 126% of that of wild type (Fig. 4B). Chlorophyll content of two transgenic lines, 35S::ECA1-#6 and 35S::ECA1-#7, was 83% and 131% of that in wild type (Fig. 4D). Furthermore, complementation lines showed abundant and elongated root hairs similar to that observed in wild type (Fig. 4C). The restoration of a wild-type phenotype was accompanied by an increase in ECA1 protein (Fig. 6B). Despite the CAMV35S promoter, the ECA1 protein level expressed in transgenic plants was comparable with that in wild-type plants, suggesting potential regulation of translation or protein degradation. Thus, the expression of the wild-type ECA1 gene from a 35S promoter was sufficient to rescue the mutant.

DISCUSSION

The mechanisms by which plants regulate the homeostasis of most essential mineral nutrients are not understood. Insufficient levels of a nutrient can lead to deficiency symptoms, whereas excessive levels are toxic (Horst, 1988; Marschner, 1995). Here, we present: (a) biochemical evidence that ECA1 can transport Mn^{2+} as well as Ca^{2+} , and (b) genetic evidence that this ER-located pump plays critical roles in growth as well as tolerance to Mn^{2+} toxicity.

T-DNA Insertion of the ECA1 Gene Reduced Ca^{2+} Pumping in the Mutant

To understand *in vivo* functions of ECA1 pump, we isolated a homozygous plant line harboring a T-DNA disruption of *ECA1* (*ECA1::T-DNA-1* = *eca1-1*). The T-DNA insertion site is predicted to disrupt the last (10th) putative transmembrane domain (see Fig. 3). Although a mutant *eca1-1* protein may be synthesized, immunoblot analyses indicates that the expression of ECA1-like protein is reduced 5- to 10-fold (Fig. 6B). It is likely that membrane proteins that fail to fold into a proper conformation are selectively removed from the ER and degraded (Kopito, 1997). Alternatively, the T-DNA insertion may disrupt a transcriptional control or create an unstable mRNA. The residual protein detected by three different antibodies is likely due to the presence of another closely related isoform that shares 97% identity to AtECA4 (AGI no. At1g07670). Regardless of the mechanism, evidence indicates that the T-DNA insertion dramatically reduces both the accumulation of the ECA1-like protein and Ca^{2+} pump activity into endomembranes (Fig. 6).

ECA1 Maintains Ion Homeostasis by Distributing Internal Ca^{2+} and Mn^{2+}

When grown in the same nutrient medium, the total ion content of wild-type or mutant plants were similar (Table I), indicating that net ion uptake was little or not altered. It is interesting that *eca1-1* mutant grown in 0.5× Murashige and Skoog medium showed little or no obvious physical differences from wild-type plants. Apparently, other intracellular cation pumps and cation/ H^+ antiporters could compensate in part for the loss of ECA1 activity when extracellular Ca^{2+} or Mn^{2+} was 1.5 mM or 50 μM , respectively. Under these conditions, it is conceivable that cells use a calmodulin-stimulated ACA2 to load Ca^{2+} into the ER lumen (Hong et al., 1999; Hwang et al., 2000), and CAX2 to remove cytosolic Mn^{2+} into small vacuoles (Hirschi et al., 2000).

Mn^{2+} Toxicity and Tolerance

However, physical symptoms of Ca^{2+} deficiency became evident in mutants when external Ca^{2+} was dropped to 0.2 to 0.4 mM. Furthermore, a 10-fold elevation in Mn^{2+} accentuated symptoms typical of Ca^{2+} , Mg^{2+} , and Fe^{2+} deficiencies (Horst and Marschner, 1978; Marschner, 1995) in the mutant, but less so in the wild-type plants. We have initially focused on the increased sensitivity to Mn^{2+} toxicity because symptoms of Ca^{2+} deficiency were less dramatic, and symptoms of Zn^{2+} deficiency or toxicity were not observed in preliminary studies. When plants were exposed to high levels of Mn^{2+} , both wild-type and mutant plants accumulated Mn^{2+} to similar, potentially toxic, levels (Table I), consistent with the idea that the *eca1-1* mutation did not alter uptake of Mn^{2+} into the plant. Thus, the difference in Mn^{2+} tolerance between mutant and wild-type plants may be caused by differential internal distribution of ions within tissues or cells. This idea was suggested for two cotton (*Gossypium hirsutum*) genotypes that differed in their ability to tolerate elevated Mn^{2+} (Foy et al., 1995).

Although measurements of ion content in the cytosol and intracellular compartments of *eca1-1* mutants are not yet available, we offer several interpretations for observed Mn sensitivity of *eca1-1*. The average Mn^{2+} content in plant tissues is low (20–50 $\mu\text{g g dry weight}^{-1}$). Mn^{2+} is accumulated in the vacuole and chloroplast in leaves (McCain and Markley, 1989), whereas cytosolic Mn^{2+} is estimated as less than 0.2 μM in roots (Quiquampoix et al., 1993). High external Mn^{2+} can compete for divalent cation-binding sites, thus reducing uptake of Ca^{2+} , Mg^{2+} , and Fe^{2+} (Marschner, 1995). In theory, this competition could also occur at sites of intracellular transport. Thus, high cytosolic Mn^{2+} potentially interferes with proper uptake and sorting of Ca^{2+} , Mg^{2+} , and Fe^{2+} into intracellular locations. Inadequate Mg^{2+} and Fe^{2+} in the chloroplasts could result in chlorosis

because they are required for chlorophyll synthesis (Csatorday et al., 1984) and as cofactors for many metalloenzymes (Marschner, 1995). Perturbations in cytosolic Ca^{2+} signaling could also result in poor growth. Interference of Ca^{2+} sequestration into organelles could potentially change the shape and frequency of Ca^{2+} oscillations that are so critical to specific signaling events (Sanders et al., 1999; Sze et al., 2000).

We propose that ECA1 could promote growth and confer Mn^{2+} tolerance in several ways. By pumping Mn^{2+} into the ER, ECA1 could reduce cytosolic Mn^{2+} to levels that do not interfere with the internal distribution of Mg^{2+} , Fe^{2+} , or Ca^{2+} . This would reduce the induced deficiency of these cations, and also restore, for instance, signal-induced Ca^{2+} transients normally seen in wild-type plants. Under conditions of Ca^{2+} deficiency when cytosolic Ca^{2+} level would be very low, the high-affinity ECA1 may be the only pump capable of loading Ca^{2+} into the ER lumen (Liang and Sze, 1998) for functioning of the secretory system and for stimuli-induced Ca release (Sze et al., 2000).

Roles of ECA1

The observation of an *eca1-1* phenotype demonstrates that ECA1 has unique functions, despite multiple calcium pumps and cation exchangers in plants. Biochemical studies demonstrate that ECA1 differs from other ACA2-like Ca^{2+} pumps in its (a) ion specificity, (b) high affinity for cations, and (c) subcellular location (Liang et al., 1997; Liang and Sze, 1998; Hwang et al., 2000; Sze et al., 2000). Although ACA2 is present in the ER of some cell types (Hong et al., 1999), it does not transport Mn^{2+} (Fig. 2) and therefore is unable to replace ECA1. Moreover, ECA1 protein may be more abundant than ECA2-ECA4 because it contributes 70% of Ca^{2+} pump activity inhibitable by CPA. As a high-affinity divalent cation pump with a K_{mCa} estimated at 0.03 μM (Liang and Sze, 1998), ECA1 would be more effective than other Ca^{2+} pumps or $\text{Ca}^{2+}/\text{H}^+$ antiporters when cytosolic [Ca^{2+}] is extremely low.

The inhibition of growth by high Mn^{2+} in *eca1-1* Arabidopsis plants and in the yeast *pmr1* mutant are remarkably similar, suggesting that disruptions in intracellular Ca^{2+} and Mn^{2+} homeostasis affect fundamental processes of cell division and cell expansion. For example, Mn^{2+} may be important for cell cycle progression in place of Ca^{2+} as shown in yeast (Loukin and Kung, 1995), and for activation of Mn-dependent glycosyltransferases involved in protein processing and cell wall synthesis (White et al., 1993; Sterling et al., 2001). Yeast *pmr1* mutants lacking a Golgi $\text{Ca}^{2+}/\text{Mn}^{2+}$ pump (Mandal et al., 2000) showed defects in glycosylation and protein sorting (Durr et al., 1998), and *Drosophila melanogaster* cells lacking a sarcoplasmic/endoplasmic reticulum-type Ca-ATPase

is defective in Notch trafficking (Periz and Fortini, 1999). Furthermore, elongating root hairs and pollen tubes depend on a tip-focused Ca^{2+} gradient that is most likely dependent on spatially localized Ca^{2+} channels (Very and Davies, 2000) and active transporters. Perturbations of the Ca^{2+} gradient alter root hair growth (Bibikova et al., 1997). Although the mechanism of this is still unknown, Ca^{2+} pumps associated with the cortical ER potentially alter the shape or frequency of Ca^{2+} oscillations that accompany tip growth (Holdaway-Clarke et al., 1997).

This study of the first plant mutant with a disruption of an ER-type Ca^{2+} pump has provided two new insights. First, despite a 70% reduction in ER-type calcium pump activity, a plant can complete its life cycle under conditions of "optimal" nutrient availability. Second, despite the presence of 14 Ca^{2+} pumps and 11 CAX1-like antiporters in Arabidopsis, the disruption of ECA1 revealed a critical role of this single pump in plants grown under conditions of low Ca^{2+} or high Mn^{2+} . Thus, ECA1 clearly functions in mineral nutrition and makes plants more adaptable to soils with variable nutrient conditions. Whether ECA1 also plays a specific role in calcium signaling will require further studies.

MATERIALS AND METHODS

Yeast (*Saccharomyces cerevisiae*) Strain, Plasmid, and Growth Medium

Yeast strains W303-1A (*MATa*, *leu2*, *his3*, *ade2*, *trp1*, and *ura3*) and K616 (*MATa pmr1::HIS3 pmc1::TRP1 cnb1::LEU2* and *ura3*) were used (Cunningham and Fink, 1994). The entire ECA1 cDNA (U96455; Liang et al., 1997) and the truncated ACA2-2 cDNA (Harper et al., 1998; L08469) from Arabidopsis were constructed into the yeast expression vector p426Gal1 (Liang et al., 1997). The K616 mutant was transformed, selected on SC-URA, and grown on medium with 2% (w/v) Gal as described (Liang et al., 1997).

PCR Analysis of T-DNA Tags in Arabidopsis

Genomic DNA was isolated from individual homozygous *eca1-1* mutant and wild-type Arabidopsis WS-F plants according to Krysan et al. (1996). The PCR primers sequences are as follows: T-DNA left border (T_L), dGAT-GCACTCGAAATCAGCCAATTTAGAC; ECA1 primer (5'), dGAGTTTC-CGGGAGAATTTGACGAATCTGT; and ECA1 primer (3'), dCCAACGC-CGAGGTAAGTAAACGCTAAT. Genomic DNA samples (40 ng) from a wild-type plant or *eca1-1* homozygous plant were used as templates in a PCR reaction using a primer pair that can amplify the entire ECA1 coding sequence (5' + 3'), or a primer pair that amplifies the sequence between the T-DNA left border and the 5' end of ECA1 (5' + T_L). PCR reactions were performed using *X-Taq* and buffers (PanVera Corporation, Madison, WI) for 41 cycles as follows: 94°C for 30 s, 65°C for 1 min, and 72°C for 2 min.

Plant Transformation

eca1-1 (−/−) (homozygous) mutant was transformed with the CAMV35S-ECA1 construct via vacuum infiltration (Bechtold et al., 1993). Kanamycin-resistant plants were selected and five independent lines expressing the ECA1 transgene were used for complementation analyses.

Growth Conditions of Arabidopsis

The homozygous *eca1-1* mutant and transformants were grown side by side with parental wild-type WS-F plants under the same conditions. Seeds

were surface sterilized, vernalized at 4°C for 48 h, and germinated vertically on 0.8% (w/v) agarose plates with 0.5× Murashige and Skoog medium (Murashige and Skoog, 1962) at pH 5.7 under constant illumination of approximately 100 $\mu\text{E m}^{-2} \text{s}^{-1}$ at 21°C. To test effects of Ca^{2+} or Mn^{2+} on growth, 5-d-old seedlings were transferred to agarose plates containing 0.5× Murashige and Skoog (control), or a modified Murashige and Skoog with 0.1 to 1 mM Ca^{2+} or supplemented with 0.5 to 2.0 mM Mn^{2+} in the absence of EGTA.

Ion Content

Mutant and wild-type seeds were supported on filter paper dipped in 0.5× Murashige and Skoog and germinated vertically in Magenta boxes for 5 d. Seedlings were transferred to either control (0.5× Murashige and Skoog alone) or medium supplemented with 0.5 mM Mn^{2+} for 2 weeks. Plants (15–20 g fresh weight) were dried at 70°C for 3 to 4 d, and ashed at 480°C for 16 h. Ash was digested with 2 mL of concentrated HNO_3 , and the dry sample was dissolved in 10 mL of 3 N HCl. Ion content was analyzed with a Liberty 150 ICP Emission Spectrometer (Varian, Palo Alto, CA) using cobalt as an internal standard.

Chlorophyll Content

Seedlings (approximately 0.5 g) were pulverized in liquid nitrogen and extracted in 2 mL and then diluted with 80% (v/v) ice-cold acetone to 3 mL. The supernatant, collected after centrifuging at 5,000g for 15 min, was examined at 663 and 645 nm, and the chlorophyll concentration was determined using the equation:

$$\mu\text{g chlorophyll mL}^{-1} = 20.2 \times A_{645} + 8.02 \times A_{663}$$

Isolation of Membrane Vesicles from Yeast and Plants

One-week-old seedlings (8 g) were homogenized in 10 mL of buffer containing 50 mM HEPES-BTP (pH 7.4), 250 mM sorbitol, 6 mM EGTA, 0.5% (w/v) bovine serum albumin, 1 mM dithiothreitol, 1 mM phenylmethylsulfonyl fluoride, 0.1 mM tosylsulfonyl phenylalanyl chloromethyl ketone, 2 mM benzamidine, 1 $\mu\text{g mL}^{-1}$ leupeptin, and 5 $\mu\text{g mL}^{-1}$ pepstatin A. The homogenate was centrifuged at 10,000g for 15 min, and the supernatant was either centrifuged at 110,000g for 50 min to get a microsomal pellet for SDS-PAGE, or layered on two 16-mL tubes with 15%/35% (w/w; 5 mL each) Suc step gradient (in buffer: 10 mM HEPES-BTP, pH 7.2; 50 mM KCl; 1 mM dithiothreitol; 1 mM phenylmethylsulfonyl fluoride; 0.1 mM tosylsulfonyl phenylalanyl chloromethyl ketone; and 2 mM benzamidine), and centrifuged at 100,000g for 3 h. Vesicles at the 15%/35% (w/w) Suc interface were collected (3–4 mL) for calcium transport. Yeast vesicles were isolated as described (Liang and Sze, 1998). Protein concentration was determined with the Bio-Rad reagent (Bio-Rad Laboratories, Hercules, CA).

Calcium Transport

Ca^{2+} uptake into membrane vesicles was measured without EGTA by the filtration method (Liang and Sze, 1998). Transport with plant vesicle was initiated with 1 mM ATP in a reaction mixture (250 μL) containing 250 mM Suc, 25 mM HEPES/BTP (pH 7.0), 10 mM KCl, 0.4 mM NaN_3 , and 10 μM $^{45}\text{CaCl}_2$ (2 $\mu\text{Ci mL}^{-1}$, PerkinElmer Life Sciences, Boston) with or without 1 mM MgSO_4 . To block $\text{H}^+/\text{Ca}^{2+}$ antiport, mixture included 0.5 μM bafilomycin A1 and 5 μM gramicidin D. CPA was incubated with membranes at 20°C for 15 min before the reaction was started. Transport of yeast vesicles was similar except that ATP and MgSO_4 is 3 mM, $^{45}\text{CaCl}_2$ is 0.3 $\mu\text{Ci mL}^{-1}$ (estimated to total calcium 0.6 μM), and the incubation time is 5 min at 20°C.

Antibody Production

Three separate rabbit polyclonal anti-ECA1 were produced. Anti-ECA1(C) number 1,374 was raised against a fusion protein, encoded by the plasmid pECA1-c, containing the last 27 residues of ECA1p fused to the C-terminal end of glutathione S-transferase (GST). Anti-ECA1(N) number 1,705 was also raised against a GST fusion protein, encoded by the plasmid

pIN-ECA1-N, containing the first 85 residues of ECA1p. Both GST fusion proteins were constructed using the parent vector pGEX-2T (Smith and Johnson, 1988). The fusion proteins were expressed in *Escherichia coli* DH10 α , and affinity purified over a glutathione-agarose column (Pharmacia Biotech, Piscataway, NJ). Anti-ECA1(M) number UMY31 was raised against a C-6 \times His tag fusion protein containing residues Met-393-Gln-605 of ECA1. The fusion protein was constructed using the parent vector pET28b at the *Nco*I/*Sal*I insertion site. The fusion protein was expressed in BL21 cells, and first purified through Probond nickel-chelating resin according to the manufacturer's protocol (Invitrogen Company Xpress System, Invitrogen, Carlsbad, CA) and purified by SDS-PAGE (approximately 25 kD). The purified fusion protein samples were injected into New Zealand white rabbits with RIBI adjuvant as recommended by the manufacturer (RIBI ImmunoChem Research). Serum of final bleeds recognized a 116-kD protein in immunoblots of microsomal proteins from ECA1 transformants.

Distribution of Materials

Upon request, all novel materials described in this publication will be made available in a timely manner for noncommercial research purposes, subject to the requisite permission from any third party owners of all or parts of the material. Obtaining any permission will be the responsibility of the requestor.

ACKNOWLEDGMENTS

We thank Dr. Rong Yu (University of Maryland, College Park) for her assistance and Drs. Carrie Green and Rufus Chaney (U.S. Department of Agriculture, Beltsville, MD) for discussions and help with ion content analyses.

Received February 22, 2002; returned for revision April 11, 2002; accepted April 24, 2002.

LITERATURE CITED

- Axelsen KB, Palmgren MG (2001). Inventory of the superfamily of P-type ion pumps in Arabidopsis. *Plant Physiol* **126**: 696–706
- Bechtold N, Ellis J, Pelletier G (1993). In planta *Agrobacterium* mediated gene transfer by infiltration of adult *Arabidopsis thaliana* plants. *C R Acad Sci Paris Life Sci* **316**: 1194–1199
- Bibikova TN, Zhigilei A, Gilroy S (1997). Root hair growth in *Arabidopsis thaliana* is directed by calcium and an endogenous polarity. *Planta* **203**: 495–505
- Bonza MC, Morandini P, Luoni L, Geisler M, Palmgren MG, De Michelis MI (2000) At-ACA8 encodes a plasma membrane-localized calcium-ATPase of Arabidopsis with a calmodulin-binding domain at the N terminus. *Plant Physiol* **123**:1495–1506
- Clarkson D (1988). The uptake and translocation of manganese by plant roots. In RD Graham, ed, *Manganese in Soils and Plants*. Kluwer Academic Publishers, Dordrecht, The Netherlands, pp 101–111
- Csatorday K, Gombos Z, Szalontai B (1984). Mn²⁺ and Co²⁺ toxicity in chlorophyll biosynthesis. *Proc Natl Acad Sci USA* **81**: 476–478
- Cunningham KW, Fink GR (1994) Calcineurin-dependent growth control in *Saccharomyces cerevisiae* mutants lacking PMCl, a homolog of plasma membrane Ca²⁺ ATPases. *J Cell Biol* **124**: 351–363
- Durr G, Strayle J, Plemper R, Elbs S, Klee SK, Catty P, Wolf DH, Rudolph HK (1998) The medial-Golgi ion pump Pmr1 supplies the yeast secretory pathway with Ca²⁺ and Mn²⁺ required for glycosylation, sorting, and endoplasmic reticulum-associated protein degradation. *Mol Biol Cell* **9**: 1149–1162
- Foy CD, Weil RR, Coradetti CA (1995) Differential Mn tolerance of cotton genotypes in nutrient solution. *J Plant Nutr* **18**: 685–706
- Geisler M, Frangne N, Gomes E, Martinoia E, Palmgren MG (2000) The ACA4 gene of Arabidopsis encodes a vacuolar membrane calcium pump that improves salt tolerance in yeast. *Plant Physiol* **124**:1814–1827
- Harper JF, Hong B, Hwang I, Guo HQ, Stoddard R, Huang JF, Palmgren MG, Sze H (1998) A novel calmodulin-regulated Ca²⁺-ATPase (ACA2) from Arabidopsis with an N-terminal autoinhibitory domain. *J Biol Chem* **273**: 1099–1106
- Hepler PK, Wayne RO (1985) Calcium and plant development. *Annu Rev Plant Physiol* **36**: 397–439
- Hirschi KD, Korenkov VD, Wilganowski NL, Wagner GJ (2000) Expression of Arabidopsis CAX2 in tobacco. Altered metal accumulation and increased manganese tolerance. *Plant Physiol* **124**: 125–133
- Holdaway-Clarke TL, Feijo JA, Hackett GR, Kunkel JG, Hepler PK (1997) Pollen tube growth and the intracellular cytosolic calcium gradient oscillate in phase while extracellular calcium influx is delayed. *Plant Cell* **9**: 1999–2010
- Hong B, Ichida A, Wang Y, Gens JS, Pickard BG, Harper JF (1999) Identification of a calmodulin-regulated Ca²⁺-ATPase in the endoplasmic reticulum. *Plant Physiol* **119**: 1165–1176
- Horst WJ (1988) The physiology of manganese toxicity. In RD Graham, RJ Hannam, NC Uren, eds, *Manganese in Soil and Plants*. Kluwer Academic Publishers, Dordrecht, The Netherlands, pp 175–188
- Horst WJ, Marschner H (1978) Effect of excess Mn on uptake and translocation of calcium in bean plants (*Phaseolus vulgaris* L.). *Z Pflanzenphysiol* **87**: 137–148
- Hwang I, Harper JF, Liang F, Sze H (2000) Calmodulin activation of an endoplasmic reticulum-located calcium pump involves an interaction with the N-terminal autoinhibitory domain. *Plant Physiol* **122**: 157–168
- Kopito RR (1997) ER quality control: the cytoplasmic connection. *Cell* **88**: 427–430
- Krysan PJ, Young JC, Tax F, Sussman MR (1996) Identification of transferred DNA insertions within Arabidopsis genes involved in signal transduction and ion transport. *Proc Natl Acad Sci USA* **93**: 8145–8150
- Lapinskas PJ, Cunningham KW, Liu XF, Fink GR, Culotta VC (1995) Mutations in PMR1 suppress oxidative damage in yeast cells lacking superoxide dismutase. *Mol Cell Biol* **15**: 1382–1388
- Liang F, Cunningham KW, Harper JF, Sze H (1997) ECA1 complements yeast mutants defective in Ca²⁺ pumps and encodes an endoplasmic reticulum-type Ca²⁺-ATPase in *Arabidopsis thaliana*. *Proc Natl Acad Sci USA* **94**: 8579–8584
- Liang F, Sze H (1998) A high-affinity Ca²⁺ pump (ECA1) from the endoplasmic reticulum is inhibited by cyclopiiazonic acid but not by thapsigargin. *Plant Physiol* **118**: 817–825
- Loukin S, Kung C (1995) Manganese effectively supports yeast cell-cycle progression in place of calcium. *J Cell Biol* **131**: 1025–1037
- MacLennan DH, Rice WJ, Green NM (1997) The mechanism of Ca²⁺ transport by sarco (endo) plasmic reticulum Ca²⁺-ATPases. *J Biol Chem* **272**: 28815–28818
- Mandal D, Woolf TB, Rao R (2000) Manganese selectivity of Pmr1, the yeast secretory pathway ion pump, is defined by residue Gln783 in transmembrane segment 6. Residue Asp778 is essential for cation transport. *J Biol Chem* **275**: 23933–23938
- Marschner H (1995) *Mineral Nutrition of Higher Plants*, Ed 2. Academic Press, San Diego
- McCain D, Markley J (1989) More manganese accumulates in maple sun leaves than in shade leaves. *Plant Physiol* **90**: 1417–1421
- Murashige T, Skoog F (1962) A revised medium for rapid growth and bioassay with tobacco tissue culture. *Physiol Plant* **15**: 473–497
- Periz G, Fortini ME (1999) Ca²⁺-ATPase function is required for intracellular trafficking of the Notch receptor in *Drosophila*. *EMBO J* **18**: 5983–5993
- Quiquampoix H, Loughman BC, Ratcliffe RG (1993) A 31P-NMR study of the uptake and compartmentation of manganese by maize roots. *J Exp Bot* **44**: 1819–1827
- Sanders D, Brownlee C, Harper JF (1999) Communicating with calcium. *Plant Cell* **11**: 691–706
- Smith DB, Johnson KS (1988) Single-step purification of polypeptides expressed in *Escherichia coli* as fusions with glutathione S-transferase. *Gene* **67**: 31–40
- Sterling JD, Quigley HF, Orellana A, Mohnen D (2001) The catalytic site of the pectin biosynthetic enzyme alpha-1,4-galacturonosyltransferase is located in the lumen of the Golgi. *Plant Physiol* **127**: 360–371
- Sze H, Liang F, Hwang I, Curran AC, Harper JF (2000) Diversity and regulation of plant Ca²⁺ pump: insights from expression in yeast. *Annu Rev Plant Physiol Plant Mol Biol* **51**: 433–462
- Very AA, Davies JM (2000) Hyperpolarization-activated calcium channels at the tip of Arabidopsis root hairs. *Proc Natl Acad Sci USA* **97**: 9801–9806
- White AR, Xin Y, Pezeshk V (1993) Xyloglucan glucosyltransferase in Golgi membranes from *Pisum sativum* (pea). *Biochem J* **294**: 231–238