

Elucidation of the Indirect Pathway of Abscisic Acid Biosynthesis by Mutants, Genes, and Enzymes¹

Steven H. Schwartz, Xiaoqiong Qin, and Jan A.D. Zeevaart*

Department of Energy-Plant Research Laboratory (S.H.S., J.A.D.Z.) and Department of Plant Biology (J.A.D.Z.), Michigan State University, East Lansing, Michigan 48824–1312; and Department of Biochemistry, University of Saskatchewan, Saskatoon, Canada S7N 5E5 (X.Q.)

Abscisic acid (ABA) was discovered independently by several groups in the early 1960s. Originally believed to be involved in the abscission of fruit and dormancy of woody plants, the role of ABA in these processes is still not clear. ABA is, however, necessary for seed development, adaptation to several abiotic stresses, and sugar sensing. The regulation of these processes is in large part mediated by changes in *de novo* synthesis of ABA.

Two pathways have been proposed for the synthesis of ABA. In the “direct pathway,” which operates in some fungi, ABA is derived from farnesyl diphosphate (Hirai et al., 2000). Because of structural similarities, an “indirect pathway” in which ABA is produced from the cleavage of carotenoids also had been proposed (Taylor and Smith, 1967). The first committed step for ABA synthesis in plants is the oxidative cleavage of a 9-cis-epoxycarotenoid (C₄₀) to produce xanthoxin (C₁₅) and a C₂₅ by-product (Fig. 1). The 4'-hydroxyl of xanthoxin is oxidized to a ketone by an NAD-requiring enzyme. As a consequence, there is a nonenzymatic desaturation of the 2'-3' bond and opening of the epoxide ring to form abscisic aldehyde. In the final step of the pathway, abscisic aldehyde is oxidized to ABA.

Evidence for the indirect pathway in plants had initially come from a variety of biochemical studies, ¹⁸O₂-labeling experiments, and the characterization of ABA-deficient mutants. In recent years, the genes encoding enzymes for many steps in the pathway have been identified. Much of the recent work in characterizing these genes has confirmed previous biochemical studies. Advances in the elucidation of the ABA biosynthetic pathway and its regulation also have allowed the manipulation of ABA levels in transgenic plants. Of particular interest is the cloning and characterization of the nine-cis-epoxycarotenoid dioxygenases (NCEDs) that catalyze the rate-limiting step in ABA synthesis. The identification of the NCEDs also has had an impact beyond plant biology.

Similar proteins are present in a diverse array of organisms. Their enzymatic activities are responsible for the synthesis of a variety of compounds from carotenoids, including vitamin A in animals.

ABA-DEFICIENT MUTANTS

Our understanding of the functions and synthesis of ABA has been greatly enhanced by the identification and characterization of ABA-deficient mutants (Table I). The ABA-deficient mutants have been identified by the following phenotypes: precocious germination, susceptibility to wilting, an increase in stomatal conductance, and an ability to germinate and grow on media containing a high concentration of Suc or salt. In recent years, these mutants have also been very useful in cloning the genes that encode ABA biosynthetic enzymes.

The pathway of ABA synthesis can be traced back to the early steps of isoprenoid synthesis in plastids (Rodríguez-Concepción and Boronat, 2002). Isoprenoids are an extremely diverse class of natural products that serve a variety of functions in plants. Although it is not a committed step in ABA synthesis, the epoxidation of zeaxanthin seems to be a reasonable place to begin a review on ABA synthesis in plants (Fig. 1). Mutants impaired in the epoxidation of zeaxanthin were first identified by an ABA-deficient phenotype. In addition, the ZEP appears to have a role in the regulation of ABA synthesis in non-chlorophyllous organs.

The *aba1* mutant in *Arabidopsis* (Karssen et al., 1983; Duckham et al., 1991; Rock and Zeevaart, 1991a) and the *aba2* mutant in *N. plumbaginifolia* (Marin et al., 1996) both contain lesions in the enzyme that catalyzes the epoxidation of zeaxanthin to antheraxanthin and violaxanthin (ZEP in Fig. 1). To avoid confusion, the genes corresponding to these mutants will subsequently be referred to as *AtZEP* and *NpZEP*, respectively.

The *atzep* mutant provided definitive evidence that ABA is derived from an epoxycarotenoid precursor (Duckham et al., 1991; Rock and Zeevaart, 1991a). This mutant also has been used extensively to characterize the role of epoxycarotenoids in the xanthophyll cycle and as components of the light-harvesting complexes (Lokstein et al., 2002).

¹ This work was supported by the National Science Foundation (grant no. IBN-9982758) and by the U.S. Department of Energy (grant no. DE-FG02-91ER20021).

* Corresponding author; e-mail zeevaart@msu.edu; fax 517-353-9168.

www.plantphysiol.org/cgi/doi/10.1104/pp.102.017921.




Figure 1. The pathway of ABA synthesis beginning with zeaxanthin. The steps in isoprenoid and carotenoid synthesis before zeaxanthin are reviewed elsewhere (Cunningham and Gantt, 1998; Rohmer, 1999; Hirschberg, 2001; Rodríguez-Concepción and Boronat, 2002). VDE, Violaxanthin de-epoxidase.

An epoxidase mutant in *N. plumbaginifolia*, *npzep*, has been identified and the corresponding gene cloned (Marin et al., 1996). The NpZEP protein, which is similar to some bacterial monooxygenases, was able to catalyze the epoxidation of zeaxanthin to antheraxanthin and violaxanthin. For this activity, it was necessary to add an additional component from chloroplasts. It was later determined that reduced ferredoxin is a necessary cofactor (Bouvier et al., 1996).

Because the level of epoxy-carotenoids in green leaves is high relative to the amount of ABA synthesized, it is considered unlikely that ZEP has a regulatory role in these tissues. The expression of ZEP transcripts in green tissue does not increase in wild tobacco (*Nicotiana plumbaginifolia*; Audran et al., 1998), tomato (Thompson et al., 2000a), or cowpea (*Vigna unguiculata*; Iuchi et al., 2000) that are subjected to osmotic stress. In etiolated tissues, the concentration of carotenoids is significantly lower and the increased expression of ZEP mRNA does correlate with elevated ABA synthesis in roots and seeds

(Audran et al., 1998; Borel et al., 2001). The overexpression of ZEP in transgenic tobacco resulted in increased seed dormancy (Frey et al., 1999), thus providing further evidence that the level of epoxy-carotenoids limits ABA synthesis in some tissues.

The Arabidopsis epoxidase, *AtZEP*, was cloned by sequence similarity to the tobacco ZEP (Audran et al., 2001; Xiong et al., 2002). Contrasting reports on the expression of *AtZEP* mRNA have appeared in the literature. In one study, it was found that the level of *AtZEP* mRNA was induced by drought stress in root tissues (Audran et al., 2001). The expression of the *AtZEP* transcript was unaffected in several ABA-deficient and -insensitive mutants (Audran et al., 2001). In another study, it was reported that the expression of *AtZEP* mRNA increased in response to osmotic stress or ABA treatment in both roots and shoots (Xiong et al., 2002). The osmotic induction of *AtZEP* transcript was impaired in ABA-deficient mutants and in the ABA-insensitive mutant, *abi1*. Several additional genes necessary for the later steps in ABA synthesis also were found to be induced by

Table 1. ABA biosynthesis mutants with position of the lesions in the pathway and functions of the wild-type proteins

Mutant	Species	Biochemical Step	Enzyme	References
<i>aba1/npq2/los6^a</i>	Arabidopsis	Epoxidation of zeaxanthin	Zeaxanthin epoxidase (ZEP)	Karssen et al. (1983); Rock and Zeevaart (1991a); Niyogi et al. (1998); Xiong et al. (2002)
<i>aba2</i>	<i>Nicotiana plumbaginifolia</i>	Epoxidation of zeaxanthin	ZEP	Marin et al. (1996)
<i>osaba1</i>	Rice (<i>Oryza sativa</i>)	Epoxidation of zeaxanthin	ZEP	Agrawal et al. (2001)
<i>vp14</i>	Maize (<i>Zea mays</i>)	Oxidative cleavage of neoxanthin or violaxanthin	NCED	Schwartz et al. (1997b); Tan et al. (1997)
<i>not (notabilis)</i>	Tomato (<i>Lycopersicon esculentum</i>)	Oxidative cleavage of neoxanthin or violaxanthin	NCED	Burbidge et al. (1999)
<i>aba2/gin1/isi4/sis4</i>	Arabidopsis	Oxidation of xanthoxin to abscisic aldehyde	Short-chain alcohol dehydrogenase	Léon-Kloosterziel et al. (1996); Schwartz et al. (1997a); Laby et al., 2000; Rook et al. (2001); Cheng et al. (2002)
<i>flc</i>	Tomato	Oxidation of abscisic aldehyde to ABA	Molybdenum cofactor (MoCo) sulfuryase	Sagi et al. (2002)
<i>aba3/los5/gin5</i>	Arabidopsis	Oxidation of abscisic aldehyde to ABA	MoCo sulfuryase	Léon-Kloosterziel et al., (1996); Schwartz et al. (1997a); Xiong et al. (2001); Cheng et al. (2002)
<i>sit</i>	Tomato	Oxidation of abscisic aldehyde to ABA	Aldehyde oxidase	Okamoto et al. (2002)
<i>aba1</i>	<i>N. plumbaginifolia</i>	Oxidation of abscisic aldehyde to ABA	MoCo sulfuryase?	Akaba et al. (1998)
<i>nar2A</i>	Barley (<i>Hordeum vulgare</i>)	Oxidation of abscisic aldehyde to ABA	MoCo synthesis	Walker-Simmons et al. (1989)
<i>ao3</i>	Arabidopsis	Oxidation of abscisic aldehyde to ABA	Abscisic aldehyde oxidase (AAO3)	Seo et al. (2000b)

^aIn several cases, allelic mutants that were isolated by different laboratories have been assigned different gene symbols.

stress and ABA (Xiong et al., 2001, 2002). The authors suggested that ABA synthesis might be subject to positive feedback regulation.

The significance of ZEP up-regulation in green leaves is uncertain. The ¹⁸O₂-labeling experiments, which were instrumental in establishing the indirect pathway of ABA synthesis, also provide some indication of flux through the pathway. The 1'-hydroxyl in ABA is derived from the epoxide in the carotenoid precursor. In ¹⁸O₂-labeling experiments, there is little incorporation of ¹⁸O at this position for time points less than 8 h (Zeevaart et al., 1989). Therefore, de novo synthesis of epoxy-carotenoids appears to be unnecessary for ABA synthesis in leaves.

THE CLEAVAGE REACTION

The first committed step in ABA synthesis is the oxidative cleavage of a 9-cis-epoxycarotenoid. For many years, the pathway of ABA synthesis had been a point of contention because of difficulties in demonstrating this activity in vitro. This problem was eventually resolved by the identification and characterization of an ABA-deficient mutant in maize, *vp14* (*viviparous 14*). Biochemical characterization of *vp14* indicated that there was no lesion in carotenoid synthesis or in the later steps of ABA synthesis (Tan et al., 1997). By the process of elimination, it appeared that *vp14* was impaired in the cleavage reaction.

The *vp14* mutant resulted from a transposon insertion, which allowed the corresponding gene to be cloned (Tan et al., 1997). At the time the *Vp14* gene was identified, the deduced amino acid sequence was most similar to lignostilbene dioxygenases (LSDs) from *Pseudomonas (Sphingomonas) paucimobilis*. The LSDs catalyze a double-bond cleavage reaction (Kamoda and Samejima, 1991) that is very similar to the cleavage reaction in ABA synthesis. The recombinant VP14 protein was able to cleave 9-cis-neoxanthin and 9-cis-violaxanthin to form xanthoxin and a C₂₅ by-product (Schwartz et al., 1997b). The characteristics of the cleavage reaction in its substrate specificity and the site of cleavage (11–12 position) were consistent with predictions. A 9-cis double bond in the carotenoid precursor was necessary for activity. The product of this cleavage reaction is cis-xanthoxin, which is readily converted to ABA [cis-(+)-S-ABA] by plants. Cleavage of an all trans-isomer would result in trans-xanthoxin, which is converted to biologically inactive trans-ABA.

Additional ABA synthetic cleavage enzymes have been identified and characterized in a variety of plant species (Qin and Zeevaart, 1999; Chernys and Zeevaart, 2000; Iuchi et al., 2000, 2001). The recombinant enzymes from these species display the same substrate specificity as VP14. The nomenclature that has been adopted for these enzymes is NCEDs,

which is consistent with either 9-cis-violaxanthin or 9-cis-neoxanthin as a substrate.

Although the NCEDs display significant substrate plasticity *in vitro*, circumstantial evidence favors neoxanthin as the primary precursor of ABA. Neoxanthin exists almost entirely as a 9-cis-isomer, whereas only a small proportion of the violaxanthin is present as a 9-cis-isomer (Strand et al., 2000). In addition, the K_m for the recombinant PvNCED1 and VP14 is lower with neoxanthin as substrate relative to 9-cis-violaxanthin (Qin and Zeevaart, 1999; Schwartz et al., 2003). Definitive evidence of the endogenous substrate would require identification of the C_{25} by-product in planta. Previous efforts to identify the C_{25} compounds in vegetative tissue have been unsuccessful. It has been suggested that these compounds are rapidly degraded after the cleavage reaction (Parry and Horgan, 1991). The C_{25} -epoxy-apocarotenal and related compounds have been identified in fruits that produce high levels of ABA during ripening (Molnár and Szabolics, 1980; Gross and Eckhardt, 1981; see also Parry and Horgan, 1991).

Carotenoids in plants are synthesized within plastids and are associated with the thylakoid and envelope membranes. Therefore, it was expected that the cleavage reaction would also occur in chloroplasts. The PvNCED1 from bean (*Phaseolus vulgaris*) was imported into pea (*Pisum sativum*) chloroplasts, where it was found to associate exclusively with the thylakoid membrane (Qin and Zeevaart, 1999). An N-terminal targeting sequence from a cowpea enzyme, VuNCED1, was capable of targeting the green fluorescent protein to chloroplasts (Iuchi et al., 2000). After *in vitro* import assays, the VP14 protein was found in the stroma and on the thylakoid membrane exposed to the stroma (Tan et al., 2001). In this study, deletion or disruption of a putative amphipathic-helix in the N terminus of VP14 interfered with the association of VP14 with thylakoids. The binding of VP14 to the thylakoid was saturable, suggesting that it associates with specific components in the thylakoid membrane that have not yet been identified.

The *not* mutant in tomato also is impaired in the cleavage step (Burbidge et al., 1999). Both the *vp14* and *not* mutants have weak phenotypes relative to other ABA-deficient mutants. The *vp14* null mutant shows only a 35% reduction of ABA levels in stressed leaves and a 70% reduction in developing embryos (Tan et al., 1997). This indicates that there are multiple NCEDs involved in ABA synthesis. In avocado (*Persea americana*), *PaNCED1* and *PaNCED3* were both shown to encode ABA biosynthetic enzymes (Chernys and Zeevaart, 2000).

In the Arabidopsis genome, there are nine hypothetical proteins that share sequence similarity to NCEDs (Fig. 2). In a phylogenetic analysis of the NCEDs and other similar proteins, five of the Arabidopsis proteins are clustered with previously charac-

terized NCEDs. It has been reported that AtNCED2, 3, 6, and 9 are able to catalyze the cleavage reaction in ABA synthesis (Iuchi et al., 2001). Based upon sequence similarity, it is expected that AtNCED5 is also an ABA synthetic enzyme. However, the biochemical role of AtNCED5 has not been verified experimentally.

Presumably, the different NCEDs in Arabidopsis are expressed in different tissues and at different developmental stages. The *AtNCED3* transcript is induced by water stress and reduced expression results in a wilted phenotype (Iuchi et al., 2001), indicating that this gene is an important regulator of ABA levels during water stress. The expression of *AtNCED9* also is elevated slightly in response to water stress (Iuchi et al., 2001). The expression pattern and physiological role of the other *AtNCED* genes has not been reported yet. One or more of these genes is expected to display elevated expression during seed development when ABA begins to accumulate.

SYNTHESIS OF APOCAROTENOIDS

A variety of natural products like ABA are derived from the oxidative cleavage of carotenoids. Collectively referred to as apocarotenoids, these compounds serve important functions in a range of organisms. Retinal is a chromophore used for phototaxis in green algae (Nagel et al., 2002; Sineshchekov et al., 2002) and for light-driven proton pumping in *Halobacterium* NRC-1. (Kolbe et al., 2000).

In animals, vitamin A is necessary for normal vision and development. Based on sequence similarity to NCEDs, a vitamin A biosynthetic enzyme was identified in fruitfly (von Lintig and Vogt, 2000) and subsequently in other species (Wyss et al., 2000; Redmond et al., 2001; Lindqvist and Andersson, 2002). Hypothetical proteins that are similar to NCEDs are also present in a variety of prokaryotes (Fig. 2). The biochemical and biological roles of the putative cleavage enzymes in prokaryotes have not been reported.

All of the proteins in the alignment (Fig. 3) have been shown to catalyze a double-bond cleavage reaction. With the exception of SpLSD, the substrates are carotenoids. The N terminus of the ZmVP14, which does not align well with the other proteins, contains a chloroplast-targeting sequence of approximately 45 amino acids (Tan et al., 2001). The AtCCD7 protein also contains a predicted chloroplast-targeting sequence. The specific functions for conserved residues have not been yet defined. There are several highly conserved His and acidic residues that may be necessary for coordinating iron in the active site.

There are four hypothetical proteins in Arabidopsis that share some degree of sequence similarity with the NCEDs, but are not thought to be involved in ABA synthesis (Fig. 2). Recent findings demonstrate




Figure 2. Unrooted phylogenetic tree of NCEs, carotenoid cleavage dioxygenases (CCDs), and related proteins. The proteins that have been demonstrated to have NCE activity branch together with other likely NCEs. Hypothetical proteins from plants that are not involved in ABA synthesis are referred to as CCDs. Alignments were performed with ClustalW and displayed with TreeView (<http://taxonomy.zoology.gla.ac.uk/rod/treeview.html>). AtCCD1, NP_191911.1; AtCCD4, NP_193652.1; AtCCD7, NP_182026; AtCCD8, NP_195007; AtNCED2, NP_193569.1; AtNCED3, NP_188062.1; AtNCED5, NP_174302.1; AtNCED6, NP_189064; AtNCED9, NP_177960; LeNCED, CAB10168; Md-FS2, CAB07784; OsCCD7, nucleotide sequence 61,958 to 59,544; OsCCD8, BAB63485; PaCCD1, AAK00622.1; PaNCED1, AAK00632.1; PaNCED3, AAK00623; PvCCD1, AAK38744; PvNCED1, AAF26356.1; VuNCED1, BAB11932.1; ZmVP14, AAB62181; Dm β -diox, CAB93141; Mm β -diox I, AAG33982; Mm β -diox II, CAC28026; Mt1, CAB09380; Mt2, CAB08511; Nos1, BAB75983; Nos2, BAB76594; Nos3, BAB73063; SpLSD, AAB35856.2; Syn1, BAA18428; Syn2, BAA18465. At, Arabidopsis; Le, tomato; Md-FS2, *Malus* \times *domestica*; Os, rice; Pa, avocado; Pv, bean; Vu, cowpea; Zm, maize; Dm, fruitfly (*Drosophila melanogaster*); Mm, *Mus musculus*; Mt, *Mycobacterium tuberculosis* H37Rv; Nos, *Nostoc* sp. PCC 7120; Sp, *S. paucimobilis*; Syn, *Synechocystis* sp. PCC 6803.

that at least two of these proteins are able to catalyze carotenoid cleavage reactions. To distinguish these enzymes from the NCEs, the nomenclature of CCDs has been adopted.

The AtCCD1 protein and a likely ortholog in bean catalyze a symmetric 9-10 (9'-10') cleavage with several different carotenoids (Schwartz et al., 2001). A variety of volatile and semivolatile compounds, such as the ionones and β -damascenone, are derived from 9-10 cleavage reactions. These compounds, which are often produced in flowers and fruits, are believed to serve as attractants for pollination and seed dispersal. Several products derived from 9-10, (9'-10') cleavage reactions also accumulate in the roots of plants inoculated with arbuscular mycorrhizal fungi (Walter et al., 2000). The function of these compounds in mycorrhizae has not been determined yet. Recently, a 7-8 (7'-8') CCD has been cloned from *Crocus sativus*, which specifically catalyzes the synthesis of crocetin dialdehyde (C₂₀) and hydroxy- β -

cyclocitral (C₁₀) from zeaxanthin (Bouvier et al., 2003). Crocetin dialdehyde is a precursor of crocin, the primary pigment in saffron.

The most disparate members of this enzyme family in Arabidopsis are AtCCD7 and AtCCD8. The recombinant AtCCD7 protein is able to cleave β , β -carotene at the 9-10 position (S.H. Schwartz and J.A.D. Zeevaart, unpublished data). The AtCCD8 gene corresponds to the *max4* mutant in Arabidopsis, which exhibits extensive lateral shoot growth. The CCD8/MAX4 gene in Arabidopsis is orthologous to the RMS1 gene in pea (O. Leyser, personal communication). Grafting experiments indicate that the *rms1* mutant is impaired in the synthesis of a signal molecule that originates in the wild-type rootstock and inhibits outgrowth of lateral buds on the mutant scion (Beveridge et al., 2000). There is also evidence for a similar signaling molecule in Arabidopsis (Turnbull et al., 2002). The biochemical function and role of AtCCD8 in this process have not been determined yet.

Figure 3. An alignment of several carotenoid cleavage enzymes with ClustalW. Residues that are identical in four or more of the proteins are shaded in black. Similar residues are shaded in gray. GenBank accession numbers and species names can be found in the legend of Figure 2. Conserved His residues are indicated with asterisks.

AtCCD1	-----	0
ZmVP14	MQGLAPPTSVSIHRHLPARSRARASNSVRFSPRAVSSVPPAECLQAPFHKVPADLPAPSRKPAIAVPGH	70
SpLSD	-----	0
Dm β -diox	-----	0
AtCCD7	-----MSLPPI	6
AtCCD1	--MAEKLSDGSI IIVHPRPSKGFSSKLLD LLERLVVKLMH---DASLFLHLYSGNFAPTRD---ETPEV	62
ZmVP14	AAAPRKAEGGKQLNLFQRAAAAALDAFEEGFVANVLERPHGLPSTADPAVQIAGNFAPVG--- ERPEV	136
SpLSD	-----MAHFQPTP---G---FSGTLR---P-LR---IEGDI	23
Dm β -diox	-----MAAGVFKSFMDFFAVKYDEQRNDPQAERLDNGRLYINCSSDVWLRSCER---EIVDP	56
AtCCD7	PKFLPPLKSPPIHHHQTPPLAPPRAAISIPDTGLGRGTILDESSTSAFRDQYSLFVSQRSETIPEV	76
AtCCD1	KDLVHGFTLPE-CLNCFVVRVGENPKFDAVAG-YHFDGDCMHEGVRKDKGKAT--VYSRYVKTSHLKEE	128
ZmVP14	HELVSGRIFPP-FIDCVYARNCANFCDDPVAG-HLFDGDCMVEHALRIRNGAAES-VACRETFETARLRQE	203
SpLSD	LDLIEEGEYPP-QLNCTFHHVHDAQPPRFEDDOFNGDCLVSLFHFHDGKID--FRQYACQDKKQVE	90
Dm β -diox	IEGHSHRTPK-WICGSLRNKGGGSKWVGDMTFGHLDFCSALLHREARNRVPT--VQNRVVDVETLRKN	123
AtCCD7	VIKTEGSLFVNPFSCYYLACGGLFTDDHGSTVPLDGHCHYLRAEHHIDGNRRKATTAARYVTRKAEKE	146
AtCCD1	EFFCAAKFMKIG-DLKGFFGLLMVNIQQLRTRKLIKLDNTYNGTANTALVYHKGQLLALQADKPYVVKV	197
ZmVP14	RATCRPVFPKAIIGELHGHSCLARLALFYARAACGLVDPSPAGVGNAGLYVYNGQLLMSDDLPYHVRV	273
SpLSD	RKAC---KS---LFCAYRNPLTDDASVQGMIR--G---TANINVMVHAGQVYMKRQDSP--- CLLI	141
Dm β -diox	BSAQRIVVTEFG---TAAVDFPCHSIFDRFAALFRPDSGDNDMSLSIYFPGDQYVYFTEPFMHRINP	188
AtCCD7	HDPVTDT-----WRFTHGGFFSVLKGKGRFNTKVMKNVANISVLKWAQCYLCLWEGGEFYEES	206
AtCCD1	LEDGD [*] LTGCIIDYDKRLT-----HSPTAHPKVDVYTC---EMFTFCYSHTP---EYITRVIRVSKDKIMH	256
ZmVP14	ADDGDELTVGRYDFDQGLG-----CAMIAHPKVDVYTC---ELHALSYDVTKR---EMLKPYFRPDKKS	333
SpLSD	MDPLTLETECYTDFDGLQS-----QTFCAHPKVDVYTCNLCFAFYAKKLT-LDMAHIESPTCKLL	204
Dm β -diox	CTLAEARICTDDEVVVM-----HTSHPHVLSGT---VYINLCTMTIRSGEAYTILCFPFGEQMF	246
AtCCD7	GSLDVGGRFVNEVNGCESDDDDSSDRDLSGEDIWDTAADLLKPILOCVFKMPPKRRHSYKWDGRRKRL	276
AtCCD1	DPVPTISEPI-----MMDHFAITVYALFMDLDMHFRPKMVKKEK	297
ZmVP14	DDVEIPLQPT-----MMDHFAITVNFVVPDHOVVKLQDMLRGG	374
SpLSD	KEIFPNQYYC-----MMDHFGVEDVAVFAVMSLSS-WDRLEQR	244
Dm β -diox	EDAHVVATLDCRWK-----LHPGYMHTGCTDHYVYVTECELSVSLTEVYKAO	294
AtCCD7	LTYFCNAEDMLLRSNFTFCEYDSEFKLIQTKFKIDDHMMHDFAPDTHYILFANRVKLNPIGSIAM	346
AtCCD1	K-----MIYSFDPTKK-ARFGLPRY-----AKDELMIREFELPNCIFIFNANAWEEDEYVLTICRL	354
ZmVP14	-----SPVVLDKERT-SREGLPKH-----AADASEMAVVDVDCFCFHUNAWAEDAEATGEVHVIVGS	430
SpLSD	L-----PFFGFDITLP-CYLEILR-----NGDARDLRVFKTGNCFVGHVMAFNEDGTRVHDMFVS	300
Dm β -diox	LGGQNLACLKWFEDRP-TLGHLEDR-----VSGKLVQVYSEAFVYHLINCCERDGHVVDICSY	355
AtCCD7	CGMSPMVSALSLNPSNESSPIYLFRFSDKYSRGRDRWVPEVSSQLWLIHSGANFMTREDNGDLKQI	416
AtCCD1	ENP-----	357
ZmVP14	CMT-----	433
SpLSD	RNN-----	303
Dm β -diox	RNPENINCMYLEAIANMQTNPYATLFRGRPLRFVPLPGTIPASIAKRGVKSFSLAGLSAPQVSRMTK	425
AtCCD7	QASACSRYWDFQKMGF-----	433
AtCCD1	-----DLDMVSGVKVEKLENFCNEI-YEMFRNMGSAQSOKKLSASA-----VDFEPIINE	406
ZmVP14	-----PADSIFNESDERLESVLTEIRLDARTGRSARRAVLPPSQQEN-----LEVGMAR	483
SpLSD	-----SFPFFDVHGAPFDPVAGGFLTRWTVDMASNGDSFEKTERLDFDR--PDEFERDE	356
Dm β -diox	HSVSYADITYMPTNGKQATAGEESPKRDAKRGRYEENLVNLMVMEGSAQAEAFQGTNG--IQLRPEMLC	493
AtCCD7	-----YDQSNKLDPSVMNLRNGDDKLLPHLVKVSMTLDSGNCNSCDVPELNGWKNKSPDFEIVNS	494
AtCCD1	CYTCKKORYVGTILDSIAKVTGIIKFDLHAEAE[*]TGKRMLEVEGGNIKIGYIDLGEGRYSSAIVVRETAE	476
ZmVP14	NLLCRESRVAMLA[*]VAPWPKESSGFAKEDLS---TCELT-----KEFYEGREGGEPFCVFMDDPAA	540
SpLSD	RYATRAYRGGWMLLDTEKPYEAPGGAFYALNTLGHIDLATGK---SSSNWAGPRCAIQEPPCTIRSPDA	424
Dm β -diox	DWGCETPRLYRMYGKNRYRYFYAISSDVDVN-FCTLIKVDNMKSLC[*]TWCEENVYSPRPIEVS[*]SPDK	562
AtCCD7	SWSGCKNRYM[*]SAASSGTRSELPHFPDMVVKFDLDSNLVR-----TNSTGARRVCGEPMVEKNSVE	557
AtCCD1	-----EDDGYLFFVHD--ENTGKSCVTVIDAKTMSAEPVAVVELFHRVYEGFHALVTEEQIQEQTLI	538
ZmVP14	AHPRGEDDGYLTFVHD--ERACTSELVAVNADIRLE--AVVQLFSRVEFSEHCTETGQEAEQA--	604
SpLSD	F---EGDGYVIALVON--HVANYSDIAIFDQHVDQGGPI[*]TRAKLEVRERQCHCNWADASRLAVAA--	485
Dm β -diox	S---EDDGVILASMLGGLNDRYVGLVIVCAKMTLGRCDFTNGPVEKCLHGWAPAPNAI-----	620
AtCCD7	EGEE-EDDGYVVVVEYA--VSVRCYMLVILDARKIGESDAVVSRETEVPRNLTPMGEHGLWASD----	618

THE LATER STEPS. CONVERSION OF XANTHOXIN TO ABA

In contrast to the cleavage reaction, the later steps in ABA synthesis have been characterized extensively by feeding potential intermediates to intact plants and cell-free extracts (Sindhu and Walton, 1987). By this approach, the sequence of reactions subsequent to cleavage has been determined. The first steps are the oxidation of the 4'-hydroxyl to a ketone followed by the nonenzymatic desaturation of 2'-3' bond and opening of the epoxide ring. The final step in the pathway is the oxidation of abscisic aldehyde to ABA.

The *aba2* mutant in Arabidopsis was first identified by screening for the ability to germinate in the presence of the GA biosynthetic inhibitor, paclobutrazol (Léon-Kloosterziel et al., 1996). By feeding potential intermediates to extracts of the *aba2* mutant, it was determined that this mutant was impaired in the

conversion of xanthoxin to abscisic aldehyde (Schwartz et al., 1997a). The *aba2* is the only mutant identified to date that is blocked at this step in the pathway. Additional alleles of *aba2* have since been identified in screens for a sugar-insensitive phenotype (Laby et al., 2000; Cheng et al., 2002; see Table I), altered stomatal conductance (Merlot et al., 2002), and germination and growth on a medium containing a high NaCl concentration (González-Guzmán et al., 2002). The gene corresponding to *aba2* has recently been cloned and the gene product was found to be similar to short chain dehydrogenases/reductases (Cheng et al., 2002; González-Guzmán et al., 2002). As expected, the ABA2 protein was able to catalyze the conversion of xanthoxin to abscisic aldehyde utilizing NAD as a cofactor. The ABA2 transcript level was not affected by stress (González-Guzmán et al., 2002) but was induced by Glc (Cheng et al., 2002). It has not yet been reported whether the

overexpression of *ABA2* would have any effect on ABA levels.

Mutants impaired in the final step of ABA synthesis, the oxidation of abscisic aldehyde to ABA, have been identified in a variety of plants. A loss of this abscisic aldehyde oxidase activity may result from a mutation in the aldehyde oxidase apoprotein or a lesion in the synthesis of a MoCo that the enzyme requires for activity. A lesion in an early step of MoCo synthesis would affect a number of activities. For example, the *nar2a* mutant in barley lacks aldehyde oxidase, xanthine dehydrogenase, and nitrate reductase activities (Walker-Simmons et al., 1989). The *aba3* mutant in Arabidopsis and the *flacca* mutant in tomato lack aldehyde oxidase activity, but the activity of nitrate reductase is unaffected. This phenotype results from a defect in the formation of a desulfo moiety of the MoCo that is specifically required by certain hydroxylases (Schwartz et al., 1997a; Akaba et al., 1998; Sagi et al., 1999). The gene corresponding to *aba3* has been cloned and the N terminus of the deduced protein is similar to the NifS sulfurase (Bittner et al., 2001; Xiong et al., 2001). Using Cys as a sulfur donor, the recombinant protein was able to activate aldehyde oxidase activity (Bittner et al., 2001). The *flacca* mutant also results from a mutation in this sulfurase (Sagi et al., 1999, 2002). It was found that *ABA3* expression increased in response to osmotic stress or ABA (Bittner et al., 2001; Xiong et al., 2001).

Four abscisic aldehyde oxidase (AAO) genes have been identified in Arabidopsis (*AAO1* through *4*). Of the aldehyde oxidases characterized so far, only *AAO3* uses abscisic aldehyde efficiently as a substrate (Seo et al., 2000a). A wilted, ABA-deficient mutant with a lesion in *AAO3* has been identified (Seo et al., 2000b), demonstrating that *AAO3* is responsible for ABA synthesis in vegetative tissues. In contrast to *aba3*, the *ao3* mutants are not subject to precocious germination. Therefore, another aldehyde oxidase appears to be necessary for ABA synthesis in some tissues. In plants subjected to dehydration, *AAO3* mRNA expression was elevated. However, the level of the corresponding protein was unaffected by water stress (Seo et al., 2000a). In addition, feeding experiments and assays with cell-free preparations indicate that the conversion of xanthoxin to ABA is unaffected by stress (Sindhu and Walton, 1987; Schwartz et al., 1997a).

Several variations in the later steps of the pathway may be responsible for a small portion of ABA synthesis. It has been suggested that oxidation of the aldehyde may occur subsequent to cleavage and before the ring modifications (Cowan, 2000), indicating that xanthoxic acid would be an intermediate in the pathway. However, the conversion of xanthoxic acid is very low in cell-free extracts (Sindhu and Walton, 1987). Also, the *ABA2* protein is unable to convert xanthoxic acid to ABA (Cheng et al., 2002). Both the

flacca and *sitiens* mutants in tomato are blocked in the final step of the pathway, the oxidation of abscisic aldehyde to ABA, and accumulate 2-trans-ABA alcohol (Linforth et al., 1987). These mutants are able to synthesize some ABA by a shunt pathway in which abscisic alcohol is oxidized to ABA (Rock et al., 1991b).

TRANSGENIC PLANTS WITH ELEVATED ABA LEVELS

Inhibitors of transcription and translation block stress-induced ABA accumulation (Quarrie and Lister, 1984; Guerrero and Mullet, 1986), indicating gene expression is up-regulated for one or more steps in ABA synthesis. The genes encoding most of the enzymes in the ABA biosynthetic pathway have now been identified. Based upon elevated expression during stress, a regulatory role has been proposed for several of the genes (Audran et al., 1998, 2001; Seo et al., 2000b; Bittner et al., 2001; Xiong et al., 2001, 2002).

In etiolated tissues, the levels of epoxy-carotenoids are low and it appears that elevated expression of ZEP may be important for ABA synthesis (Frey et al., 1999). In green tissue, however, most of the biochemical evidence indicates that the NCED-catalyzed cleavage reaction is the primary regulatory step in ABA synthesis. In all instances studied to date, stress-induced ABA accumulation correlates well with increased expression of *NCED* mRNA (Tan et al., 1997; Qin and Zeevaart, 1999; Chernys and Zeevaart, 2000; Iuchi et al., 2000, 2001; Thompson et al., 2000a) and also with *NCED* protein levels (Qin and Zeevaart, 1999). The expression of *PanNCED1* and *PanNCED3* also increased before the accumulation of high ABA levels during fruit ripening in avocado (Chernys and Zeevaart, 2000).

The overexpression of *NCEDs* is sufficient for elevated ABA synthesis. Overexpression of the *LeNCED1* in tomato (Thompson et al., 2000b), *PvNCED1* in tobacco (Qin and Zeevaart, 2002), and *AtNCED3* in Arabidopsis (Iuchi et al., 2001) resulted in increased ABA levels. In *N. plumbaginifolia*, induced expression of *PvNCED1* resulted in decreased stomatal conductance and increased stress tolerance (Fig. 4). Similar results were obtained with the other species listed above. For the overexpression of *LeNCED1* (Thompson et al., 2000b) and *PvNCED1* (Qin and Zeevaart, 2002), increased seed dormancy was also reported.

CATABOLISM OF ABA

The level of ABA in plants is controlled not only by its synthesis, but also through its catabolism. One of the primary catabolites of ABA is phaseic acid (PA). The conversion of ABA to PA begins with the hydroxylation of the 8' position by ABA 8'-hydroxylase. The 8' hydroxyl appears to be an unstable intermediate that spontaneously rearranges to form




Figure 4. Enhanced drought tolerance and recovery of *N. plumbaginifolia* plants overexpressing *PvNCED1*. A, Homozygous plants transformed with vector only (*GVG*) or with *PvNCED1* gene under control of a dexamethasone-inducible promoter *GVG* 7 d after spraying with water or 30 μM dexamethasone (DEX), and without watering. B, Difference in recovery of control and plant overexpressing *PvNCED1* 9 d after daily watering was resumed.

PA. The ABA 8'-hydroxylase is a cytochrome P450 (Krochko et al., 1998), which may be induced by ABA (Windsor and Zeevaart, 1997). This negative feedback regulation is consistent with time course measurements of ABA and PA accumulation in stressed plants (Zeevaart, 1980) and recent work with *NCED* overexpression in plants (Qin and Zeevaart, 2002). ABA may also be inactivated by the formation of ABA Glc ester in some tissues. An ABA glucosyltransferase gene from adzuki bean (*Vigna angularis*) has been cloned recently (Xu et al., 2002). Interestingly, this gene is also up-regulated by ABA. The physiological significance of ABA Glc ester formation and the potential for engineering ABA levels by decreased glucosylation may now be investigated.

FUTURE DIRECTIONS

There are several steps in ABA biosynthesis preceding the cleavage reaction that are not well characterized. The epoxy-carotenoid precursor must have a 9-cis configuration to be cleaved by an *NCED* and for subsequent conversion to ABA [cis-(+)-S-ABA]. The formation of these 9-cis isomers has not yet been established. An enzyme that catalyzes a similar reaction, the cis/trans isomerization of prolycopene to lycopene, has recently been identified (Isaacson et al., 2002; Park et al., 2002). This isomerase appears to be

necessary only in non-photosynthetic tissue. In light-grown tissue, photo-isomerization of lycopene is sufficient. It has not been established whether the 9-cis isomerization of neoxanthin and violaxanthin is an enzymatic reaction. Alternatively, the 9-cis conformations of some epoxy-carotenoids could be stabilized by carotenoid-binding proteins.

In most plant tissues, neoxanthin is the predominant carotenoid with a 9-cis conformation and is considered the most likely precursor of ABA (Strand et al., 2000). Neoxanthin is derived through the opening of an epoxy ring in violaxanthin followed by an intramolecular rearrangement to form an allenic bond. Allenic carotenoids, such as neoxanthin, are among the most abundant carotenoids in nature. Therefore, an understanding of their synthesis and functions in photosynthetic organisms is of considerable interest. Two genes that encode neoxanthin synthases (*NSY*) have been identified in potato (*Solanum tuberosum*) and tomato (Al-Babili et al., 2000; Bouvier et al., 2000). The *NSY* gene products are similar to lycopene cyclases from various plants and a capsanthin-capsorubin synthase from pepper (*Capsicum annuum*). Transient expression in tobacco and in vitro assays both demonstrated that the tomato *NSY* was capable of converting violaxanthin to neoxanthin (Bouvier et al., 2000). No lycopene cyclase activity was found by co-expression in a lycopene-accumulating strain of *Escherichia coli* (Bouvier et al., 2000). However, the *NSY* gene corresponds to the *old-gold* mutant in tomato, which accumulates higher levels of lycopene due to the loss of a fruit-specific lycopene β -cyclase, *CYC-B* (Ronen et al., 2000; Hirschberg, 2001). It has been suggested that the *NSY* is a bifunctional enzyme capable of converting lycopene to β , β -carotene or violaxanthin to neoxanthin (Hirschberg, 2001). Presumably, there is an additional gene responsible for neoxanthin synthesis in plants, because the *old-gold* mutant is able to produce neoxanthin. Moreover, no ortholog of the *NSY* gene is apparent in the Arabidopsis genome.

The oxidative cleavage products of carotenoids serve important roles in both plants and animals. Based upon sequence similarity to *NCEDs*, putative cleavage enzymes have been identified in a number of plants and prokaryotes. The characterization of *CCDs* in plants suggests that apocarotenoids have various roles in growth and development. The synthesis of apocarotenoids is well documented in cyanobacteria and a carotenoid cleavage activity has been described in the cyanobacterium *Microcystis PCC7806* (Jüttner and Höflacher, 1985). However, the biological functions of these compounds in cyanobacteria and other prokaryotes have not yet been determined.

Despite the important roles that apocarotenoids serve in various organisms and the growing number of putative cleavage enzymes appearing in the sequence databases, little is known about the mecha-

nism by which these enzymes catalyze reactions. In an isotopic labeling experiment with β,β -carotene 15, 15'-dioxygenase from chicken (*Gallus gallus*), approximately 50% of the cleavage products contained oxygen derived from O_2 (Leuenberger et al., 2001). In this experiment, the second oxygen was derived from water, and the authors proposed a monooxygenase mechanism. However, no reducing equivalents are required for assays with any of the recombinant enzymes that have been characterized. In addition, $^{18}O_2$ -labeling experiments with plants indicate that the initial cleavage product in ABA synthesis, xanthoxin, results entirely from O_2 (Zeevaert et al., 1989). At this point, the mechanism by which these enzymes catalyze reactions is still uncertain.

The biochemical aspects of ABA synthesis, such as the intermediates in the pathway and the sequence of reactions, have become well established. The genes that encode most of the enzymes in the pathway have now been cloned. Although elevated expression in response to osmotic stress has been reported for several of these genes, the significance of this up-regulation is still uncertain. Previous biochemical studies and the recent work with transgenic plants clearly demonstrate that transcriptional regulation of the *NCEDs* is the major control point in ABA synthesis. The initial perception of stress and the signal transduction pathway leading to elevated *NCED* expression remain to be elucidated.

ACKNOWLEDGMENTS

We thank Dr. Nam-Hai Chua (Rockefeller University, New York) for providing the pTA7002 plasmid and Dr. Ottoline Leyser (University of York, UK) for sharing her unpublished results with the *atccd8/max4* mutant.

Received November 19, 2002; returned for revision December 9, 2002; accepted January 13, 2003.

LITERATURE CITED

- Agrawal GK, Yamazaki M, Kobayashi M, Hirochika R, Miyao A, Hirochika H (2001) Screening of the rice viviparous mutants generated by endogenous retrotransposon *Tos17* insertion: tagging of a zeaxanthin epoxidase gene and a novel *OsTATC* gene. *Plant Physiol* **125**: 1248–1257
- Akaba S, Leydecker MT, Moureaux T, Oritani T, Koshiba T (1998) Aldehyde oxidase in wild type and *aba1* mutant leaves of *Nicotiana plumbaginifolia*. *Plant Cell Physiol* **39**: 1281–1286
- Al-Babili S, Huguency P, Schledz M, Welsch R, Frohnmeyer H, Laule O, Beyer P (2000) Identification of a novel gene coding for neoxanthin synthase from *Solanum tuberosum*. *FEBS Lett* **485**: 168–172
- Audran C, Borel C, Frey A, Sotta B, Meyer C, Simonneau T, Marion-Poll A (1998) Expression studies of the zeaxanthin epoxidase gene in *Nicotiana plumbaginifolia*. *Plant Physiol* **118**: 1021–1028
- Audran C, Liotenberg S, Gonneau M, North H, Frey A, Tap-Waksman K, Vartanian N, Marion-Poll A (2001) Localisation and expression of zeaxanthin epoxidase mRNA in *Arabidopsis* in response to drought stress and during seed development. *Aust J Plant Physiol* **28**: 1161–1173
- Beveridge CA, Symons GM, Turnbull CGN (2000) Auxin inhibition of decapitation-induced branching is dependent on graft-transmissible signals regulated by genes *Rms1* and *Rms2*. *Plant Physiol* **123**: 689–697
- Bittner F, Oreb M, Mendel RR (2001) ABA3 is a molybdenum cofactor sulfurase required for activation of aldehyde oxidase and xanthine dehydrogenase in *Arabidopsis thaliana*. *J Biol Chem* **276**: 40381–40384
- Borel C, Audran C, Frey A, Marion-Poll A, Tardieu F, Simonneau T (2001) *N. plumbaginifolia* zeaxanthin epoxidase transgenic lines have unaltered baseline ABA accumulations in roots and xylem sap, but contrasting sensitivities of ABA accumulation to water deficit. *J Exp Bot* **52**: 427–434
- Bouvier F, D'Harlingue A, Backhaus RA, Kumagai MH, Camara B (2000) Identification of neoxanthin synthase as a carotenoid cyclase paralog. *Eur J Biochem* **267**: 6346–6352
- Bouvier F, d'Harlingue A, Huguency P, Marin E, Marion-Poll A, Camara B (1996) Xanthophyll biosynthesis: cloning, expression, functional reconstitution, and regulation of β -cyclohexenyl carotenoid epoxidase from pepper (*Capsicum annuum*). *J Biol Chem* **271**: 28861–28867
- Bouvier F, Suire C, Mutterer J, Camara B (2003) Oxidative remodeling of chromoplast carotenoids: identification of the carotenoid dioxygenase *CsCCD* and *CsZCD* genes involved in crocus secondary metabolite biogenesis. *Plant Cell* **15**: 47–62
- Burbidge A, Grieve TM, Jackson A, Thompson A, McCarty DR, Taylor IB (1999) Characterization of the ABA-deficient tomato mutant *notabilis* and its relationship with maize *Vp14*. *Plant J* **17**: 427–431
- Cheng W-H, Endo A, Zhou L, Penney J, Chen H-C, Arroyo A, Leon P, Nambara E, Asami T, Seo M et al. (2002) A unique short-chain dehydrogenase/reductase in *Arabidopsis* glucose signaling and abscisic acid biosynthesis and functions. *Plant Cell* **14**: 2723–2743
- Chernys JT, Zeevaert JAD (2000) Characterization of the 9-*cis*-epoxycarotenoid dioxygenase gene family and the regulation of abscisic acid biosynthesis in avocado. *Plant Physiol* **124**: 343–353
- Cowan AK (2000) Is abscisic aldehyde really the immediate precursor to stress-induced ABA? *Trends Plant Sci* **5**: 191–192
- Cunningham FX Jr, Gantt E (1998) Genes and enzymes of carotenoid biosynthesis in plants. *Annu Rev Plant Physiol Plant Mol Biol* **49**: 557–583
- Duckham SC, Linforth RST, Taylor IB (1991) Abscisic-acid-deficient mutants at the *aba* gene locus of *Arabidopsis thaliana* are impaired in the epoxidation of zeaxanthin. *Plant Cell Environ* **14**: 601–606
- Frey A, Audran C, Marin E, Sotta B, Marion-Poll A (1999) Engineering seed dormancy by the modification of zeaxanthin epoxidase gene expression. *Plant Mol Biol* **39**: 1267–1274
- González-Guzmán M, Apostolova N, Bellés JM, Barrero JM, Piqueras P, Ponce MR, Micol JL, Serrano R, Rodríguez PL (2002) The short-chain alcohol dehydrogenase ABA2 catalyzes the conversion of xanthoxin to abscisic aldehyde. *Plant Cell* **14**: 1833–1846
- Gross J, Eckhardt G (1981) Structures of persicaxanthin, persicachrome and other apocarotenols of various fruits. *Phytochemistry* **20**: 2267–2269
- Guerrero F, Mullet JE (1986) Increased abscisic acid biosynthesis during plant dehydration requires transcription. *Plant Physiol* **80**: 588–591
- Hirai N, Yoshida R, Todoroki Y, Ohigashi H (2000) Biosynthesis of abscisic acid by the non-mevalonate pathway in plants, and by the mevalonate pathway in fungi. *Biosci Biotechnol Biochem* **64**: 1448–1458
- Hirschberg J (2001) Carotenoid biosynthesis in flowering plants. *Curr Opin Plant Biol* **4**: 210–218
- Isaacson T, Ronen G, Zamir D, Hirschberg J (2002) Cloning of *tangerine* from tomato reveals a carotenoid isomerase essential for the production of β -carotene and xanthophylls in plants. *Plant Cell* **14**: 333–342
- Iuchi S, Kobayashi M, Taji T, Naramoto M, Seki M, Kato T, Tabata S, Kakubari Y, Yamaguchi-Shinozaki K, Shinozaki K (2001) Regulation of drought tolerance by gene manipulation of 9-*cis*-epoxycarotenoid dioxygenase, a key enzyme in abscisic acid biosynthesis in *Arabidopsis*. *Plant J* **27**: 325–333
- Iuchi S, Kobayashi M, Yamaguchi-Shinozaki K, Shinozaki K (2000) A stress-inducible gene for 9-*cis*-epoxycarotenoid dioxygenase involved in abscisic acid biosynthesis under water stress in drought-tolerant cowpea. *Plant Physiol* **123**: 553–562
- Jüttner F, Höflacher B (1985) Evidence of β -carotene 7,8 (7',8') oxygenase (β -cyclocitral, crocetindial generating) in *Microcystis*. *Arch Microbiol* **141**: 337–343
- Kamoda S, Samejima M (1991) Cloning of a lignostilbene- α,β -dioxygenase gene from *Pseudomonas paucimobilis* TMY1009. *Agric Biol Chem* **55**: 1411–1412
- Karssen CM, Brinkhorst-van der Swan DLC, Breeklund AE, Koornneef M (1983) Induction of dormancy during seed development by endogenous

- abscisic acid: studies on abscisic acid deficient genotypes of *Arabidopsis thaliana* (L.) Heynh. *Planta* **157**: 158–165
- Kolbe M, Besir H, Essen LO, Oesterhelt D** (2000) Structure of the light-driven chloride pump halorhodopsin at 1.8 Å resolution. *Science* **288**: 1390–1396
- Krochko JE, Abrams GD, Loewen MK, Abrams SR, Cutler AJ** (1998) (+)-Abscisic acid 8'-hydroxylase is a cytochrome P450 monooxygenase. *Plant Physiol* **118**: 849–860
- Laby RJ, Kincaid MS, Kim DG, Gibson SI** (2000) The *Arabidopsis* sugar-insensitive mutants *sis4* and *sis5* are defective in abscisic acid synthesis and response. *Plant J* **23**: 587–596
- Léon-Kloosterziel KM, Alvarez-Gil M, Ruijs GJ, Jacobsen SE, Olszewski NE, Schwartz SH, Zeevaart JAD, Koornneef M** (1996) Isolation and characterization of abscisic acid-deficient *Arabidopsis* mutants at two new loci. *Plant J* **10**: 655–661
- Leuenberger MG, Engeloch-Jarret C, Woggon WD** (2001) The reaction mechanism of the enzyme-catalyzed central cleavage of β -carotene to retinal. *Angew Chem Int Ed* **40**: 2614–2617
- Lindqvist A, Andersson S** (2002) Biochemical properties of purified recombinant human β -carotene 15,15'-monooxygenase. *J Biol Chem* **277**: 23942–23948
- Linforth RST, Bowman WR, Griffin DA, Marples BA, Taylor IB** (1987) 2-trans-ABA alcohol accumulation in the wilted tomato mutants *flacca* and *sitiens*. *Plant Cell Environ* **10**: 599–606
- Lokstein H, Tian L, Polle JEW, DellaPenna D** (2002) Xanthophyll biosynthetic mutants of *Arabidopsis thaliana*: altered nonphotochemical quenching of chlorophyll fluorescence is due to changes in photosystem II antenna size and stability. *Biochim Biophys Acta* **1553**: 309–319
- Marin E, Nussaume L, Quesada A, Gonneau M, Sotta B, Huguency P, Frey A, Marion-Poll A** (1996) Molecular identification of zeaxanthin epoxidase of *Nicotiana plumbaginifolia*, a gene involved in abscisic acid biosynthesis and corresponding to the *ABA* locus of *Arabidopsis thaliana*. *EMBO J* **15**: 2331–2342
- Merlot S, Mustilli AC, Genty B, North H, Lefebvre V, Sotta B, Vavasseur A, Giraudat J** (2002) Use of infrared thermal imaging to isolate *Arabidopsis* mutants defective in stomatal regulation. *Plant J* **30**: 601–609
- Molnár P, Szabolics J** (1980) β -Citaurin epoxide, a new carotenoid from Valencia orange peel. *Phytochemistry* **19**: 633–637
- Nagel G, Ollig D, Fuhrmann M, Kateriya S, Musti AM, Bamberg E, Hegemann P** (2002) Channel rhodopsin-1: a light-gated proton channel in green algae. *Science* **296**: 2395–2398
- Niyogi KK, Grossman AR, Björkman O** (1998) *Arabidopsis* mutants define a central role for the xanthophyll cycle in the regulation of photosynthetic energy conversion. *Plant Cell* **10**: 1121–1134
- Okamoto M, Min X, Seo M, Nakabayashi K, Kamiya Y, Nambara E, Koshiba T** (2002) Complementation of a tomato ABA-deficient *sitiens* mutant by an *Arabidopsis* aldehyde oxidase gene, *AAO3*. *Plant Cell Physiol* **43**: S42
- Park H, Kreunen SS, Cuttriss AJ, DellaPenna D, Pogson BJ** (2002) Identification of the carotenoid isomerase provides insight into carotenoid biosynthesis, prolamellar body formation, and photomorphogenesis. *Plant Cell* **14**: 321–332
- Parry AD, Horgan R** (1991) Carotenoid metabolism and the biosynthesis of abscisic acid. *Phytochemistry* **30**: 815–821
- Qin X, Zeevaart JAD** (1999) The 9-*cis*-epoxycarotenoid cleavage reaction is the key regulatory step of abscisic acid biosynthesis in water-stressed bean. *Proc Natl Acad Sci USA* **96**: 15354–15361
- Qin X, Zeevaart JAD** (2002) Overexpression of a 9-*cis*-epoxycarotenoid dioxygenase gene in *Nicotiana plumbaginifolia* increases abscisic acid and phaseic acid levels and enhances drought tolerance. *Plant Physiol* **128**: 544–551
- Quarrie SA, Lister PG** (1984) Effects of inhibitors of protein synthesis on abscisic acid accumulation in wheat. *Z Pflanzenphysiol* **114**: 309–314
- Redmond TM, Gentleman S, Duncan T, Yu S, Wiggert B, Gantt E, Cunningham FX** (2001) Identification, expression, and substrate specificity of a mammalian β -carotene 15,15'-dioxygenase. *J Biol Chem* **276**: 6560–6565
- Rock CD, Heath TG, Gage DA, Zeevaart JAD** (1991b) Abscisic alcohol is an intermediate in abscisic acid biosynthesis in a shunt pathway from abscisic aldehyde. *Plant Physiol* **97**: 670–676
- Rock CD, Zeevaart JAD** (1991a) The *aba* mutant of *Arabidopsis thaliana* is impaired in epoxy-carotenoid biosynthesis. *Proc Natl Acad Sci USA* **88**: 7496–7499
- Rodríguez-Concepción M, Boronat A** (2002) Elucidation of the methylerythritol phosphate pathway for isoprenoid biosynthesis in bacteria and plastids. A metabolic milestone achieved through genomics. *Plant Physiol* **130**: 1079–1089
- Rohmer M** (1999) The discovery of a mevalonate-independent pathway for isoprenoid biosynthesis in bacteria, algae and higher plants. *Nat Prod Rep* **16**: 565–574
- Ronen G, Carmel-Goren L, Zamir D, Hirschberg J** (2000) An alternative pathway to β -carotene formation in plant chromoplasts discovered by map-based cloning of *Beta* and *old-gold* color mutations in tomato. *Proc Natl Acad Sci USA* **97**: 11102–11107
- Rook F, Corke F, Card R, Munz G, Smith C, Bevan MW** (2001) Impaired sucrose-induction mutants reveal the modulation of sugar-induced starch biosynthetic gene expression by abscisic acid signalling. *Plant J* **26**: 421–433
- Sagi M, Fluhr R, Lips SH** (1999) Aldehyde oxidase and xanthine dehydrogenase in a *flacca* tomato mutant with deficient abscisic acid and wilted phenotype. *Plant Physiol* **120**: 571–577
- Sagi M, Scaccocchio C, Fluhr R** (2002) The absence of molybdenum cofactor sulfuration is the primary cause of the *flacca* phenotype in tomato plants. *Plant J* **31**: 305–317
- Schwartz SH, Léon-Kloosterziel KM, Koornneef M, Zeevaart JAD** (1997a) Biochemical characterization of the *aba2* and *aba3* mutants in *Arabidopsis thaliana*. *Plant Physiol* **114**: 161–166
- Schwartz SH, Qin XQ, Zeevaart JAD** (2001) Characterization of a novel carotenoid cleavage dioxygenase from plants. *J Biol Chem* **276**: 25208–25211
- Schwartz SH, Tan BC, Gage DA, Zeevaart JAD, McCarty DR** (1997b) Specific oxidative cleavage of carotenoids by VP14 of maize. *Science* **276**: 1872–1874
- Schwartz SH, Tan BC, McCarty DR, Welch W, Zeevaart JAD** (2003) Substrate specificity and kinetics for VP14, a carotenoid cleavage dioxygenase in the ABA biosynthetic pathway. *Biochim Biophys Acta* **1619**: 9–14
- Seo M, Koiwai H, Akaba S, Komano T, Oritani T, Kamiya Y, Koshiba T** (2000a) Abscisic aldehyde oxidase in leaves of *Arabidopsis thaliana*. *Plant J* **23**: 481–488
- Seo M, Peeters AJM, Koiwai H, Oritani T, Marion-Poll A, Zeevaart JAD, Koornneef M, Kamiya Y, Koshiba T** (2000b) The *Arabidopsis* aldehyde oxidase 3 (*AAO3*) gene product catalyzes the final step in abscisic acid biosynthesis in leaves. *Proc Natl Acad Sci USA* **97**: 12908–12913
- Sindhu RK, Walton DC** (1987) Conversion of xanthoxin to abscisic acid by cell-free preparations from bean leaves. *Plant Physiol* **85**: 916–921
- Sineshchekov OA, Jung KH, Spudich JL** (2002) Two rhodopsins mediate phototaxis to low- and high-intensity light in *Chlamydomonas reinhardtii*. *Proc Natl Acad Sci USA* **99**: 8689–8694
- Strand A, Kvernberg K, Karlsen AM, Liaaen-Jensen S** (2000) Geometrical *E/Z* isomers of (6*R*)- and (6*S*)-neoxanthin and biological implications. *Biochem Syst Ecol* **28**: 443–455
- Tan BC, Cline K, McCarty DR** (2001) Localization and targeting of the VP14 epoxycarotenoid dioxygenase to chloroplast membranes. *Plant J* **27**: 373–382
- Tan BC, Schwartz SH, Zeevaart JAD, McCarty DR** (1997) Genetic control of abscisic acid biosynthesis in maize. *Proc Natl Acad Sci USA* **94**: 12235–12240
- Taylor HF, Smith TA** (1967) Production of plant growth inhibitors from xanthophylls: a possible source of dormin. *Nature* **215**: 1513–1514
- Thompson AJ, Jackson AC, Parker RA, Morpeth DR, Burbidge A, Taylor IB** (2000a) Abscisic acid biosynthesis in tomato: regulation of zeaxanthin epoxidase and 9-*cis*-epoxycarotenoid dioxygenase mRNAs by light/dark cycles, water stress and abscisic acid. *Plant Mol Biol* **42**: 833–845
- Thompson AJ, Jackson AC, Symonds RC, Mulholland BJ, Dadswell AR, Blake PS, Burbidge A, Taylor IB** (2000b) Ectopic expression of a tomato 9-*cis*-epoxycarotenoid dioxygenase gene causes over-production of abscisic acid. *Plant J* **23**: 363–374
- Turnbull CGN, Booker JP, Leyser HM** (2002) Micrografting techniques for testing long-distance signalling in *Arabidopsis*. *Plant J* **32**: 255–262
- von Lintig J, Vogt K** (2000) Filling the gap in vitamin A research: molecular identification of an enzyme cleaving β -carotene to retinal. *J Biol Chem* **275**: 11915–11920
- Walker-Simmons M, Kudrna DA, Warner RL** (1989) Reduced accumulation

- of ABA during water stress in a molybdenum cofactor mutant of barley. *Plant Physiol* **90**: 728–733
- Walter MH, Fester T, Strack D** (2000) Arbuscular mycorrhizal fungi induce the non-mevalonate methylerythritol phosphate pathway of isoprenoid biosynthesis correlated with accumulation of the “yellow pigment” and other apocarotenoids. *Plant J* **21**: 571–578
- Windsor ML, Zeevaert JA** (1997) Induction of ABA 8'-hydroxylase by (+)-S-,(-)-R- and 8'-8'-8'-trifluoro-S-abscisic acid in suspension cultures of potato and *Arabidopsis*. *Phytochemistry* **45**: 931–934
- Wyss A, Wirtz G, Woggon WD, Brugger R, Wyss M, Friedlein A, Bachmann H, Hunziker W** (2000) Cloning and expression of β,β -carotene 15,15'-dioxygenase. *Biochem Biophys Res Commun* **271**: 334–336
- Xiong L, Ishitani M, Lee H, Zhu JK** (2001) The *Arabidopsis* *LOS5/ABA3* locus encodes a molybdenum cofactor sulfuryase and modulates cold stress- and osmotic stress-responsive gene expression. *Plant Cell* **13**: 2063–2083
- Xiong L, Lee H, Ishitani M, Zhu JK** (2002) Regulation of osmotic stress-responsive gene expression by the *LOS6/ABA1* locus in *Arabidopsis*. *J Biol Chem* **277**: 8588–8596
- Xu ZJ, Nakajima M, Suzuki Y, Yamaguchi I** (2002) Cloning and characterization of the abscisic acid-specific glucosyltransferase gene from adzuki bean seedlings. *Plant Physiol* **129**: 1285–1295
- Zeevaert JAD** (1980) Changes in the levels of abscisic acid and its metabolites in excised leaf blades of *Xanthium strumarium* during and after water stress. *Plant Physiol* **66**: 672–678
- Zeevaert JAD, Heath TG, Gage DA** (1989) Evidence for a universal pathway of abscisic acid biosynthesis in higher plants from ^{18}O incorporation patterns. *Plant Physiol* **91**: 1594–1601