

The Cape Verde Islands Allele of Cryptochrome 2 Enhances Cotyledon Unfolding in the Absence of Blue Light in *Arabidopsis*^{1[w]}

Javier F. Botto*, Carlos Alonso-Blanco, Ignacio Garzarón, Rodolfo A. Sánchez, and Jorge J. Casal

IFEVA, Consejo Nacional de Investigaciones Científicas y Técnicas and Facultad de Agronomía, Universidad de Buenos Aires, Avenida San Martín 4458, 1417 Buenos Aires, Argentina (J.F.B., I.G., R.A.S., J.J.C.); and Departamento de Biotecnología, Instituto Nacional de Investigación y Tecnología Agraria y Alimentaria, Carretera de A Coruña, Km 7, Madrid 28040, Spain (C.A.-B.)

We analyzed the natural genetic variation between Landsburg *erecta* (*Ler*) and Cape Verde Islands (*Cvi*) accessions by studying 105 recombinant inbred lines to search for players in the regulation of sensitivity to light signals perceived by phytochromes in etiolated seedlings of *Arabidopsis*. In seedlings grown under hourly pulses of far-red (FR) light, we identified three quantitative trait loci (QTLs; *VLF3*, *VLF4*, and *VLF5*) for hypocotyl growth inhibition and three different QTLs (*VLF6*, *VLF7*, and *VLF1*) for cotyledon unfolding. This indicates that different physiological outputs have selective regulation of sensitivity during de-etiolation. *Ler* alleles, compared with *Cvi* alleles, of *VLF3*, *VLF4*, *VLF5*, *VLF7*, and *VLF1* enhanced, whereas the *Ler* allele of *VLF6* reduced, the response to pulses of FR. We confirmed and narrowed down the position of some QTLs by using near-isogenic lines. *VLF6* mapped close to the *CRY2* (cryptochrome 2) gene. Transgenic *Ler* seedlings expressing the *Cvi* allele of *CRY2* showed enhanced cotyledon unfolding under hourly pulses of FR compared with the wild type or transgenics expressing the *CRY2-Ler* allele. This response required phytochrome A. The *cry1 cry2* double mutant lacking both cryptochromes showed reduced cotyledon unfolding under FR pulses. Because the *CRY2-Cvi* is a gain-of-function allele compared with *CRY2-Ler*, cryptochrome activity correlates positively with cotyledon unfolding under FR pulses. We conclude that the blue light photoreceptor cryptochrome 2 can modulate seedling photomorphogenesis in the absence of blue light. In addition to the nuclear loci, we identified cytoplasmic effects on seedling de-etiolation.

Seedlings grown in darkness show elongated shoot axis and rudimentary development of the foliage and of the photosynthetic machinery (Nemhauser and Chory, 2002). This etiolated body form is considered to facilitate penetration through the soil and seedling emergence. Light induces the transition from this so-called skotomorphogenic pattern to the photomorphogenic pattern of development. The seedling becomes de-etiolated as rapid axis growth ceases the leaves or cotyledons unfold, expand, and become green, whereas the cellular and biochemical components of the photosynthetic machinery are established. De-etiolation is initiated by red (R) and far-red (FR) light perceived by phytochromes (Quail, 2002), and blue light perceived primarily by cryptochromes (Cashmore et al., 1999) and secondarily by phototropins (Parks and Spalding, 1999; Briggs and Christie, 2002) and even by phytochromes. These

photoreceptors form an interconnected signaling network (Casal, 2000). There are five phytochromes (phyA–phyE) in *Arabidopsis* (Mathews and Sharrock, 1997). De-etiolation induced by R requires phyB (Reed et al., 1994) and phyA (Mazzella et al., 1997), whereas de-etiolation under FR requires phyA (Whitelam et al., 1993).

Three photobiologically discrete components of the phytochrome-mediated de-etiolation response can be distinguished in *Arabidopsis* and other species based on the effects of different R/FR treatments and different frequencies of excitation (Casal et al., 2003). The very low-fluence response (VLFR) is mediated by phyA and saturates with very low fluences of R or FR. Long-term effects on hypocotyl growth inhibition or cotyledon unfolding require these pulses to be repeated. The low-fluence response (LFR) of phyB is induced by R and not by FR, which is actually able to revert the effect of R. The high-irradiance response (HIR) is also mediated by phyA and requires sustained excitation with FR. HIR and VLFR can be dissected genetically and molecularly at the level of phyA domains (Casal et al., 2002; Yanovsky et al., 2002) and cis-acting regions of target gene promoters (Cerdán et al., 2000). The VLFR pathway would initiate de-etiolation when seedling emergence from the soil is impending. The LFR of phyB mediates de-etiolation under the high R to FR ratios of open

¹ This work was supported by the University of Buenos Aires (grant no. G406 to J.F.B.), by the Agencia Nacional de Promoción Científica y Tecnológica (grant nos. BID 1201/OC-AR and PICT 06739 to J.J.C.), and by the Spanish Ministerio de Ciencia y Tecnología (salary contract “Ramón y Cajal” to C.A.-B.).

[w] The online version of this article contains Web-only data.

* Corresponding author; e-mail botto@ifeva.edu.ar; fax 54–011–4514–8730.

Article, publication date, and citation information can be found at <http://www.plantphysiol.org/cgi/doi/10.1104/pp.103.029546>.

places, and the HIR of phyA does the same under the low R to FR ratios of dense canopies (Yanovsky et al., 1995; Smith and Whitelam, 1997). The contribution of each of these pathways differs with the species and the specific process under phytochrome control.

A source of significant genetic variation in de-etiolation responses can be found among naturally occurring populations of a given species. For example, in a set of 97 accessions of *Arabidopsis*, the promotion of hypocotyl growth by low R to FR ratios typical of dense canopies compared with high R to FR ratios of open places ranged from 50% to 600% (Botto and Smith, 2002). Compared with other accessions, the Lm-2 has reduced HIR caused by a single amino acid change affecting the biochemical properties of phyA (Maloof et al., 2001). Recombinant inbred lines (RILs) are available and allow a detailed investigation of variation between two parental strains (Alonso-Blanco and Koornneef, 2000; Maloof, 2003). This variability provides an excellent opportunity to identify loci involved in the fine regulation of VLFR, LFR, and HIR, particularly because the contribution of each one of these signaling pathways can be estimated by simultaneous analysis of different samples of each line under various light/dark conditions. The later is not possible in a mutant screening protocol. The first genetic dissection between VLFR and HIR was obtained by using RILs between Landsberg *erecta* (*Ler*) and Columbia (*Col*) to identify two quantitative trait loci (QTLs; *VLF1* and *VLF2*) controlling the VLFR of seed germination, hypocotyl growth inhibition, and cotyledon opening (Yanovsky et al., 1997). In addition, several QTLs for hypocotyl length were mapped based on the genetic variation between *Ler* and the Cape Verde Islands (*Cvi*) accessions under different light/hormone environments (Borevitz et al., 2002). Here, we uncover new loci that selectively regulate subsets of phytochrome-mediated de-etiolation responses by using *Cvi/Ler* RILs and near-isogenic lines (NILs). The identification of one of the genes involved in QTL analysis demonstrates a previously unsuspected role for a known player in photomorphogenesis. In addition to several nuclear loci, we identified cytoplasmic effects on seedling de-etiolation.

RESULTS

Hypocotyl Growth Inhibition and Cotyledon Unfolding in *Ler*, *Cvi*, and RILs

We measured hypocotyl length and the angle between cotyledons in seedlings of *Ler* and *Cvi* accessions and of 105 RILs grown in darkness or under hourly pulses of FR, hourly pulses of R, or continuous FR. In dark-grown seedlings, the hypocotyl was shorter in *Ler* compared with *Cvi* parental accessions. The difference was even larger under hourly pulses of FR and R, but it was not significant under continuous FR (Fig. 1). The cotyledons were closed in dark-

ness. Cotyledon unfolding was enhanced in *Cvi* compared with *Ler* under hourly pulses of FR (Fig. 1). However, the differences were not significant under hourly pulses of R or continuous FR. Actually, R pulses failed to enhance cotyledon unfolding compared with FR pulses in *Cvi* (Fig. 1).

The frequency distribution of RILs was very different for each physiological output. For hypocotyl length, the range of variation was large (7–8 mm in darkness and 4.7 mm under continuous FR) and similar for each light condition (Fig. 1). For cotyledon unfolding, the RILs showed a full-scale variation (7°–178°) under hourly pulses of FR, a narrow range under continuous FR (128°–180°) and an intermediate variation under hourly pulses of R (Fig. 1; Supplemental Table I).

The RILs showed transgressive variation in both directions compared with the parental lines for both traits in almost all conditions, with the exception of cotyledon angle in darkness (no unfolding) and under continuous FR (parental lines showed saturated response; Fig. 1). This suggests that parental lines contain a balance of different alleles at several loci with contrasting effects on a given physiological output. In addition, cotyledon angle and hypocotyl length values of the RIL population showed no obvious correlation under any light condition (Supplemental Fig. 1). This suggests that different loci are involved in the control of each physiological output, an expectation that was met by the results of QTL mapping described below.

Different QTLs Control Hypocotyl Growth and Cotyledon Unfolding Responses

For mapping purposes, we used two complementary approaches based either on: (a) the absolute values of hypocotyl length and cotyledon angle in darkness, pulses of FR and R, and continuous FR; or (b), the differences between darkness and pulses of FR, between pulses of FR and pulses of R, and between pulses of FR and continuous FR. These differences calculate the contribution of phyA acting in the VLFR mode, phyB mediating the LFR, and phyA acting in the HIR mode, respectively (Casal et al., 1998). Conservatively, our analysis detected five QTLs controlling hypocotyl growth and three QTLs controlling cotyledon unfolding (Fig. 2; Supplemental Figs. 2–5; Supplemental Tables II and III). The total additive effects of these QTLs accounted for a maximum of 41% of the phenotypic variation of a given trait (Fig. 2; Supplemental Tables II and III). No significant two-way QTL × QTL interaction among the various nuclear QTLs identified was detected ($P > 0.001$).





Figure 1. Phenotypes of Ler, Cvi, and Cvi/Ler RILs. Histograms show the distributions of mean hypocotyl length and mean cotyledon angle in different dark/light environments. The mean (arrows) and SDs (horizontal lines) of parental accessions are indicated. Photographs of representative seedlings of both parental accessions grown in different conditions are shown at the top.

***HYP1* and *HYP2* Control Hypocotyl Growth in Darkness**

Two loci mapping to chromosomes 1 and 2 were detected in darkness and almost under every light condition (Fig. 2). Based on Borevitz et al. (2002), who observed loci with pleiotropic effects in darkness and/or continuous light environments (i.e. white, blue, R, and FR) at similar positions using the Cvi/Ler RIL population, the loci were named *HYP1* (chromosome 1) and *HYP2* (chromosome 2).

The Ler allele of *HYP1* increased hypocotyl length compared with the Cvi allele, and the opposite was true for *HYP2* (Fig. 2). *HYP1* was not significant for

hypocotyl length under hourly pulses of R. This situation reflects the fact that the Ler allele of *HYP1* increased hypocotyl length in darkness and under FR, and it also increased the inhibition of hypocotyl growth caused by R compared with FR pulses (i.e. LFR; Fig. 2). The effect on the LFR seemed rather specific because neither the VLFR nor the HIR were affected by *HYP1*. *HYP2* mapped close to the ER (*ERECTA*) marker (Fig. 2), but because Ler and La(ER+) seedlings showed no differences in hypocotyl growth (mean ± SE, darkness, 11.9 ± 0.4 and 12.3 ± 1.3 mm; hourly FR pulses, 8.4 ± 0.7 and 8.5 ± 0.9 mm; hourly R pulses, 6.9 ± 0.3 and 6.2 ± 0.5 mm,



respectively), we conclude that *HYP2* is not the *ERECTA* locus.

VLF3, *VLF4*, and *VLF5* Control the VLFR of Hypocotyl Growth

Three QTLs mapping to chromosomes 1, 3, and 5 affected the VLFR of hypocotyl growth (Fig. 2). Following the terminology used by Yanovsky et al. (1997), who identified *VLF1* and *VLF2* in *Ler/Col* RILs, the loci at novel positions were named *VLF3*, *VLF4*, and *VLF5*, respectively. The three *VLF* QTLs together accounted for 26.3% of the total phenotypic variation observed in the VLFR. For all three loci, *Ler* alleles increased the VLFR compared with *Cvi* alleles (Fig. 2).

VLF4 was also detected for hypocotyl length under hourly pulses of FR or R, and this indicates that the magnitude of the effect on the VLFR was enough to modify the absolute hypocotyl length under these conditions. To confirm *VLF4*, we used NIL DOG6 carrying a small *Cvi* fragment between 40 and 65 cM in chromosome 3. NIL DOG6 showed reduced VLFR of hypocotyl growth compared with *Ler* ($12\% \pm 3\%$ and $27\% \pm 4\%$ inhibition, respectively), thus confirming the presence of *VLF4*.

VLF5 was localized toward the top of chromosome 5 close to *VLF2* (maximum LOD score at 25 and 7 cM, respectively). However, *VLF2* affected primarily the VLFR of cotyledon unfolding and secondarily the VLFR of hypocotyl growth, whereas *VLF5* only affected hypocotyl growth. *VLF5* also mapped close to the flowering repressor *FLC* (flowering locus C), but *flc* mutants had no obvious VLFR phenotype suggesting that *VLF5* and *FLC* are two different loci (data not shown). To confirm the position of *VLF5*, we used NIL 130 carrying an introgressed *Cvi* fragment of 30 cM at the top of the chromosome 5. As expected, NIL 130 showed reduced VLFR of hypocotyl growth compared with *Ler* ($16\% \pm 2\%$ and $27\% \pm 4\%$ inhibition, respectively).

VLF1, *VLF6*, and *VLF7* Control the VLFR of Cotyledon Unfolding

Three QTLs, *VLF6* and *VLF7* mapping to chromosome 1 and *VLF1* mapping to chromosome 2, affected the VLFR of cotyledon unfolding (Fig. 2). The VLFR of cotyledon unfolding was enhanced by the *Cvi*

allele of *VLF6* and the *Ler* alleles of *VLF7* and *VLF1*. Interestingly, the *Cvi* allele of *VLF6* increased the VLFR but reduced the LFR compared with the *Ler* allele (Fig. 2). A similar feature has been observed for other loci affecting the VLFR (e.g. *EVE1*; Luccioni et al., 2002) and has been interpreted as the result of an early element of the VLFR signaling pathway negatively regulating the LFR of phyB (Cerdán et al., 1999). The *Cvi* allele of *VLF7* also negatively affected the HIR, but it was not significant for the LFR (Fig. 2).

The locus *VLF1* was identified in chromosome 2 with an LOD score significant under hourly pulses of R (Fig. 2) and incipient but below the threshold under hourly pulses of FR (LOD = 1.55). To confirm the presence of *VLF1*, we measured cotyledon unfolding in seedlings of NIL 43 carrying a 20-cM *Cvi* fragment around the *ERECTA* locus in the *Ler* background. The cotyledon unfolding of NIL 43 seedlings was significantly reduced under hourly pulses of FR (i.e. VLFR) compared with *Ler* ($58^\circ \pm 8^\circ$ and $99^\circ \pm 6^\circ$, $P < 0.001$, respectively), and the effect was not increased under hourly pulses of R ($93^\circ \pm 9^\circ$ and $127^\circ \pm 4^\circ$, $P < 0.01$, respectively). This indicates that this locus affects mainly the VLFR. In addition, since NIL 43 showed reduced VLFR of cotyledon unfolding but normal hypocotyl length compared with *Ler* in all light/dark conditions (e.g. darkness, 11.5 ± 1.3 and 11.6 ± 0.5 mm; hourly FR pulses, 8.4 ± 0.8 and 8.4 ± 0.7 mm, respectively). *VLF1* and *HYP2* appear to be different loci (i.e. NIL 43 retained the *VLF1* but not the *HYP2* phenotype). To further refine the position of *VLF1*, we analyzed the segregation of six markers in the F_3 families from an F_2 population of 40 individuals derived from a cross between NIL 43 and *Ler*. The LOD score was below the threshold for the three markers placed toward the top end of the insertion (F3P11, MSAT2.36, and PLS7), and this excludes the *PHYB* gene as a candidate for this locus. The LOD score was significant for F13B15, which showed no segregation with the bottom end markers MSAT241 and nga1126 (data not shown). The lines with *Ler* alleles for the markers F13B15 to nga1126 showed stronger cotyledon unfolding under FR pulses than the lines with *Cvi* alleles ($85^\circ \pm 5^\circ$ and $33^\circ \pm 10^\circ$; $P = 0.003$, respectively). Thus, *VLF1* would be placed in the region between PLS7 and the bottom end of the insertion.

Figure 2. QTL mapping of hypocotyl growth and cotyledon angle for the *Cvi/Ler* RIL population grown under different light/dark environments. Mapping was based either on absolute values of hypocotyl length and cotyledon angle in darkness, hourly pulses of FR, hourly pulses of R, and continuous FR, or on the specific photobiological response (VLFR, LFR, and HIR). VLFR, Difference between darkness and pulses of FR. LFR, Difference between pulses of FR and pulses of R. HIR, Difference between pulsed and continuous FR. Rectangles, Two-log of the odds (LOD) support interval. The white arrowhead points to the position of the highest LOD score. The numbers above the arrowhead indicate the LOD score. The numbers below the white arrowhead indicate the percentage of explained variance for each QTL. Additive allele effects of each QTL are shown close to the 2-LOD support interval. The black arrowheads point upwards when the value of a given trait is increased by *Ler* compared with *Cvi* alleles and downwards when the value of a given trait is increased by *Cvi* compared with *Ler* alleles. Note that for hypocotyl growth, higher values of VLFR, LFR, or HIR mean shorter hypocotyls.

VLF6 Is the CRY2 Gene

To narrow down the position of *VLF6*, we used NIL 45- and EDI-NIL-containing Cvi regions introgressed into a *Ler* genetic background on the top of chromosome 1. Both NILs increased significantly cotyledon unfolding compared with *Ler* (Fig. 3), confirming the presence of *VLF6* at the top of chromosome 1. Because this chromosomal region contains the *CRY2* (cryptochrome 2) gene, which is known to be polymorphic between *Ler* and Cvi (El-Assal et al., 2001), we used transgenic seedlings in the *Ler* background carrying an additional copy of the *CRY2* from *Ler* (*CRY2-Ler*) or from Cvi (*CRY2-Cvi*) to investigate whether this blue light photoreceptor gene was playing a role in the absence of blue light. Cotyledon unfolding was enhanced by the Cvi but not by the *Ler* transgene (Fig. 3). The natural substitution of Val (*Ler*) for Met (Cvi) at position 367 creates a dominant gain-of-function of *CRY2* allele (El-Assal et al., 2001). Further support for a role of *CRY2* in the control of the VLFR of cotyledon unfolding was provided by transgenic lines bearing as a transgene either the *CRY2-Ler* allele, where the Val-367 was changed to Met-367, or the *CRY2-Cvi* allele bearing the Met-367 to Val-367 replacement (El-Assal et al., 2001). The




Figure 3. *VLF6* is the *CRY2* gene. Cotyledon angle in seedlings of *Ler*, NIL-EDI, NIL 45, transgenic lines carrying genomic alleles of *CRY2* from *Ler* or Cvi (*CRY2-Ler* and *CRY2-Cvi*, respectively) or with the amino acid 367 replaced to Val (*CRY2-Cvi-367V*) or with the amino acid 367 replaced to Met (*CRY2-Ler-367M*), the *cry1*, *cry2*, and *cry1 cry2* mutants grown under hourly pulses of FR. *phyA-201* and *CRY2-Cvi* in the *phyA-201* background are also included. Data are means \pm SE of three independent experiments with at least three replicate boxes each. Photographs of representative seedlings are shown at the top.




Figure 4. The Cvi allele of *CRY2* enhances the VLFR but reduces the LFR. Cotyledon angle in seedlings of *Ler*, NIL-EDI, *CRY2-Cvi*, and *CRY2-Ler-367M* grown under hourly pulses of FR, R, or R-FR mixtures providing a series of calculated Pfr/P (3%, 10%, 33%, 61%, and 87%). The standard FR source used in the rest of the experiments corresponds to a calculated Pfr/P = 10%. Inset, Slopes (Δ cotyledon angle/ Δ Pfr/P) of the VLFR (0%–10% Pfr/P) and LFR (10%–87% Pfr/P). Data are means \pm SE of three independent experiments with at least three replicate boxes each.

sole substitution of this amino acid was enough to modify the effect of the transgene on cotyledon unfolding in a manner consistent with a critical role of this residue in the whole *CRY2* gene (Fig. 3). The single *cry1* (cryptochrome 1) or *cry2* loss-of-function mutants had no significant effects, but the *cry1 cry2* double mutant showed reduced cotyledon unfolding under FR pulses ($P < 0.01$, Fig. 3). Because the Cvi is a gain-of-function allele compared with the *Ler* allele, the results of the loss-of function mutants indicate a positive correlation between cryptochrome activity and the VLFR of cotyledon unfolding. The *phyA* mutation was epistatic over the enhanced cotyledon unfolding under hourly FR (Fig. 3) but not on the flowering phenotype of the Cvi allele (data not shown; see also S.E.-D. El-Assal, C. Alonso-Blanco, A.J.M. Peeters, C. Wagemaker, J.L. Weller, and M. Koornneef, unpublished data).

The analysis of RILs indicated that the *Ler* allele of *VLF6* reduced the VLFR but increased the LFR of cotyledon unfolding (Fig. 2). This issue was investigated in further detail by characterizing the cotyledon unfolding response under hourly pulses of R, FR, or R-FR mixtures, providing a series of calculated proportions of phytochrome in the Pfr form (Fig. 4). Given the unexpected effects of *cry2* in the absence of blue light, the first important conclusion from these experiments is that the Cvi allele of *CRY2* enhances cotyledon unfolding even under hourly pulses of long-wavelength FR provided by a well-known filter (Schott RG9) that cuts off all visible light (Fig. 4, calculated Pfr/P = 3%; data not shown), indicating that the observed effects were not due to any contamination with blue light. The EDI-NIL showed en-

hanced VLFR but did not show LFR (Fig. 4, inset). This is a genuine negative effect on the LFR because these seedlings were well below the maximum attainable cotyledon unfolding (Fig. 4). The transgenic line overexpressing *CRY2-Ler* behaved like WT (calculated Pfr/P 10% = $60^\circ \pm 14^\circ$ and calculated Pfr/P 87% = $139^\circ \pm 5^\circ$).

Cytoplasmic Effects on Cotyledon Unfolding

Because the RILs were obtained from reciprocal crosses (Alonso-Blanco et al., 1998a), cytoplasmic effects could be analyzed. Noteworthy, significant LOD scores were detected for VLFR and LFR (LOD = 2 and 2.6, respectively) No significant two-way interaction was found between the cytoplasm and the nuclear QTLs identified ($P > 0.001$). The RILs derived from crosses involving a *Ler* mother showed enhanced VLFR and reduced LFR when compared with the lines derived from a *Cvi* mother (Fig. 5).

DISCUSSION

The analysis of VLFR, LFR, and HIR during de-etiolation in RILs derived from the cross between *Cvi* and *Ler* has yielded three important conclusions: First, cryptochromes can control selective aspects of photomorphogenesis in the absence of blue light; second, multiple QTLs control specific features of the de-etiolation syndrome; and third, maternal extranuclear information controls some de-etiolation responses.

The existence of specific blue-light photoreceptors had been indicated first by elegant physiological experiments where the effects of blue light could not be accounted for by the action of phytochromes (Thomas and Dickinson, 1979) and by identification of the *hy4* mutant of *Arabidopsis* affecting hypocotyl-growth responses to blue light but not to R or FR (Koornneef et al., 1980). Cloning of a T-DNA mutant allele of *hy4* finally allowed the discovery of *cry1* (Ahmad and Cashmore, 1993; Lin et al., 1995) fol-

lowed shortly afterward by *cry2*, which affects de-etiolation under very low fluences of blue light and the photoperiodic response of flowering (Lin et al., 1995, 1998). The Val residue at position 376 is highly conserved among cryptochrome genes, but it is substituted to Met in the *Cvi* allele of *CRY2* (El-Assal et al., 2001). The EDI-NIL carrying *CRY2-Cvi* is hypersensitive to continuous blue light during de-etiolation (El-Assal et al., 2001). Here, we show that an effect of *cry2* on cotyledon unfolding can also be observed under hourly pulses of FR in the absence of any blue light. This is particularly obvious for the *Cvi* compared with the *Ler* allele of *CRY2*, but it is also detectable for the *Ler* compared with the *cry2* mutant allele in the *cry1* mutant background (Fig. 3). This observation cannot be accounted for by the absorption properties of cryptochromes measured in vitro (Lin et al., 1995; Ahmad et al., 2002). However, overexpression of the carboxy termini of cryptochromes can yield hypersensitivity to FR (Yang et al., 2000).

The *phyA* mutation is epistatic to the effects of *CRY2-Cvi* reported here (Fig. 3), indicating that *cry2* is a positive regulator of *phyA*-mediated perception and/or signaling even in the absence of blue light. Genetic interaction between *phyA* and *cry2* had been observed for hypocotyl growth inhibition in seedlings de-etiolating under white light, i.e. in the presence of blue light (Mazzella et al., 2001). In concordance, two detailed works studying the dynamics of the hypocotyl inhibition under blue light demonstrated that *phyA* fully impairs *cry2* function between 30 and 120 min after the initiation of light treatment (Folta and Spalding, 2001a, 2001b).

In mammals, cryptochromes have a function as components of the clock observed even in darkness in addition to their function in the light input (van der Horst et al., 1999). In *Arabidopsis*, *cry1* and *cry2* are not key components of the clock (Devlin and Kay, 2000; Yanovsky et al., 2000a) but are involved in the light input to the clock (Yanovsky et al., 2001). In *Arabidopsis*, *cry1* and *cry2* mutants show an extended period of the circadian rhythm of expression of a light-harvesting chlorophyll-binding protein gene under R (Devlin and Kay, 2000). However, the latter effects are larger than those of the *phyA* mutation and can be observed in the *phyA* background (Devlin and Kay, 2000), indicating a fundamental difference between the effects of cryptochromes on clock period and cotyledon unfolding in the absence of blue light. El-Assal et al. (2001) have shown that *CRY2-Cvi* accelerates flowering under short photoperiods. The *phyA* mutation is not epistatic to the latter phenotype (data not shown; see also S.E.-D. El-Assal, C. Alonso-Blanco, A.J.M. Peeters, C. Wagemaker, J.L. Weller, and M. Koornneef, unpublished data), which, therefore, would be more directly mediated by *CRY2-Cvi*.

We have identified two novel QTLs for the VLFR of cotyledon unfolding (*VLF6* = *CRY2* and *VLF7*) and




Figure 5. Cytoplasmic effects on cotyledon unfolding. Average VLFR, LFR, and HIR of RILs with an *Ler* or *Cvi* cytoplasm. Data are means \pm SE of 25 (*Ler*) or 81 (*Cvi*) RILs.

three novel QTLs for the VLFR of hypocotyl growth (*VLFR3*, *VLFR4*, and *VLFR5*). A third locus controlling cotyledon unfolding is tentatively equated to the *VLFR1* locus also polymorphic between *Ler* and *Col* (Yanovsky et al., 1997). *Ler*, compared with *Cvi* alleles of *VLFR1*, *VLFR3*, *VLFR4*, *VLFR5*, and *VLFR7*, increased VLFR, whereas the opposite was true for *VLFR6*. In addition, we have confirmed and extended the role of *HYP1* and *HYP2* in the control of hypocotyl growth (Borevitz et al., 2002). The *Ler* allele of *HYP1* increased hypocotyl length in darkness, pulses of FR, or continuous FR, but not under pulses of R. This is due to enhanced LFR (phyB-mediated inhibition of hypocotyl growth caused by R compared with FR pulses; Fig. 2). Some of the QTLs affecting hypocotyl growth reported previously were not detected here, and this is likely to reflect differences in growth conditions. In particular, Borevitz et al. (2002) measured hypocotyl length after 7 d of growth in agar containing one-half-strength Murashige and Skoog salts; i.e. conditions that could enhance the impact of loci controlling seed carbohydrate reserves compared with the shorter experiments used here (3 d, plain agar). Actually, Borevitz et al. (2002) noted the coincidence between one of their loci and a previously identified seed-sized locus (Alonso-Blanco et al., 1999).

VLFR6 = *CRY2-Cvi* enhanced the VLFR and reduced the LFR (Fig. 4). This is not surprising because the *phyA* or *fly1* mutations that eliminate or reduce VLFR also enhance the LFR (Mazzella et al., 1997), the overexpression of *PHYA* (Casal and Sánchez, 1994; Casal et al., 2002), and the *eve1* (Luccioni et al., 2002) and *spa1* (Baumgardt et al., 2002) mutations that enhance VLFR reduce the LFR. These observations have been interpreted as negative regulation of phyB-mediated LFR by some component of the VLFR signaling cascade (Cerdán et al., 1999). Other loci as *VLFR1*, *VLFR2*, *VLFR3*, *VLFR5*, and *VLFR7* affect VLFR but not LFR, suggesting that they would regulate VLFR downstream the point of the VLFR pathway controlling LFR signaling. The possibility of a negative regulation of HIR signaling by VLFR signaling also has been proposed (Luccioni et al., 2002). However, although *CRY2-Cvi* and *VLFR7-Ler* reduced HIR, available information does not rule out a trivial explanation based on physical constraints for further cotyledon unfolding under HIR conditions (continuous FR), when the VLFR is already enhanced.

Most photomorphogenic mutants, including those with enhanced or reduced VLFR, have both altered hypocotyl growth and altered cotyledon unfolding responses to light (Baumgardt et al., 2002; Luccioni et al., 2002). In this report, we have demonstrated the power of QTL mapping to uncover novel loci that selectively affect one of the two physiological responses. This finding suggests a scenario where plants would bear a relatively large number of QTLs

with developmental specific action, providing flexibility to the de-etiolation responses.

At least between *Ler* and *Cvi*, there is a clear imbalance in the variability for the different response modes. Six QTLs affect primarily the VLFR, but only one (*HYP1*) shows an effect on the LFR that cannot be accounted for on the basis of a primary action on VLFR (as discussed in the previous paragraph), and none affects selectively the HIR. This is also true for the variability between *Ler* and *Col* (Yanovsky et al., 1997). This might merely reflect the genetic information of *Ler*, present in both crosses, or a more general tendency for genetic variability in sensitivity to weak signals compared with maximum attainable response. It will be interesting to explore this issue as new RILs become available.

Because the RILs used here are derived from reciprocal crosses between the parental lines (Alonso-Blanco et al., 1998a), it is possible to evaluate the occurrence of cytoplasmic effects. The lines derived from a cross involving a *Ler*, compared with *Cvi* mother, showed enhanced VLFR and reduced LFR and HIR (Fig. 5). At least in principle, the effects on LFR and HIR could derive from those on VLFR as described in previous paragraphs for the effects of nuclear loci. The reports of cytoplasmic effects are scant (Corey et al., 1976). A similar analysis conducted for seed traits that are affected by the genotype of the mother plant concluded against the occurrence of a cytoplasmic control for those traits (Alonso-Blanco et al., 1999). Retrograde signals emanate from the chloroplast to control nuclear gene expression (Surpin et al., 2002). Mutations at a nuclear gene coding a soluble ATP-binding cassette protein that localizes at the chloroplast periphery impair phyA-mediated de-etiolation under continuous FR (Møller et al., 2001). Therefore, it is tempting to speculate that chloroplast loci polymorphic between *Ler* and *Cvi* could modulate retrograde signaling and, hence, modulate phytochrome-mediated responses.

MATERIALS AND METHODS

Plant Material

The parental lines *Ler* and *Cvi* and a set of 105 RILs derived from reciprocal crosses between the two genotypes (Alonso-Blanco et al., 1998a) were obtained from Arabidopsis Biological Resource Center (Ohio State University, Columbus). The *La(ER+)*, NIL 42, NIL 45, NIL-EDI (Alonso-Blanco et al., 1998b; Swarup et al., 1999), and NIL DOG6 (Alonso-Blanco et al., 2003) were described previously. NIL 43 carrying a single introgression of about 20 cM from physical positions 8.3 to 11.7 Mb of chromosome 2 from *Cvi* into a *Ler* background was kindly provide by Dr. Andrew Millar (University of Warwick, UK). NIL 130 carries a single introgression from the top of chromosome 5 of *Cvi* into a *Ler* genetic background, with a recombination breakpoint between markers AD129 and HH.480. The transgenic lines carrying *CRY2-Ler* or *CRY2-Cvi* transgenes or *CRY2* transgenes mutated in a single amino acid at position 367 in the *Ler* background were described before (El-Assal et al., 2001) and kindly provide by Dr. Maarten Koornneef (University of Wageningen, The Netherlands). The *phyA-201* allele in *Ler* (Reed et al., 1994) was used to produce *CRY2-Cvi* transgenic plants in the *phyA* background. F₃ families from an F₂ population of 40

individuals derived from a cross between NIL 43 and *Ler* were genotyped for six markers covering regularly the Cvi region introgressed in the NIL 43. The microsatellite markers MSAT2.36, MSAT2.41 (Loudet et al., 2002) and nga1126 (Bell and Ecker, 1994) and the INDELs F3P11, PLS7, and F13B15 (AtDB) were used.

Growth Conditions, Light Treatments, and Physiological Measurements

Twenty seeds were sown in clear plastic boxes (40 × 33-mm² × 15-mm height) containing 3 mL of 0.8% (w/v) agar. The boxes were incubated at 4°C in darkness for 3 d and exposed to an R pulse followed by 24 h of darkness to promote seed germination. The seedlings were either kept in the dark or exposed to various light treatment for 3 d. Hypocotyl length was measured to the nearest 0.5 mm with a ruler, and, to eliminate defective seedlings, only the largest 10 seedlings of each box were averaged. The angle between the cotyledons was measured with a protractor.

Hourly pulses of R (3 min, 20 μmol m⁻² s⁻¹), hourly pulses of FR (3 min, 50 μmol m⁻² s⁻¹), continuous FR (2.5 μmol m⁻² s⁻¹), and the mixtures of R-FR (3 min, at least 11 μmol m⁻² s⁻¹) used to obtain intermediate calculated proportions of phytochrome in the Pfr form were provided by the sources described earlier (Yanovsky et al., 2000b). Long-wavelength FR (3 min, 40 μmol m⁻² s⁻¹) was provided by incandescent lamps in combination with a water filter and an RG9 filter (Schott, Mainz, Germany).

Analysis of QTLs and Cytoplasmic Effects

Each trait was analyzed separately by using the mean values from three independent experiments. All the lines were included in parallel in each experiment. Each experiment included 10 seedlings per RIL. Hypocotyl length data were normally distributed. Cotyledon angle data were transformed (arcsin √) to improve the normality of distributions, but none of the conclusions was changed when using the original data. Therefore, results are presented in figures with the original scale. A set of 99 markers covering nearly all of the *Arabidopsis* genetic map at average intervals of 5 cM was selected from the Cvi/*Ler* RIL map (Alonso-Blanco et al., 1998a). The computer software MapQTL version 4.0 (van Ooijen, 2000) was used to identify and locate QTLs on the linkage map by using interval mapping and multiple-QTL model mapping methods as described previously (Alonso-Blanco et al., 2003). LOD threshold values applied to declare the presence of a QTL were estimated with the permutation tests implemented in MapQTL version 4. The quantitative trait data of the RILs were permuted 1,000 times over the genotypes and empirical LOD thresholds corresponding to a genome wide significance $\alpha = 0.05$ were estimated between 2.5 and 2.7 for the various data sets. Two-LOD support intervals were established as ≈95% QTL confidence intervals. The estimated additive genetic effect and the percentage of variance explained by each QTL and the total variance explained by all the QTLs affecting a trait were obtained with MapQTL in the final multiple-QTL model in which one cofactor marker was fixed per QTL. Additive genetic effects presented correspond to the differences between the estimated means of the two homozygous RIL genotypic groups, at each particular QTL, divided into two. Two-way interactions among the QTLs identified were tested by ANOVA using the corresponding two markers as factors. In addition, two-way interactions were searched for among all pair-wise combinations of the 99 nuclear markers and the cytoplasmic genotype, using the computer program EPISTAT (Chase et al., 1997) with log-likelihood ratio threshold corresponding to a significance $P < 0.001$. Because 81 of the RILs carry *Ler* cytoplasm and 25 carry Cvi cytoplasm, cytoplasmic genetics effects were analyzed in the RIL population using the cytoplasmic genotype as a factor in one-way ANOVA and in multiple factor linear models in combination with the nuclear QTL markers affecting each trait.

ACKNOWLEDGMENTS

We thank Ana María Rodríguez for her technical assistance. We also thank Maarten Koornneef, Salah E.-D. El-Assal, and Leonie Bentsink for kindly providing the seeds of *CRY2* transgenic lines and NIL *DOG6*, Andrew Millar for the seeds of NIL 43, and the *Arabidopsis* Biological Resource Center (Ohio State University, Columbus) for the RILs.

Received July 3, 2003; returned for revision August 3, 2003; accepted August 28, 2003.

LITERATURE CITED

- Ahmad M, Cashmore AR (1993) HY4 gene of *Arabidopsis thaliana* encodes a protein with characteristics of a blue-light photoreceptor. *Nature* **366**: 162–166
- Ahmad M, Grancher N, Heil M, Black RC, Giovani B, Galland P, Lardemer D (2002) Action spectrum for cryptochrome-dependent hypocotyl growth inhibition in *Arabidopsis*. *Plant Physiol* **129**: 1–12
- Alonso-Blanco C, Bentsink L, Hanhart CJ, Blankestijn-deVries H, Koornneef M (2003) Analysis of natural allelic variation at seed dormancy loci of *Arabidopsis thaliana*. *Genetics* **164**: 711–729
- Alonso-Blanco C, Blankestijn-deVries H, Hanhart CJ, Koornneef M (1999) Natural allelic variation at seed size loci in relation to other life history traits of *Arabidopsis thaliana*. *Proc Natl Acad Sci USA* **96**: 4710–4717
- Alonso-Blanco C, El-Assal SE-D, Coupland G, Koornneef M (1998b) Analysis of natural allelic variation at flowering time loci in the Landsberg *erecta* and Cape Verde islands ecotypes of *Arabidopsis thaliana*. *Genetics* **149**: 749–764
- Alonso-Blanco C, Koornneef M (2000) Naturally occurring variation in *Arabidopsis*: an underexploited resource for plant genetics. *Trends Plant Sci* **5**: 22–29
- Alonso-Blanco C, Peeters AJM, Koornneef M, Lister C, Dean C, van der Bosch N, Pot J, Kuiper MTR (1998a) Development of an AFLP based linkage map of *Ler*, Col and Cvi *Arabidopsis thaliana* ecotypes and construction of a *Ler*/Cvi recombinant inbred line population. *Plant J* **14**: 259–271
- Baumgardt RL, Oliverio KA, Casal JJ, Hoecker U (2002) SPA1, a component of phytochrome A signal transduction, regulates the light signaling current. *Planta* **21**: 745–753
- Bell CJ, Ecker JR (1994) Assignment of 30 microsatellite loci to the linkage map of *Arabidopsis*. *Genomics* **19**: 137–144
- Botto JF, Smith H (2002) Differential genetic variation in adaptive strategies to a common environmental signal in *Arabidopsis* accessions: phytochrome-mediated shade avoidance. *Plant Cell Environ* **25**: 53–64
- Borevitz JO, Maloof JN, Lutes J, Dabi T, Redfern JL, Trainer GT, Werner JD, Asami T, Berry CC, Weigel D, Chory J et al. (2002) Quantitative trait loci controlling light and hormone response in two accessions of *Arabidopsis thaliana*. *Genetics* **160**: 683–696
- Briggs WR, Christie JM (2002) Phototropins 1 and 2: versatile plant blue-light receptors. *Trends Plant Sci* **7**: 204–210
- Casal JJ (2000) Phytochromes, cryptochromes, phototropin: photoreceptor interactions in plants. *Photochem Photobiol* **71**: 1–11
- Casal JJ, Davis SJ, Kirchenbauer D, Viczian A, Yanovsky MJ et al (2002) The serine-rich N-terminal domain of oat phytochrome A helps regulate light responses and subnuclear localization of the photoreceptor. *Plant Physiol* **129**: 1127–1137
- Casal JJ, Luccioni LG, Oliverio KA, Boccalandro HE (2003) Light, phytochrome signalling and photomorphogenesis in *Arabidopsis*. *Photochem Photobiol Sci* **2**: 625–636
- Casal JJ, Sánchez RA (1994) Overexpression of oat phytochrome A gene differentially affects stem growth responses to red/far-red ratio signals characteristic of sparse or dense canopies. *Plant Cell Environ* **17**: 407–419
- Casal JJ, Sánchez RA, Botto JF (1998) Modes of action of phytochromes. *J Exp Bot* **49**: 127–138
- Cashmore AR, Jarillo JA, Wu Y-J, Liu D (1999) Cryptochromes: blue light receptors for plants and animals. *Science* **284**: 760–765
- Cerdán PD, Staneloni RJ, Ortega J, Bunge MM, Rodríguez-Batiller J, Sánchez RA, Casal JJ (2000) Sustained but not transient phytochrome A signaling targets a region of a *Lhcb1*2* promoter that is not necessary for phytochrome B action. *Plant Cell* **12**: 1203–1211
- Cerdán PD, Yanovsky MJ, Reymundo FC, Nagatani A, Staneloni RJ, Whitelam GC, Casal JJ (1999) Regulation of phytochrome B signaling by phytochrome A and FHY1 in *Arabidopsis thaliana*. *Plant J* **18**: 499–507
- Chase K, Adler FR, Lerk KG (1997) EPISTAT: a computer program for identifying and testing interactions between pairs of quantitative trait loci. *Theor Appl Genet* **242**: 81–89
- Corey LA, Matzinger DF, Cockerham CC (1976) Maternal and reciprocal effects on seedling characters in *Arabidopsis thaliana* (L.) HEYNH. *Genetics* **82**: 677–683

- Devlin PF, Kay SA** (2000) Cryptochromes are required for phytochrome signaling to the circadian clock but not for rhythmicity. *Plant Cell* **12**: 2499–2510
- El-Assal SE-D, Alonso-Blanco C, Peeters AJM, Raz V, Koornneef M** (2001) A QTL for flowering time in *Arabidopsis* reveals a novel allele of *CRY2*. *Nat Genet* **29**: 435–439
- Folta KM, Spalding EP** (2001a) Opposing roles of phytochrome A and phytochrome B in early cryptochrome-mediated growth inhibition. *Plant J* **28**: 333–340
- Folta KM, Spalding EP** (2001b) Unexpected roles for cryptochrome 2 and phototropin revealed by high-resolution analysis of blue light-mediated hypocotyl growth inhibition. *Plant J* **26**: 471–478
- Koornneef M, Rolff E, Spruit CJP** (1980) Genetic control of light-inhibited hypocotyl elongation in *Arabidopsis thaliana* (L.) Heynh. *Z Pflanzenphysiol* **100**: 147–160
- Lin C, Robertson DE, Ahmad M, Raibekas AA, Jorns MS, Dutton PL, Cashmore AR** (1995) Association of flavin adenine dinucleotide with the *Arabidopsis* blue light receptor *CRY1*. *Science* **269**: 968–970
- Lin C, Yang H, Guo H, Mockler T, Chen J, Cashmore AR** (1998) Enhancement of blue-light sensitivity of *Arabidopsis* seedlings by a blue light receptor cryptochrome 2. *Proc Natl Acad Sci USA* **93**: 15491–15496
- Loudet O, Chaillou S, Goubely C, Depeiges A, Picard G** (2002) Bay-0 x Shahdara recombinant inbred line population: a powerful tool for the genetic dissection of complex traits in *Arabidopsis*. *Theor Appl Genet* **104**: 1173–1184
- Luccioni LG, Oliverio KA, Yanovsky MJ, Boccalandro H, Casal JJ** (2002) Brassinosteroid mutants uncover fine tuning of phytochrome signaling. *Plant Physiol* **128**: 173–181
- Maloo JN** (2003) QTL for plant growth and morphology. *Curr Opin Plant Biol* **6**: 85–90
- Maloo JN, Borevitz JO, Babi T, Lutes J, Nehring RB, Redfern JL, Trainer GT, Wilson JM, Asami T, Berry CC et al.** (2001) Natural variation in light sensitivity of *Arabidopsis*. *Nat Genet* **29**: 441–446
- Mathews S, Sharrock RA** (1997) Phytochrome gene diversity. *Plant Cell Environ* **20**: 666–671
- Mazzella MA, Alconada Magliano TM, Casal JJ** (1997) Dual effect of phytochrome A on hypocotyl growth under continuous red light. *Plant Cell Environ* **20**: 261–267
- Mazzella MA, Cerdán PD, Staneloni RJ, Casal JJ** (2001) Hierarchical coupling of phytochromes and cryptochromes reconciles stability and light modulation of *Arabidopsis* development. *Development* **128**: 2291–2299
- Møller SG, Kunkel T, Chua NH** (2001) A plastidic ABC protein involved in intercompartmental communication of light signaling. *Genes Dev* **15**: 90–103
- Nemhauser J, Chory J** (2002) In CR Somerville, EM Meyerowitz, eds, *The Arabidopsis Book*. American Society of Plant Biologists, Rockville, MD, doi/10.1199/tab.0054, <http://www.aspb.org/publications/arabidopsis>
- Parks BM, Spalding EP** (1999) Sequential and coordinated action of phytochrome A and B during *Arabidopsis* stem growth revealed by kinetics analysis. *Proc Natl Acad Sci USA* **96**: 14142–14146
- Quail PH** (2002) Phytochrome photosensory signaling networks. *Nat Rev* **3**: 85–93
- Reed JW, Nagatani A, Elich TD, Fagan M, Chory J** (1994) Phytochrome A and phytochrome B have overlapping but distinct functions in *Arabidopsis* development. *Plant Physiol* **104**: 1139–1149
- Smith H, Whitelam GC** (1997) The shade avoidance syndrome: multiple responses mediated by multiple phytochromes. *Plant Cell Environ* **20**: 840–844
- Surpin M, Larkin RM, Chory J** (2002) Signal transduction between the chloroplast and the nucleus. *Plant Cell* **14**: S327–S338
- Swarup K, Alonso-Blanco C, Lynn JR, Michaels SD, Amasino RM, Koornneef M, Millar A** (1999) Natural allelic variation identifies new genes in the *Arabidopsis* circadian system. *Plant J* **20**: 67–77
- Thomas B, Dickinson HG** (1979) Evidence for two photoreceptors controlling growth in de-etiolated seedlings. *Planta* **146**: 545–550
- van der Horst GT, Muijtjens M, Kobayashi K, Takano R, Kanno S, Takao M, Dewit J, Verkerk A, Eker APM, Van Leenen D et al.** (1999) Mammalian *Cry1* and *Cry2* are essential for maintenance of circadian rhythms. *Nature* **398**: 627–630
- van Ooijen** (2000) MapQTL (r) Version 4.0: User Friendly Power in QTL Mapping: Addendum to the Manual of Version 3.0. Plant Research International, Wageningen, The Netherlands.
- Whitelam GC, Johnson E, Peng J, Carol P, Anderson ML, Cowl JS, Harberd NP** (1993) Phytochrome A null mutants of *Arabidopsis* display a wild-type phenotype in white light. *Plant Cell* **5**: 757–768
- Yang H-Q, Wu Y-J, Tang R-H, Liu D, Liu Y, Cashmore AR** (2000) The C termini of *Arabidopsis* cryptochromes mediate a constitutive light response. *Cell* **103**: 815–827
- Yanovsky MJ, Casal JJ, Luppi JP** (1997) The VLF loci, polymorphic between ecotypes *Landsberg erecta* and *Columbia*, dissect two branches of phytochrome A signal transduction that correspond to very-low-fluence and high-irradiance responses. *Plant J* **12**: 101–109
- Yanovsky MJ, Casal JJ, Whitelam GC** (1995) Phytochrome A, phytochrome B and *HY4* are involved in hypocotyl growth responses to natural radiation in *Arabidopsis*: weak de-etiolation of the *phyA* mutant under dense canopies. *Plant Cell Environ* **18**: 788–794
- Yanovsky MJ, Luppi JP, Kirchbauer D, Ogorodnikova OB, Sineshchekov VA, Adam E, Kircher S, Staneloni RJ, Schäfer E, Nagy F, Casal JJ** (2002) Missense mutation in the PAS2 domain of phytochrome A impairs subnuclear localization and a subset of responses. *Plant Cell* **14**: 1591–1603
- Yanovsky MJ, Mazzella MA, Casal JJ** (2000a) A quadruple photoreceptor mutant still keeps track of time. *Curr Biol* **10**: 1013–1015
- Yanovsky MJ, Mazella MA, Whitelam GC, Casal JJ** (2001) Resetting of the circadian clock by phytochromes and cryptochromes in *Arabidopsis*. *J Biol Rhythms* **16**: 523–530
- Yanovsky MJ, Whitelam GC, Casal JJ** (2000b) *fhy3-1* retains inductive responses of phytochrome A. *Plant Physiol* **123**: 235–242