

A Nodule-Specific Dicarboxylate Transporter from Alder Is a Member of the Peptide Transporter Family¹

Jeeyon Jeong², SuJeong Suh², Changhui Guan, Yi-Fang Tsay, Nava Moran, Chang Jae Oh, Chung Sun An, Kirill N. Demchenko, Katharina Pawlowski³, and Youngsook Lee^{3*}

Department of Life Science, Pohang University of Science and Technology, Hyoja-dong, san 31, Pohang 790–784, Korea (J.J., S.S., Y.L.); Department of Molecular Biology, Agricultural University, Dreijenlaan 3, 6703 HA Wageningen, The Netherlands (C.G., K.P.); Max-Planck-Institut für Züchtungsforschung, Carl-von-Linné-Weg 10, 50829 Köln, Germany (C.G.); Institute of Molecular Biology, Academia Sinica, Taipei, Taiwan (Y.-F.T.); Institute of Plant Sciences and Genetics in Agriculture, Faculty of Agricultural, Food, and Environmental Quality Sciences, The Hebrew University of Jerusalem, Rehovot 76100, Israel (N.M.); School of Biological Sciences, Seoul National University, Kwanak-gu, Seoul 151–742, Korea (C.J.O., C.S.A.); and Albrecht von Haller Institute for Plant Sciences, Plant Biochemistry, Göttingen University, Justus-von-Liebig-Weg 11, 37077 Göttingen, Germany (K.N.D., K.P.)

Alder (*Alnus glutinosa*) and more than 200 angiosperms that encompass 24 genera are collectively called actinorhizal plants. These plants form a symbiotic relationship with the nitrogen-fixing actinomycete *Frankia* strain HFPAr13. The plants provide the bacteria with carbon sources in exchange for fixed nitrogen, but this metabolite exchange in actinorhizal nodules has not been well defined. We isolated an alder cDNA from a nodule cDNA library by differential screening with nodule versus root cDNA and found that it encoded a transporter of the PTR (peptide transporter) family, *AgDCAT1*. *AgDCAT1* mRNA was detected only in the nodules and not in other plant organs. Immunolocalization analysis showed that *AgDCAT1* protein is localized at the symbiotic interface. The *AgDCAT1* substrate was determined by its heterologous expression in two systems. *Xenopus laevis* oocytes injected with *AgDCAT1* cRNA showed an outward current when perfused with malate or succinate, and *AgDCAT1* was able to complement a dicarboxylate uptake-deficient *Escherichia coli* mutant. Using the *E. coli* system, *AgDCAT1* was shown to be a dicarboxylate transporter with a K_m of 70 μM for malate. It also transported succinate, fumarate, and oxaloacetate. To our knowledge, *AgDCAT1* is the first dicarboxylate transporter to be isolated from the nodules of symbiotic plants, and we suggest that it may supply the intracellular bacteria with dicarboxylates as carbon sources.

Some plants and microorganisms engage in reciprocal symbiosis for the purpose of exchanging nutrients. For example, in nitrogen-fixing nodules, the intracellular bacteria supply the host plant with combined nitrogen and are in turn provided with carbon sources (Mylona et al., 1995). There are two types of nodule symbioses between nitrogen-fixing soil bacteria and higher plants, namely, the symbiosis between legumes and rhizobia, and the actinorhizal symbiosis between actinomycetes of the genus *Frankia* and a diverse group of angiosperms collectively called actinorhizal plants. In both cases, nutri-

ent exchange between the host and its microsymbiont is controlled by the plant plasma membrane-derived interface enclosing the microsymbiont. In most legume symbioses, where the bacteria are taken up into the plant cells in a complete endocytotic process (Verma, 1992), this interface is the peribacteroid membrane (PBM). In primitive legume symbioses (de Faria et al., 1987) and the actinorhizal symbioses (Mylona et al., 1995), the interface is reported to be the invaginated and incompletely enclosed plasma membrane of the infected cell.

The nutrient exchange between the symbiotic partners requires transporters of the carbon sources and trace elements that flow from the plant to the microsymbiont along with the transporters of the products of bacterial nitrogen fixation that flow from the microsymbiont to the plant (Pawlowski and Bisseling, 1996). Soybean (*Glycine max*) nodules, which represent legume symbioses, show evidence of the physiological or biochemical activities of transporters of ammonium, which is the product of bacterial nitrogen fixation (Tyerman et al., 1995), and of transporters of dicarboxylates, the form of the carbon source that is supplied to the nitrogen-fixing microorganisms (Streeter, 1995). However, the genetic identities

¹ This work was supported in part by the National Science Council of Taiwan (grant no. NSC86–2811–B–001–076R to Y.L. when at Academia Sinica), by the National Research Laboratory program of Ministry of Science and Technology of Korea (to Y.L.), and by the Dutch Organization for Scientific Research (to C.G. and K.P. when at Wageningen University).

² These authors contributed equally to the work.

³ These authors contributed equally to the manuscript.

* Corresponding author; e-mail ylee@postech.ac.kr; fax 82–54–2792199.

Article, publication date, and citation information can be found at <http://www.plantphysiol.org/cgi/doi/10.1104/pp.103.032102>.

of these transporters remain unclear. It was initially suggested that nodulin-26 in the PBM of soybean nodules mediates the transport of dicarboxylates (Ou Yang et al., 1991), but it was later revealed that this molecule is an aquaporin that probably participates in osmoregulation (Rivers et al., 1997; Dean et al., 1999).

Dicarboxylates also appear to be the carbon source that actinorhizal plants provide to their *Frankia* sp. microsymbionts as suggested by studies on the enzymatic activities of *Frankia* sp. isolated from alder (*Alnus glutinosa*) nodules (Huss-Danell, 1997). Here, we identify a dicarboxylate transporter that is localized in the nitrogen-fixing nodules of alder. The nodule-specific expression and substrate specificity of this transporter support the notion that it participates in the flow of carbon sources from the plant cytoplasm to the symbiotic bacteria.

RESULTS

Isolation and Characterization of cDNAs Encoding a Nodule-Specific Transporter from Alder

An alder nodule cDNA library was screened differentially using nodule versus root cDNA. Seven clones representing nodule-specific genes were identified and characterized. By sequencing the ends of the inserts, three were found to contain full-size cDNAs of the same gene, which was later termed *AgDCAT1* (*Alnus glutinosa* dicarboxylate transporter 1) as the data we will describe in this paper indicate that it transports dicarboxylates. The insert of the three clones was sequenced completely (EMBL accession no. AGL488290). When we searched the GenBank database, we found that the protein encoded by the cloned cDNAs is a novel member of the PTR (peptide transporter) family (Fig. 1) because it contains 12 putative transmembrane-spanning domains with a large hydrophilic loop between transmembrane domains VI and VII and the signature motif for the PTR family, F-Y-x-x-I-N-x-G-S-L, within transmembrane domain V. Furthermore, the central loop of *AgDCAT1* contains the protein kinase C recognition motif (T-x-R/K) that is also conserved in the PTR family transporters (Steiner et al., 1995). When compared with the members of the PTR family that have been characterized already, it had the highest homology to CHL1, the nitrate transporter of *Arabidopsis* (Tsay et al., 1993; Frommer et al., 1994). DNA gel-blot analysis indicated that the corresponding gene is encoded by a small gene family (Fig. 2A). RNA gel-blot analysis showed expression of this gene only in nodules and not in roots, shoot tips, flowers, or developing fruits (Fig. 2B; data not shown for flowers and developing fruits).

AgDCAT1-Specific Antisera Recognize Two Bands in a Western Blot

Antisera were raised against two different peptides of *AgDCAT1* (TGM13 and CED15; see "Materials and

Methods") and used for hybridization with blots containing plasma membrane proteins from alder nodules. In a dilution of 1:500, both the anti-TGM13 and anti-CED15 antisera hybridized with the same two bands that correspond to polypeptides with an apparent molecular masses of 54 and 61 kD (Fig. 2C). Peptide competition experiments were performed to test antibody specificity. The pre-incubation of anti-TGM13 with the immunizing peptide used to raise it prevented the appearance of both hybridizing bands in western blots (data not shown). The calculated molecular mass of *AgDCAT1* is 64 kD. Hence, the 54-kD band might result from proteolysis of *AgDCAT1*. However, it should be noted that the estimation of the molecular mass of hydrophobic polypeptides on SDS-PAGE gels can be imprecise because they tend to run faster than typical standard proteins of similar molecular mass (Takagi, 1991). At any rate, both bands could correspond to *AgDCAT1* and probably result from posttranslational modification of the original *AgDCAT1* protein or its processed form. Glycosylation, for example, can change the apparent molecular mass of a transporter by up to 30 kD (Arima et al., 2002). Phosphorylation can also change the apparent molecular mass of a protein by several kilodaltons (Husain et al., 1996).

Immunolocalization of AgDCAT1 in Infected Alder Nodule Cells

When *Frankia* sp.-infected alder root nodule sections were incubated with anti-TGM13 antiserum, the immunoreactant was found to be localized at the infected cells (Fig. 3, A, B, D, and E) and, more specifically, at the interface between the plant cell and the bacteria (Fig. 3, G and H). Together with our western-blot experiment that showed the presence of the protein in the plasma membrane-enriched fraction of the nodules (Fig. 2C), these images strongly support the notion that *AgDCAT1* is localized at the plant plasma membrane-derived interface that encloses the microsymbiont. In contrast, hardly any fluorescence (except for the autofluorescence from the xylem cell wall) was visible from a negative control section from which the primary antiserum was omitted at the first incubation step (Fig. 3, C and F).

Functional Expression of AgDCAT1 in *Xenopus laevis* Oocytes

Members of the PTR family transport diverse substrates, namely, nitrate, amino acids or oligopeptides and their substrate specificities have been well characterized by their functional expression in *X. laevis* oocytes (Williams and Miller, 2001). Therefore, we used the *X. laevis* oocyte system to identify the substrate of *AgDCAT1*. Oocytes injected with *AgDCAT1* cRNA did not respond to allantoin, citrulline, ammonia, dipeptides, or Glu ($n = 13-18$). However, when

A





Figure 1. AgDCAT1 sequence analysis. A, Comparison of the amino acid sequences of AgDCAT1, the Arabidopsis nitrate transporter CHL1 (Tsay et al., 1993; AtCHL1A), and the Arabidopsis peptide transporter PTR2-B (Frommer et al., 1994; AtPTR2B). Gaps to optimize the alignment were introduced by using the Program ClustalW (EMBL), and the editor GeneDoc was used to present the alignment (Nicholas et al., 1997). Identical amino acids at conserved positions are labeled by inverse print, whereas chemically similar amino acids are shaded in gray. A signature motif of the PTR family within transmembrane domain V, F-Y-x-x-I-N-x-G-S-L, is shown in bold letters (amino acids 194–203). The protein kinase C recognition motif (T-x-R) in the central loop is shown in italics (amino acids 239–241). B, Hydropathy profile of AgDCAT1. Hydrophilicity was determined by the method of Kyte and Doolittle using a window of 19 amino acid residues. The numbers I to XII refer to the putative membrane-spanning segments.

B



the injected oocytes were perfused with 10 mM malate (pH 5.5), an outward current was generated (Fig. 4A, left; $n = 23$), as would be expected because of the influx of a negatively charged substrate. Succinate mimicked malate (Fig. 4A, middle; $n = 10$), but Cl^- or SO_4^- failed to generate similar outward currents ($n = 10$; data not shown). Noninjected oocytes or water-injected oocytes did not respond to either

malate or succinate (Fig. 4B, left and middle; $n = 18$). The responses of the injected oocytes to NO_3^- were small and inconsistent ($n = 20$; data not shown). Upon perfusion with the buffer alone and changing the pH from 7.4 to 5.5 (Fig. 4A, right), the injected oocyte developed a small inward current. Although this could result from an AgDCAT1-associated change in leak conductance as has been found for a




Figure 2. DNA, RNA, and protein gel-blot hybridization analysis. A, DNA gel blot containing the total DNA of alder digested with *Eco*RI (E), *Bam*HI (B), and *Hind*III (H), respectively, was hybridized with the 32 P-labeled insert of pAgDCAT1. The *AgDCAT1* cDNA contains one *Bam*HI site and two *Hind*III sites (223 bp apart). The numbers indicate the sizes of the marker fragments in kb. B, RNA gel blot containing the total RNA of shoot tips, nodules, and roots of alder was hybridized with the 32 P-labeled insert of pAgDCAT1. Hybridization with a heterologous ubiquitin (*ubi*) probe (Kouchi and Hata, 1993) was used to quantify the amount of mRNA on the filter. C, Western blots of plasma membrane proteins from alder nodules. Plasma membrane proteins extracted from alder nodules were separated by SDS-PAGE, transferred to a nitrocellulose membrane, and probed using the antisera specified. The numbers indicate protein molecular mass standards. The molecular masses of the two bands are approximately 54 and 61 kD. No signal was detected using plasma membrane proteins from alder leaves (data not shown).

variety of other transporters (e.g. Sonders et al., 1997), further study is required to clarify this phenomenon. No distinct response was observed from the noninjected oocyte (Fig. 4B, right). These results suggest that *AgDCAT1* is a dicarboxylate-specific transporter, with a much higher preference for malate and succinate than for the dicarboxylic amino acid Glu or for dipeptides.

To test if the oocytes indeed took up malate via *AgDCAT1*, we measured [14 C]malate accumulation in *AgDCAT1*-expressing oocytes. Thus, water- and *AgDCAT1*-injected oocytes were incubated in a bath containing 1 mM malate at pH 5.5 for 30 min. Although the malate accumulation of three separate batches of *AgDCAT1*-expressing oocytes was $205\% \pm 96\%$ (SE) of that of water-injected oocytes, the difference between the two groups was not statistically significant ($P > 0.05$, $n = 18$). Voltage clamping of the oocytes at -60 , -30 , or 0 mV during the uptake period did not sufficiently reduce the variability among the oocytes; therefore, we were not able to detect the significant accumulation of malate in *AgDCAT1*-expressing oocytes under these experimental conditions either (data not shown). This may

be because of variability between the different oocyte batches and/or an additional modulatory interaction of the transporter with its own substrate malate, as has been shown before for other transporters (Lin et al., 1996; Sonders and Amara, 1996; Sonders et al., 1997).

AgDCAT1 Mediates Malate Transport in Budding Yeast (*Saccharomyces cerevisiae*) and an *Escherichia coli* *dct* Mutant

For further functional analysis of *AgDCAT1*, the gene was introduced into a wild-type strain of budding yeast that lacks an active dicarboxylate transporter (Camarasa et al., 2001) and into CBT315, a dicarboxylate transport mutant (*dctA*) strain of *E. coli* (Lo et al., 1972). Both systems showed enhanced malate uptake when *AgDCAT1* was introduced. Yeast cells transformed with pFL61-*AgDCAT1* took up 2-fold more malate than those transformed with empty vector (pFL61) when [14 C]malate uptake was measured for 2 min with 1 mM malate (Fig. 5). Because *E. coli* showed more reproducible and higher rates of malate uptake in comparable conditions (see below), *E. coli* was chosen for further analysis. *E. coli* has been used successfully for the functional characterization of other plant transporters (Kim et al., 1998; Uozumi, 2001).

To test the ability of *AgDCAT1* to complement CBT315, *AgDCAT1* was cloned into the *E. coli* expression vector pKK223-3 under the control of the *tac* promoter (Brosius and Holy, 1984), yielding pAgDCAT1. The wild type that had been transformed with the empty vector, K-12(pKK223-3), was used as a positive control, and the empty vector transformant of CBT315, CBT315(pKK223-3), was used as a negative control. Because the *tac* promoter in pKK223-3 confers a low level of expression even in the absence of its inducer, isopropylthio- β -D-thiogalactoside, the experiments were performed in the absence of isopropylthio- β -D-thiogalactoside. The growth of the three transformants on M9 medium with 10 mM malate as the sole carbon source was then compared. K12(pKK223-3) and CBT315(pAgDCAT1) were able to grow under these conditions, whereas CBT315(pKK223-3) was not (Fig. 6A). Thus, *AgDCAT1* complements the dicarboxylate uptake deficiency of CBT315, even when it is expressed at a very low level.

The function of *AgDCAT1* in *E. coli* was analyzed further by comparing the malate uptake of CBT315(pKK223-3) and CBT315(pAgDCAT1). To obtain the initial rate of malate transport, time-dependent uptake was measured over 15 min in the presence of 1 mM malate (Fig. 6B). In CBT315(pAgDCAT1), malate uptake increased linearly during the first 10 min in the presence of 1 mM malate. During this period, CBT315(pKK223-3) cells accumulated only a background level of malate.




Figure 3. Immunolocalization of AgDCAT1 in *Frankia* sp.-infected alder nodules using anti-TGM13 antiserum. A, Bright-field image of a section of a whole alder nodule. B, Fluorescence image corresponding to A. C, Negative control nodule section from which the primary antiserum was omitted at the first incubation step. D, Enlarged image of a part of A enclosed in a white frame. E, Enlarged image of a part of B enclosed in a white frame. F, Enlarged image of a part of C enclosed in a white frame. G, Bright-field image of an infected cell filled with symbiotic vesicles. H, Fluorescence image corresponding to G. Bars in A to C = 500 μm , in D to F = 50 μm , and in G and H = 20 μm .

Based on this, all of the subsequent uptake assays lasted 10 min (Fig. 6B). The concentration-dependent malate uptake by CBT315(pAgDCAT1) showed saturation kinetics (Fig. 6C) with an apparent K_m of 70 μM according to the Lineweaver-Burk analysis (Fig. 6C, inset). This result indicates that AgDCAT1 mediates intermediate affinity malate transport, although it should be noted that the apparent K_m value may not reflect the true K_m within living plants.

To determine the substrate specificity of AgDCAT1, competition assays were performed by adding potential competing substrates in a 3-fold molar excess to a reaction mixture containing 0.5 mM malate. As shown in Figure 7, the addition of 1.5 mM unlabeled malate to the reaction mixture decreased the uptake of radiolabeled malate to about 25%. The inhibitory effects of fumarate and succinate were at

the same level as that of non-radiolabeled malate (Student's t test, $P = 0.07$ for succinate and 0.42 for fumarate). Oxaloacetate and pyruvate also competed with malate, reducing the uptake of radiolabeled malate to approximately 44% and 63%, respectively. Based on these results, and on the assumption of classical competitive inhibition and the K_m and V_{max} determined from the kinetics of malate transport (Fig. 6), the apparent binding constant (K_i) of the competing substrate was estimated in several trials so as to produce the fraction of residual malate transport as shown in Figure 7. Thus, the K_i value obtained was 145 and 315 μM for oxaloacetate and pyruvate, respectively. Other organic acids competed with malate only weakly. For example, malonate, an analog of L-malate, left 78% of the malate uptake intact, and citrate and oxalate inhibited only 8% (as a




Figure 4. Representative current traces from *AgDCAT1*-injected or noninjected oocytes perfused with a malate- or succinate-containing solution. A, Outward currents were generated from an *AgDCAT1*-injected oocyte upon treatment with 10 mM malate (pH 5.5) or 10 mM succinate (pH 5.5). When only the pH of the perfusion solution was changed from 7.4 to 5.5 (right), the injected oocyte developed a small inward current. B, No distinct response was observed from a noninjected oocyte. The oocytes were first perfused with 230 mM mannitol, 0.3 mM CaCl_2 , and 10 mM MES/Tris (pH 7.4), and then either with the same solution adjusted to pH 5.5 (right) or with 10 mM malate or succinate plus 220 mM mannitol, 0.3 mM CaCl_2 , and 10 mM MES/Tris (pH 5.5; left and middle).

consequence, their calculated K_i values were 0.66 mM for malonate and 2 mM for citrate and oxalate). Substrates of other members of the PTR family, namely, nitrate, and di-Gly, did not compete, nor did nitrogenous compounds such as Glu or allantoin. It is interesting that Glu, a dicarboxylate, did not compete with malate. We speculate that the transport of this large dicarboxylic acid may be abrogated by its charged amino group. Thus, our results can be summarized as follows: In addition to malate, *AgDCAT1* transports (or at least binds competitively at the transport-relevant site) several other C4-dicarboxylates, namely, succinate and fumarate with




Figure 5. Enhanced malate uptake by *AgDCAT1*-transformed yeast. DTY165 cells were transformed with pFL61-*AgDCAT1* (*AgDCAT1*) or pFL61 (control), and [^{14}C]malate uptake was measured for 2 min with 1 mM malate. Average values of two independent tests in triplicates are shown. Error bars = SE; $n = 6$.






Figure 6. Expression and characterization of *AgDCAT1* by using an *E. coli* *dct* mutant. A, Complementation test on M9 media with L-malic acid as a sole carbon source. The *E. coli* wild-type K-12(pKK223-3) and its *dct* mutant strain CBT315 that harbored p*AgDCAT1* grew on this medium, but CBT315(pKK223-3) did not. B, Time dependence of malate uptake at 1 mM malate (pH 5.9). Black circles, *AgDCAT1*-expressing CBT315(p*AgDCAT1*) cells; white circles, CBT315(pKK223-3) negative control. C, Rate of malate uptake over 10 min. Lineweaver-Burk plot analysis of the concentration-dependent uptake data revealed the K_m and V_{max} values to be 70 μM and 6.3 $\text{nmol min}^{-1} \text{mg}^{-1}$ protein, respectively (inset). Black circles, *AgDCAT1*-expressing CBT315(p*AgDCAT1*) cells; white circles, CBT315(pKK223-3) negative control. Assays were performed in triplicate and repeated four times. Error bars = SE; $n = 12$.

the same affinity, oxaloacetate with a 2-fold lower affinity, and malonate and oxalate with a 10- to 30-fold lower affinity.




Figure 7. Substrate specificity of AgDCAT1 in *E. coli*. [¹⁴C]Malate uptake in CBT315(pAgDCAT1) was measured with 0.5 mM malate (pH 5.9) in the presence of a 3-fold molar excess of non-labeled competing substrates. Mean values of three independent experiments in triplicate are shown. Error bars = SE; *n* = 9.

Malate Transport via AgDCAT1 Is Dependent on Membrane Potential

In the case of barley (*Hordeum vulgare*) mesophyll vacuoles (Martinoia et al., 1991), malate transport depends on the membrane potential. Thus, to determine whether the membrane potential could promote the AgDCAT1-mediated malate flux, CBT315(pAgDCAT1) cells were treated with the K⁺ ionophore valinomycin in the presence of different concentrations of extracellular K⁺. This method of altering the membrane potential is well established in *E. coli* (Ahmed and Booth, 1983; Li and Pajor, 2001). Malate uptake by CBT315(pAgDCAT1) in the presence of valinomycin was significantly lower at 5 mM of external K⁺, which should hyperpolarize the membrane potential, than at 50 or 100 mM external K⁺, which should depolarize the membrane potential (Fig. 8). A control experiment in which the osmolarity of the external solutions had been adjusted yielded similar results (data not shown). These data indicate that a depolarized membrane potential promotes dicarboxylate uptake via AgDCAT1.

DISCUSSION

In this paper, we have characterized AgDCAT1, a nodule-specific gene from alder that encodes a member of the PTR family of plasma membrane transporters. Its expression in *E. coli* showed that AgDCAT1 mediates the transport of dicarboxylates, including malate, succinate, fumarate, and oxaloacetate. Dicar-

boxylates have been suggested previously to be the carbon sources for *Frankia* sp. in the infected cells of alder nodules (Huss-Danell, 1997). We found that AgDCAT1 is specifically expressed in the alder nodules and that the protein product of the gene is localized at the symbiotic interface. Thus, AgDCAT1 appears to participate in the supply of carbon sources from alder to its nitrogen-fixing microsymbiont.

AgDCAT1 has similarities and differences to the dicarboxylate transporters found in the PBM of soybean symbiosomes. They both transport many different dicarboxylates and are energized by depolarized membrane potential, which suggests that they transport anionic forms of dicarboxylates. However, they differ somewhat in their substrate preference and in their affinity for various substrates. AgDCAT1 shows a similar affinity for malate, fumarate, and succinate. In contrast, the dicarboxylate transporter of PBM prefers malate and oxaloacetate over succinate and fumarate, and its affinity for the dicarboxylates is much lower than that of AgDCAT1 (Ou Yang et al., 1990). It would be very interesting to compare the structures of the dicarboxylate transporters of these different symbiotic systems when the dicarboxylate transporter of the soybean PBM is identified.

In vivo, the prevailing conditions favor malate efflux from the plant cell because malate concentrations are higher in the cell than in the apoplast, and the membrane potential is negative in the cytoplasmic side. So far, however, we have shown that AgDCAT1 participates in the uptake of malate into cells. To be able to use our findings to understand the physiologically relevant function of AgDCAT1, we need to postulate that AgDCAT1 is capable of the bidirectional transport of malate. Supporting this is that many transporters are not mechanically fixed in the direction of transport; rather, they are capable of bidirectional transport depending on the prevailing




Figure 8. Effect of membrane potential on the malate transport of AgDCAT1 in *E. coli*. CBT315(pAgDCAT1) cells were pretreated for 2 min with 50 μM valinomycin (Val) in the presence of 5, 50, or 100 mM external K⁺, and then the [¹⁴C]malate solution was added to a final malate concentration of 1 mM (pH 5.9). The assays were repeated three times in triplicate, and mean values were compared with the control ([K⁺] = 50 mM, without valinomycin) are shown. Error bars = SE; *n* = 9.

conditions (Yu and Choi, 1997; Sensi et al., 1997; Schwartz et al., 2002). To test if AgDCAT1 can export malate, we assayed malate efflux via AgDCAT1 from malate-preloaded *E. coli* and yeast. However, we could not detect any AgDCAT1-mediated efflux from these cells (data not shown), perhaps because these cells metabolize malate too efficiently. Thus, further experiments are necessary to determine whether AgDCAT1 is capable of exporting dicarboxylates. The heterologous expression of a zinc transporter from soybean nodules has resulted in a similar topological problem. When it was expressed in yeast, it catalyzed the uptake of Zn^{2+} , but when it was in the PBM, it catalyzed the export of Zn from the cytoplasm (Moreau et al., 2002). It was suggested that this Zn transporter is capable of bidirectional transport and that the direction of the transport is determined by the membrane potential and the concentration gradient of the substrate. This may also be true for AgDCAT1. The negative membrane potential of the plasma membrane of the infected cell, which is the interface between the host and the actinorhizal symbiont, should inhibit malate uptake into the infected cell and should instead promote malate efflux from the cell to the microsymbiont. Consistent with this explanation is our experiment using valinomycin in combination with low external K^+ , which showed that hyperpolarization of the membrane potential decreased malate uptake via AgDCAT1 (Fig. 8). In addition, the concentration gradient of dicarboxylates, which is established by the uptake of malate by the microsymbiont, should promote the malate efflux from the cell.

Because *AgDCAT1* is expressed specifically in nodules, the question arises whether it evolved from a gene for a transporter with different substrate specificity or whether it represents a nodule-specific duplication product of a malate transporter gene expressed elsewhere. *AgDCAT1* is not homologous to any known dicarboxylate transporter, including a plant mitochondrial di- and tricarboxylate transporter (Grobler et al., 1995; Fiermonte et al., 1999; Palmieri et al., 2000; Jausch et al., 2002; Picault et al., 2002). Because other PTR family dicarboxylate transporters have not been identified to date, we suggest that AgDCAT1 may have evolved from a PTR transporter with another substrate specificity to accommodate the need for a dicarboxylate transporter in nodule symbiosis.

To our knowledge, AgDCAT1, a nodule-specific transporter from alder, is the first plasma membrane dicarboxylate transporter that has been characterized in higher plants. Although the molecular details of the dicarboxylate transport via AgDCAT1 remain to be understood, the present data suggest that AgDCAT1 is the first transporter to be identified that is likely to be responsible for supplying an intracellular nitrogen-fixing microsymbiont with carbon sources. Thus, our study improves the understanding of the nutrient exchange that takes place during

root nodule symbioses and sheds new light on the diversity of substrates that are transported by members of the PTR family.

MATERIALS AND METHODS

Plant Material

Alder (*Alnus glutinosa*) seeds were collected from a local source (Weerribben, The Netherlands). The plants were grown in a greenhouse at 25°C under cycles of 16 h of light and 8 h of dark. Seeds were germinated in trays containing sterile gravel that were wetted with sterile tap water. After 3 weeks, the seedlings were transferred to sterile gravel that had been wetted with one-quarter-strength Hoagland solution (Hoagland and Arnon, 1938), and each plantlet was infected with 1 mL of a 1:5 diluted dispersed culture of *Frankia* sp. HFPAr13 (Berry and Torrey, 1979) that had been grown in phosphorus medium without nitrogen (Meesters et al., 1985). The nodules were harvested 6 to 8 weeks after infection. For control experiments, roots were harvested from noninoculated seedlings 2 to 3 weeks after germination.

Isolation of Nucleic Acids, Cloning, and Sequencing Procedures

DNA and RNA isolation from alder were performed as described previously (Ribeiro et al., 1995). DNA manipulations were carried out as described by Sambrook et al. (1989). Nucleotide sequences were determined using an automatic sequencer (model 373A, Applied Biosystems, Foster City, CA). Sequence data were analyzed using the programs of the Wisconsin Genetics Computer Group (Devereux et al., 1984). Database searches were performed by using the BLAST algorithm (Altschul et al., 1990) on the nucleotide sequence databases of the National Center of Biotechnology Information, National Library of Medicine, National Institutes of Health (Bethesda, MD).

Nucleic Acid Hybridization Methods

Total RNA was denatured in dimethyl sulfoxide (DMSO)/glyoxal and electrophoresed on 1.2% (w/v) agarose gels (Sambrook et al., 1989). DNA was separated on 0.8% (w/v) agarose gels (Sambrook et al., 1989). Nucleic acids were transferred to GeneScreen (New England Nuclear, Boston) filters (RNA) or to Amersham Hybond N^+ (Amersham, Little Chalfont, UK) filters (DNA). RNA-blot hybridizations were performed in buffer containing 50% (w/v) formamide at 42°C (Sambrook et al., 1989). DNA-blot hybridizations were performed according to the protocol provided by the manufacturer for Amersham Hybond N^+ . Filters were washed at 65°C with decreasing salt concentrations down to $0.5\times$ SSC and 0.1% (w/v) SDS.

Western Blot of AgDCAT1

Two anti-AgDCAT1 antisera were produced by Peptron (Taejun, South Korea) by immunizing rabbits with peptides corresponding to the C terminus of AgDCAT1 (CEDAHQKINGKEEKV, denoted as CED15) or the loop between transmembrane domains 6 and 7 (TGMDRGADGLTIS, denoted as TGM13). Plasma membrane proteins were isolated from 10 g of alder nodules using a two-phase system according to Robinson and Hinz (2001), concentrated using 72% (w/v) trichloroacetic acid, and separated on SDS-polyacrylamide gels. Protein transfer and immunodetection were performed as described by Sambrook et al. (1989).

Immunolocalization of AgDCAT1 in Alder Nodule

The fixation of nodules was carried out according to Stadler et al. (1995). Parts of nodules were briefly degassed in 3 mL of fixative (3:1 [v/v] ethanol:acetic acid), and the tissues were fixed at room temperature for 1 h. After three 30-min washes in 70% (v/v) ethanol containing 1 mM dithiothreitol (DTT) and one overnight washing step, the tissues were dehydrated on ice with 80%, 85%, 90%, and 95% (v/v) ethanol containing 1 mM DTT for 20 min with each solution and finally incubated twice for 20 min in 99.8% (v/v) ethanol containing 10 mM DTT. The tissue was infiltrated with methacrylate in three sequential incubations at 4°C using a methacrylate mix (75% [v/v] butyl methacrylate, 25% [v/v] methyl methacrylate, 0.5%

[w/v] benzoine ethyl ether, and 10 mM DTT) at increasing methacrylate mix to ethanol ratios (first incubation, overnight, 1:2 [w/v]; second incubation, 6 h, 1:1 [w/v]; and third incubation, overnight, 2:1 [w/v]). After one additional incubation for 6 h and two final overnight incubations at 4°C, all in 100% (w/v) methacrylate mix, the samples were transferred to ultrathin PCR tubes. The methacrylate was polymerized by incubation at 4°C for 15 h under UV light (310 nm) in 100% (v/v) methacrylate mix. Semithin sections (2 µm) were cut with an ultramicrotome (Ultracut R, Leica, Bensheim, Germany) and placed on poly-L-Lys-coated slides.

To remove the methacrylate from the semithin sections, the coverslips were incubated for 2 min in 100% (v/v) acetone. Rehydration involved incubations with an ethanol series (100%, 70%, and 30% [v/v]) for 30 s each. The coverslips were then washed with Tris-buffered saline buffer (50 mM Tris-HCl [pH 7.5] and 150 mM NaCl) for 30 s and incubated in blocking buffer (1% [w/v] nonfat milk powder in Tris-buffered saline) for 45 min. After an overnight incubation with anti-TGM13 antiserum (diluted 1:100 or 1:500 [v/v] in blocking buffer), the coverslips were washed three times with blocking buffer and incubated for 1 h with the anti-rabbit IgG-fluorescein isothiocyanate isomer 1 conjugate (Sigma, St. Louis; diluted 1:300 [v/v] in blocking buffer). After five final 5-min washes with blocking buffer, the coverslips were rinsed with water and mounted with the 10-µL ProLong-Antifade Kit (Molecular Probes, Leiden, Netherlands). Photographs were taken with a fluorescence microscope (Carl Zeiss, Axioskope 2, Zeiss, Göttingen, Germany) equipped with filter blocks for excitation at 455 to 495 nm and emission at 505 to 555 nm.

Functional Characterization of AgDCAT1 in *Xenopus laevis* Oocytes

AgDCAT1 was functionally expressed in oocytes and was performed as described by Tsay et al. (1993). Female *X. laevis* were obtained from Xenopus I (Merkle Road, Dexter, MI). The AgDCAT1 coding region from the original cDNA clone was subcloned in the EcoRI site of the *X. laevis* oocyte expression vector pGEMHE (Liman et al., 1992). pGEMHE-AgDCAT1 was linearized with *NheI*, and capped cRNA was transcribed in vitro using the T7 RNA transcription kit (Ambion, Austin, TX). Fifty nanoliters of the cRNA (1 µg µL⁻¹) were injected into collagenase-digested oocytes. Current measurements were performed 2 to 3 d after injection.

The conventional two-electrode voltage clamp technique was performed using a TEV200A amplifier (Dagan, Minneapolis). Oocytes were incubated in a solution containing 230 mM mannitol, 0.3 mM CaCl₂, and 10 mM MES/Tris (pH 7.4), then perfused with 10 mM malate or another substrate plus 220 mM mannitol, 0.3 mM CaCl₂, and 10 mM MES/Tris (pH 5.5 or 7.4). The oocytes were held at -60 mV, and the currents were recorded and measured using Axotape (Axon Instruments, Inc., Union City, CA).

Expression and [¹⁴C]Malate Transport in Yeast

AgDCAT1 was cloned into the *NotI* site of pFL61 (Minet et al., 1992) to produce pFL61-AgDCAT1, and this was introduced into yeast (*Saccharomyces cerevisiae*) strain DTY165 using the lithium acetate method (Schiestl and Gietz, 1989). As a negative control, pFL61 vector alone was transformed into the same strain. DTY165 cells expressing AgDCAT1 or vector alone were cultured in SD-*ura* broth and harvested at mid-log phase. The cells were then washed twice with 100 mM potassium phosphate buffer (pH 5.9), resuspended in the same buffer to a final concentration of 1 mg protein mL⁻¹, and kept on ice until the uptake assays. [¹⁴C]Malate transport assays were performed according to Camarasa et al. (2001), with modifications. Thus, the cell suspensions were pre-incubated at 28°C for 2 min; then, uptake was initiated by adding L-[1,4(2,3)-¹⁴C] malic acid (Amersham) diluted with 1 mM cold malate to a specific activity of 0.1 µCi nmol⁻¹. After 2 min, the reaction was stopped by adding ice-cold 100 mM LiCl. The samples were immediately filtered through 0.45-µm nitrocellulose membrane filters, washed with 4 mL of ice-cold 100 mM LiCl, and incubated in scintillation cocktail solution (Research Products International Corp., Mount Prospect, IL). The radioactivity was measured by using a liquid scintillation counter (Tri-CARB2100TR, Packard Bioscience Co., Meriden, CT).

Complementation of *Escherichia coli*

E. coli K-12 (DCT) and its dicarboxylate transport mutant strain CBT315 (CGSC5269) were obtained from the *E. coli* Genetic Stock Center (Yale

University, New Haven, CT). AgDCAT1 cDNA was cloned into the EcoRI site of the pKK223-3 vector, yielding pAgDCAT1, and CBT315 then was transformed with pAgDCAT1. The pKK223-3 vector alone was transformed into CBT315 as a negative control and into K-12 as a positive control. The phenotypes were compared on M9 medium (Sambrook et al., 1989) with malate as the sole carbon source.

[¹⁴C]Malate Transport Assays in *E. coli*

[¹⁴C]Malate transport assays were performed as described (Labarre et al., 1996) with modifications. CBT315(pKK223-3) and CBT315(pAgDCAT1) cells were cultured to mid-log phase in Luria-Bertani medium, washed twice with 50 mM potassium phosphate buffer (pH 5.9), and 0.75 mg protein mL⁻¹ was resuspended in the same buffer for the assays. After pre-incubation for 3 min at 30°C, uptake was initiated by adding L-[1,4(2,3)-¹⁴C] malate] diluted with the appropriate concentration of cold malate to a specific activity of 0.05 to 0.1 µCi nmol⁻¹. Further steps were performed as described in the assays in yeast.

The K_m and V_{max} values for malate were determined according to the Lineweaver-Burk plot based on AgDCAT1-mediated malate uptake at external concentrations of 0.025 to 10 mM. To test the effect of membrane potential on malate uptake via AgDCAT1, membrane potential was clamped at different levels using valinomycin at different concentrations of extracellular K⁺. After the pre-incubation step, the cells were treated with valinomycin in different concentrations of potassium phosphate buffer (pH 5.9) for 2 min; then, uptake was initiated as described above. Valinomycin was dissolved to a final concentration of 50 µM in 0.1% (v/v) DMSO. This concentration is in the range used for experiments with bacteria (Lee et al., 1999; Fischer et al., 2000). DMSO (0.1% [v/v]) alone did not have any effect on malate uptake. Cell viability was tested by the BacLight bacteria kit (Molecular Probes, Eugene, OR), and it did not differ between control cells and cells treated with valinomycin.

Distribution of Materials

Upon request, all novel materials described in this publication will be made available in a timely manner for noncommercial research purposes, subject to the requisite permission from any third party owners of all or parts of the material. Obtaining any permissions will be the responsibility of the requestor.

ACKNOWLEDGMENTS

We thank Tony van Kampen (Department of Molecular Biology, Agricultural University, Wageningen, The Netherlands) for DNA sequencing and Gary Stacey and Serry Koh (Department of Microbiology, University of Tennessee, Knoxville) for testing AgDCAT1 for peptide transport activity in yeast. We thank Christian Knop (Göttingen University, Göttingen, Germany) for help with protein isolation and immunodetection. We would also like to thank Arie Moran (Department of Physiology, Ben Gurion University, Beer-Sheva, Israel) for helpful suggestions and comments.

Received August 22, 2003; returned for revision October 12, 2003; accepted December 23, 2003.

LITERATURE CITED

- Ahmed S, Booth IR (1983) The use of valinomycin, nigericin and trichloro-carbanilide in control of the proton motive force in *Escherichia coli* cells. *Biochem J* 212: 105–112
- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ (1990) Basic local alignment search tool. *J Mol Biol* 215: 304–310
- Arima K, Hines ER, Kiela PR, Drees JB, Collins JF, Ghishan FK (2002) Glucocorticoid regulation and glycosylation of mouse intestinal type IIb Na-P-i cotransporter during ontogeny. *Am J Physiol* 283: G426–G434
- Berry AM, Torrey JG (1979) Isolation and characterization *in vivo* and *in vitro* of an actinomycetous endophyte from *Alnus rubra* Bong. In JC Gordon, DT Wheeler, DA Perry, eds, *Symbiotic Nitrogen Fixation in the Management of Temperate Forests*. Oregon State University, Corvallis, pp 69–84
- Brosius J, Holy A (1984) Regulation of ribosomal RNA promoters with a synthetic lac operator. *Proc Natl Acad Sci USA* 81: 6929–6933

- Camarasa C, Bidard F, Bony M, Barre P, Dequin S** (2001) Characterization of *Schizosaccharomyces pombe* malate permease by expression in *Saccharomyces cerevisiae*. *Appl Environ Microbiol* **67**: 4144–4151
- Dean RM, Rivers RL, Zeidel ML, Roberts DM** (1999) Purification and functional reconstitution of soybean nodulin 26: an aquaporin with water and glycerol transport properties. *Biochemistry* **38**: 347–353
- de Faria SM, McInroy SG, Sprent JI** (1987) The occurrence of infected cells, with persistent infection threads in legume root nodules. *Can J Bot* **65**: 553–558
- Devereux J, Haeberli P, Smithies O** (1984) A comprehensive set of sequence analysis programs for the VAX. *Nucleic Acids Res* **12**: 387–395
- Fiermonte G, Dolce V, Arrigoni R, Runswick MJ, Walker JE, Palmieri F** (1999) Organization and sequence of the gene for the human mitochondrial dicarboxylate carrier: evolution of the carrier family. *Biochem J* **344**: 953–960
- Fischer S, Graber P, Turina P** (2000) The activity of the ATP synthase from *Escherichia coli* is regulated by the transmembrane proton motive force. *J Biol Chem* **275**: 30157–30162
- Frommer WB, Hummel S, Rentsch D** (1994) Cloning of an *Arabidopsis* histidine transporting protein related to nitrate and peptide transporters. *FEBS Lett* **347**: 185–189
- Grobler J, Bauer F, Subden RE, Van Vuuren HJ** (1995) The *mae1* gene of *Schizosaccharomyces pombe* encodes a permease for malate and other C4 dicarboxylic acids. *Yeast* **11**: 1485–1491
- Hoagland DR, Arnon DT** (1938) The water-culture method for growing plants without soil. University of California, Berkeley, California Agriculture Experiment Station Circular 347
- Husain J, Lo R, Grbavec D, Stifani S** (1996) Affinity for the nuclear compartment and expression during cell differentiation implicate phosphorylated Groucho/TLE1 forms of higher molecular mass in nuclear functions. *Biochem J* **317**: 523–531
- Huss-Danell K** (1997) Actinorhizal symbioses and their N₂ fixation. *New Phytol* **136**: 375–405
- Janausch IG, Zientz E, Tran QH, Kröger A, Unden G** (2002) C4-dicarboxylate carriers and sensors in bacteria. *Biochim Biophys Acta* **1553**: 39–56
- Kim EJ, Kwak JM, Uozumi N, Schroeder JI** (1998) AtKUP1: an Arabidopsis gene encoding high-affinity potassium transport activity. *Plant Cell* **10**: 51–62
- Kouchi H, Hata S** (1993) Isolation and characterization of novel nodulin cDNAs representing genes expressed at early stages of soybean nodule development. *Mol Gen Genet* **238**: 106–119
- Labarre C, Guzzo J, Cavin J-F, Divies C** (1996) Cloning and characterization of the genes encoding the malolactate enzyme and the malate permease of *Leuconostoc oenos*. *Appl Environ Microbiol* **62**: 1274–1282
- Lee AI, Delgado A, Gunsalus RP** (1999) Signal-dependent phosphorylation of the membrane-bound NarX two-component sensor-transmitter protein of *Escherichia coli*: nitrate elicits a superior anion ligand response compared to nitrite. *J Bacteriol* **181**: 5309–5316
- Li H, Pajor AM** (2001) Functional characterization of CitM, the Mg²⁺-citrate transporter. *J Membr Biol* **185**: 9–16
- Liman ER, Tytgat J, Hess P** (1992) Subunit stoichiometry of a mammalian K⁺ channel determined by construction of multimeric cDNAs. *Neuron* **9**: 861–871
- Lin F, Lester HA, Mager S** (1996) Single-channel currents produced by the serotonin transporter and analysis of a mutation affecting ion permeation. *Biophys J* **71**: 3126–3135
- Lo TCY, Rayman MK, Sanwal BD** (1972) Transport of succinate in *Escherichia coli*: I. Biochemical and genetic studies of transport in whole cells. *J Biol Chem* **247**: 6323–6331
- Martinoia E, Vogt E, Rentsch D, Amrhein N** (1991) Functional reconstitution of the malate carrier of barley mesophyll vacuoles in liposomes. *Biochim Biophys Acta* **1062**: 271–278
- Meesters TM, van Genesen ST, Akkermans ADL** (1985) Growth, acetylene reduction activity and localization of nitrogenase in relation to vesicle formation in *Frankia* strains Cc1.17 and Cp1.2. *Arch Microbiol* **143**: 137–179
- Minet M, Dufour ME, Lacroute F** (1992) Complementation of *Saccharomyces cerevisiae* auxotrophic mutants by *Arabidopsis thaliana* cDNAs. *Plant J* **2**: 417–422
- Moreau S, Thomson RM, Kaiser BN, Trevaskis B, Guerinot ML, Udvardi MK, Puppo A, Day DA** (2002) GmZIP1 encodes a symbiosis-specific zinc transporter in soybean. *J Biol Chem* **277**: 4738–4746
- Mylona P, Pawlowski K, Bisseling T** (1995) Symbiotic nitrogen fixation. *Plant Cell* **7**: 869–885
- Nicholas KB, Nicholas HB Jr, Deerfield DW II** (1997) GeneDoc: analysis and visualization of genetic variation. *EMBnet NEWS* **4**: 14
- Ou Yang LJ, Udvardi MK, Day DA** (1990) Specificity and regulation of the dicarboxylate carrier on the peribacteroid membrane of soybean nodules. *Planta* **182**: 437–444
- Ou Yang LJ, Whelan J, Weaver CD, Roberts DM, Day DA** (1991) Protein phosphorylation stimulates the rate of malate uptake across the peribacteroid membrane of soybean nodules. *FEBS Lett* **293**: 188–190
- Palmieri L, Runswick MJ, Fiermonte G, Walker JE, Palmieri F** (2000) Yeast mitochondrial carriers: bacterial expression, biochemical identification and metabolic significance. *J Bioenerg Biomembr* **32**: 67–77
- Pawlowski K, Bisseling T** (1996) Rhizobial and actinorhizal symbioses: what are the shared features? *Plant Cell* **8**: 1899–1913
- Picault P, Palmieri L, Pisano I, Hodges M, Palmieri F** (2002) Identification of a novel transporter for dicarboxylates and tricarboxylates in plant mitochondria. *J Biol Chem* **277**: 24204–24211
- Ribeiro A, Akkermans ADL, van Kammen A, Bisseling T, Pawlowski K** (1995) A nodule-specific gene encoding a subtilisin-like protease is expressed in early stages of actinorhizal nodule development. *Plant Cell* **7**: 785–794
- Rivers RL, Dean RM, Chandy G, Hall JE, Roberts DM, Zeidel ML** (1997) Functional analysis of nodulin 26, an aquaporin in soybean root nodule symbiosomes. *J Biol Chem* **272**: 16256–16261
- Robinson DG, Hinz G** (2001) Organelle isolation. *In* B Satiat-Jeunemaitre, C Hawes, eds, *Plant Cell Biology: A Practical Approach*, Ed 2. IRL Press, Oxford, pp 295–324
- Sambrook J, Fritsch E, Maniatis T** (1989) *Molecular Cloning: A Laboratory Manual*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY
- Schiestl RH, Gietz RG** (1989) High efficiency transformation of intact yeast cells using single stranded nucleic acids as a carrier. *Curr Genet* **16**: 339–346
- Schwartz GJ, Tsuruoka S, Vijayakumar S, Petrovic S, Mian A, Al-Awgati Q** (2002) Acid incubation reverses the polarity of intercalated cell transporters, an effect mediated by hensin. *J Clin Invest* **109**: 89–99
- Sensi SL, Canzoniero LMT, Yu SP, Ying HS, Koh J-Y, Kerchner GA, Choi DW** (1997) Measurements of intracellular free Zn in living cortical neurons: routes of entry. *J Neurosci* **17**: 9554–9564
- Sonders MA, Amara SG** (1996) Channels in transporters. *Curr Opin Neurobiol* **6**: 294–302
- Sonders MS, Zhu S-J, Zahniser NR, Kavanaugh MP, Amara SG** (1997) Multiple ionic conductances of the human dopamine transporter: the action of dopamine and psychostimulants. *J Neurosci* **17**: 960–974
- Stadler R, Wolf K, Hilgarth C, Tanner W, Sauer N** (1995) Subcellular localization of the inducible *Chorella* HUP1 monosaccharide-H⁺ symporter and cloning of a co-induced galactose-H⁺ symporter. *Plant Physiol* **107**: 33–41
- Steiner HY, Naider F, Becker JM** (1995) The PTR family: a new group of peptide transporters. *Mol Microbiol* **16**: 825–834
- Streeter JG** (1995) Recent developments in carbon transport and metabolism in symbiotic systems. *Symbiosis* **19**: 175–196
- Takagi T** (1991) New aspects of electrophoresis of membrane proteins: electrophoretic light scattering and polyacrylamide gel electrophoresis in the presence of sodium dodecyl sulfate analogues. *In* A Chrambach, MJ Dunn, BJ Radola, eds, *Advances in Electrophoresis*, Vol 4. VCH Publishers, New York, pp 391–406
- Tsay Y-F, Schroeder JI, Feldmann KA, Crawford NM** (1993) The herbicide sensitivity gene *CHL1* of *Arabidopsis* encodes a nitrate-inducible nitrate transporter. *Cell* **12**: 705–713
- Tyerman SD, Whitehead LF, Day DA** (1995) A channel-like transporter for NH₄⁺ on the symbiotic interface of N₂-fixing plants. *Nature* **378**: 629–632
- Uozumi N** (2001) *E. coli* as an expression system for K⁺ transport systems from plants. *Am J Physiol* **281**: C733–C739
- Verma DPS** (1992) Signals in root nodule organogenesis and endocytosis of *Rhizobium*. *Plant Cell* **4**: 1039–1057
- Williams L, Miller A** (2001) Transporters responsible for the uptake and partitioning of nitrogenous solutes. *Annu Rev Plant Physiol Plant Mol Biol* **52**: 659–688
- Yu SP, Choi DW** (1997) Na⁺/Ca²⁺ exchange currents in cortical neurons: concomitant forward and reverse operation and effect of glutamate. *Eur J Neurosci* **9**: 1273–1281