

# An Auxin Transport Independent Pathway Is Involved in Phosphate Stress-Induced Root Architectural Alterations in Arabidopsis. Identification of *BIG* as a Mediator of Auxin in Pericycle Cell Activation<sup>1</sup>

José López-Bucio<sup>2</sup>, Esmeralda Hernández-Abreu<sup>2</sup>, Lenin Sánchez-Calderón, Anahí Pérez-Torres, Rebekah A. Rampey, Bonnie Bartel, and Luis Herrera-Estrella\*

Departamento de Ingeniería Genética, Unidad Irapuato del Centro de Investigación y Estudios Avanzados del Instituto Politécnico Nacional, 36500 Irapuato, Guanajuato, Mexico (J.L.-B., E.H.-A., L.S.-C., A.P.-T., L.H.-E.); and Department of Biochemistry and Cell Biology, Rice University, Houston, Texas 77005–1892 (R.A.R., B.B.)

*Arabidopsis* (*Arabidopsis thaliana*) plants display a number of root developmental responses to low phosphate availability, including primary root growth inhibition, greater formation of lateral roots, and increased root hair elongation. To gain insight into the regulatory mechanisms by which phosphorus (P) availability alters postembryonic root development, we performed a mutant screen to identify genetic determinants involved in the response to P deprivation. Three *low phosphate-resistant root* lines (*lpr1-1* to *lpr1-3*) were isolated because of their reduced lateral root formation in low P conditions. Genetic and molecular analyses revealed that all *lpr1* mutants were allelic to *BIG*, which is required for normal auxin transport in *Arabidopsis*. Detailed characterization of lateral root primordia (LRP) development in wild-type and *lpr1* mutants revealed that *BIG* is required for pericycle cell activation to form LRP in both high (1 mM) and low (1  $\mu$ M) P conditions, but not for the low P-induced alterations in primary root growth, lateral root emergence, and root hair elongation. Exogenously supplied auxin restored normal lateral root formation in *lpr1* mutants in the two P treatments. Treatment of wild-type *Arabidopsis* seedlings with brefeldin A, a fungal metabolite that blocks auxin transport, phenocopies the root developmental alterations observed in *lpr1* mutants in both high and low P conditions, suggesting that *BIG* participates in vesicular targeting of auxin transporters. Taken together, our results show that auxin transport and *BIG* function have fundamental roles in pericycle cell activation to form LRP and promote root hair elongation. The mechanism that activates root system architectural alterations in response to P deprivation, however, seems to be independent of auxin transport and *BIG*.

Phosphorus (P) is one of the most limiting nutrients for plant growth in many natural and agricultural ecosystems. In response to low P availability in the soil, plants have developed several adaptive strategies to increase P acquisition, including enhanced expression of P transport genes, exudation of P-solubilizing substances, such as organic acids and phosphatases, and alterations in postembryonic root development (for review, see Raghothama, 1999; López-Bucio et al., 2000; Abel et al., 2002; López-Bucio et al., 2003; Vance et al., 2003).

Plants typically respond to P deficiency by allocating more carbon to roots, thus increasing their root-

to-shoot ratio. In addition, low P availability can dramatically alter the spatial configuration of the root system by increasing root hairs and promoting lateral root formation. Such plastic root alterations are believed to play a crucial role in exploring increased soil volumes in search of nutrient-rich patches (Neumann and Martinoia, 2002; López-Bucio et al., 2003). In *Lupinus albus*, P deficiency promotes the formation of cluster roots, also termed proteoid roots (Johnson et al., 1996). Increased root branching in response to P limitation has also been reported for *Arabidopsis* (*Arabidopsis thaliana*). Growth of *Arabidopsis* in a moderate-to-limiting supply of P results in a redistribution of root growth from the primary to lateral roots (Williamson et al., 2001). Reduced primary root elongation in low P conditions was accompanied by increased root branching, perhaps concentrating root biomass near the soil surface for more efficient nutrient foraging (López-Bucio et al., 2002, 2003).

Despite several detailed anatomical studies on root architectural alterations in response to P limitation, little is known about the physiological and molecular mechanisms that coordinate these developmental changes. Phytohormones control most root system characteristics in angiosperms, including primary

<sup>1</sup> This work was supported in part by the Howard Hughes Medical Institute (grant no. Nbr55003677), by the European Commission (grant no. ICA-4-CT2000-30017), and by the Robert A. Welch Foundation (C-1309).

<sup>2</sup> Present address: Instituto de Investigaciones Químico-Biológicas, Universidad Michoacana de San Nicolás de Hidalgo. Edificio B3, Ciudad Universitaria, C.P. 58030 Morelia, Michoacán, México.

\* Corresponding author; e-mail lherrera@ira.cinvestav.mx; fax 462-624-58-46.

Article, publication date, and citation information can be found at [www.plantphysiol.org/cgi/doi/10.1104/pp.104.049577](http://www.plantphysiol.org/cgi/doi/10.1104/pp.104.049577).

root growth and the formation of lateral roots. For example, auxin, ethylene, and cytokinin are thought to play important roles in P deficiency-induced alterations in lateral root development and cluster root development (López-Bucio et al., 2003; Vance et al., 2003). In Lupin and Arabidopsis, an important role for auxins in the formation of lateral roots in response to P limitation has been suggested. In low P conditions, the auxin transport inhibitors 2,3,5-triiodobenzoic acid (TIBA) and *N*-(1-naphthyl) phthalamic acid (NPA) inhibit the formation of proteoid or lateral roots in Lupin and Arabidopsis, respectively (Gilbert et al., 2000; Skene and James, 2000; López-Bucio et al., 2002). In addition, auxin treatments of P-starved plants and gene expression analysis support the conclusion that auxin signaling is involved in the adaptive response of the root system architecture to P limitation (López-Bucio et al., 2002; Al-Ghazi et al., 2003).

Although these data collectively support the hypothesis that auxin is involved in the responses of the root system architecture to P limitation, several mutants defective in either auxin transport or signaling display normal changes in root system architecture in response to P availability. For example, the auxin transport mutants *aux1* and *eir1* and the auxin response mutants *axr1* and *axr4*, which have reduced lateral root formation in P-sufficient conditions, show typical primary root growth inhibition and enhanced production of lateral roots on low P media (Williamson et al., 2001; López-Bucio et al., 2002). To date, it is not clear whether changes in local concentration, transport, or sensitivity to auxin play important roles in the root architectural changes that occur during P deprivation.

To elucidate the low P-signaling pathway responsible for alterations in root development, we conducted a visual screen for Arabidopsis mutants with reduced lateral root formation when grown in P-limiting conditions. Here, we report the isolation of mutants that have reduced lateral root formation in low P conditions. Recombination mapping identified the gene defective in the mutants as *BIG*, a gene required for normal auxin transport (Ruegger et al., 1997; Gil et al., 2001) and lateral root development (Ruegger et al., 1997). Analysis of root system architecture and *DR5:uidA* expression in wild-type and *BIG* plants in high and low P conditions indicates that low P-induced alterations in postembryonic root development are likely independent of auxin transport and *BIG* function. Moreover, our results point to a key role of *BIG* in pericycle cell activation to form lateral root primordia (LRP), a process that is not modified by P availability but is required for increased LRP emergence in low P conditions.

## RESULTS

### Isolation of Arabidopsis Mutants with Altered Lateral Root Formation in Response to Low P Conditions

To identify mutants that are defective in lateral root formation in low P conditions, we screened ethyl


methane sulfonate (EMS) and T-DNA insertion mutant collections (Krysan et al., 1999) by observing the root architecture of plants growing over the surface of agar plates with low (1  $\mu$ M) P content. Three mutant lines showing a reduced number and density of lateral roots during P deprivation were identified (Fig. 1, A and B). Other aspects of root architecture induced by low P, such as primary root growth inhibition and root hair growth stimulation, were apparently unaffected in the mutants. The mutants were back-crossed to either wild-type Columbia (Col-0) or Wassilewskija (Ws) four times prior to detailed phenotypic analyses. In F<sub>2</sub> progeny from these crosses, the 3 lines segregated the mutant phenotype in a 1:3 ratio, indicating that each resulted from a single recessive nuclear mutation. All three mutants appeared to belong to the same phenotypic class, showing reduced primary root growth, absence of lateral roots, and very short root hairs in high (1 mM) P conditions (Fig. 1C). Complementation tests revealed that the three lines were allelic (data not shown). We named this locus *low phosphate-resistant root 1* (*lpr1*), and the three alleles were designated *lpr1-1* (Col-0), *lpr1-2* (Ws), and *lpr1-3* (Col-0). Adult *lpr1* mutants had shoot developmental alterations (Fig. 1, D–F). Mutant plants have a compact rosette caused by a reduction in leaf petiole length and smaller, round leaves (Fig. 1E) compared to wild type (Fig. 1D). At later stages of development, *lpr1* plants showed reduced apical dominance (Fig. 1F). Aside from these morphological alterations, *lpr1* mutants were fertile and produced fully viable seed (data not shown). Since the three *lpr1* alleles appeared to be phenotypically indistinguishable, we performed most of our detailed studies on *lpr1-1* (Col-0) and *lpr1-2* (Ws).

### Recombinant Mapping of *lpr1* Locus: *lpr1* Is Allelic to *doc1/tir3/big*

The chromosomal location of the *lpr1* locus was mapped genetically. We out-crossed an *lpr1* mutant plant (Col-0 background) to Ws. DNA from F<sub>2</sub> progeny with the *lpr1* mutant phenotype was analyzed for linkage to published markers (Konieczny and Ausubel, 1993; Bell and Ecker, 1994). The *lpr1* mutation mapped to the top of chromosome 3, north of *nga172* (Fig. 2). The *doc1/tir3* locus that encodes *BIG*, a calossin-like protein required for polar auxin transport in Arabidopsis, also maps to this region (Gil et al., 2001). To test the possibility that *lpr1* mutants were allelic to *doc1/tir3*, we crossed *lpr1-1* with *doc1* (Li et al., 1994). F<sub>1</sub> plants from this cross had *lpr1* mutant phenotypes (data not shown), indicating that *lpr1* mutants represent new mutant alleles of *BIG*.


### Effects of Phosphate Availability on Root System Architecture in Wild Type and *BIG* Mutants

To more clearly define the alterations in the root architectural response to low P caused by mutations in



**Figure 1.** *lpr1-1* mutant characterization. Twelve-day-old wild-type (Col-0; A) and *lpr1-1* (B) seedlings growing on media containing low P. C, Eighteen-day-old Col-0 (left) and *lpr1-1* (right) seedlings growing in high P. Note that no lateral roots are visible in the mutant at this advanced stage of seedling development. Col-0 (D) and *lpr1-1* (E) grown under 16-h photoperiod for 3 weeks at 22°C/18°C. F, Mature Col-0 and *lpr1-1* plants.


*BIG*, we performed temporal and single-point measurements of primary root growth, lateral root density, and root hair length. To test whether mutations in *BIG* could alter primary root growth in response to varying P availability, a kinetic analysis of primary root growth in wild-type and *lpr1-2* seedlings was carried out in high and low P conditions. As previously reported for *tir3* (Ruegger et al., 1997), primary roots in the *lpr1-2* mutants were 30% to 40% shorter than wild type at 12 d in high P conditions (Fig. 3A). In low P conditions, the primary roots of both wild-type and mutant plants



**Figure 2.** Recombinant mapping of *lpr1-1*. The *lpr1-1* mutation was localized to the top arm of chromosome 3 using published markers (Konieczny and Ausubel, 1993; Bell and Ecker, 1994). The *lpr1* mutation was mapped to north of *nga172*, a region that includes the *BIG* locus (Ruegger et al., 1997; Gil et al., 2001). Fractions represent the number of recombinant chromosomes over the total number of chromosomes scored.

were similar, growing for 6 to 9 d before arresting at lengths of only 1.5 to 1.8 cm (Fig. 3A). In addition, root hair length on primary roots of wild-type and *lpr1-2* plants were measured following 10 d of growth on high and low P media. In high P conditions, in *lpr1-2*, root hairs were 50% shorter than wild type (Fig. 3B). In contrast, low P conditions restored root hair elongation to wild-type levels (Fig. 3B). Similar results were obtained for *doc1* (data not shown).

To further analyze the effects of low P availability on lateral root development and the effects of mutations in *BIG* on this process, the density of emerged lateral roots in wild-type and *lpr1-2* seedlings was determined. After 15 d of growth in low P conditions, wild-type seedlings showed a highly branched root system harboring second- and third-order lateral roots (data not shown), with a 5- to 6-fold increased density of lateral roots (Fig. 3C). At this stage, in high P conditions, first-order lateral roots had emerged from the primary root. As previously reported for *tir3* (Ruegger et al., 1997), *lpr1-2* mutant seedlings produced less than 10% of the lateral roots observed in wild type in high P conditions. In low P, the density of lateral roots was 25% of wild type (Fig. 3C). Interestingly, although *lpr1-2* produced fewer lateral roots compared to wild type in both low and high P conditions, exposure to low P conditions induced a 3- to 5-fold increase in



**Figure 3.** Effects of P availability on the root system architecture of wild-type (Ws) and *lpr1-2* plants. A, Primary root growth kinetic assay. Seedlings were grown for the indicated times over the surface of agar plates containing high or low P and primary root lengths were measured. B, Mean root hair length at 10 d in the above experiment. C, Mean lateral root density expressed as the number of lateral roots per centimeter, recorded at 15 d in the experiment. In B and C, black bars represent growth in high P and white bars represent growth in low P. All values shown represent the mean of 20 seedlings  $\pm$  SD. Letters represent means statistically different ( $P < 0.05$ ).

lateral root density in the mutant, similar to that observed for wild type. Most lateral roots that emerged in low P in the mutants elongated normally and were located close to the root tip (Fig. 1B). Similar results were obtained with *doc1* (data not shown). Taken together, these results show that the primary root and root hair responses to low P were unaltered in

*lpr1-2* mutants and that mutations in *BIG* may alter more specific developmental traits related to lateral root formation.


#### Effect of P Availability on LRP Initiation

To more closely analyze the effects of P availability on lateral root development, and to investigate whether *lpr1-1* is defective in LRP formation or in the subsequent elongation of these primordia, LRP originating in the primary root were counted at 4, 6, and 8 d after germination (see "Materials and Methods"). The developmental stage of each LRP was classified according to Zhang et al. (1999): stage A, up to 3 cell layers; stage B, unemerged, of more than 3 cell layers; stage C, emerged lateral roots of less than 0.5 mm in length; stage D, lateral root longer than 0.5 mm. In wild-type plants grown in high P conditions, the number of LRP at the four developmental stages increased with time. At 8 d, however, most LRP remained at an early developmental stage (Fig. 4A). Interestingly, low P conditions stimulated LRP transition from stage A to stage D. This effect was clearly observed at 4 d and increased at 8 d after germination. At day 8, most wild-type LRP were classified as stage D (Fig. 4B). In high P conditions, *lpr1-1* mutants showed a reduction in total LRP formation when compared to wild type; no LRPs at stages B, C, and D were observed at 4, 6, or 8 d (Fig. 4A). At 6 and 8 d of growth in low P conditions, we observed an increase in LRP transition from early-to-late stages of development in *lpr1-1* roots (Fig. 4B).

The LRP density was calculated by dividing the number of LRP by the length of the primary root to normalize for the effects of P availability on root length. LRP density significantly increased in plants grown at low P when compared with high P-grown plants. Although low P conditions increased the density of LRP in *lpr1-1* mutants when compared to high P conditions, *lpr1* always showed lower LRP density than wild type at the two P treatments (Fig. 4C). These results indicate that low P conditions modify the Arabidopsis root system architecture, most likely accelerating the emergence of LRP from the primary root and mutations in *BIG* interfere with the process of LRP initiation.

#### *BIG* Mutants Display Reduced Auxin Maximum in Primary and Lateral Roots

To evaluate the contribution of auxin responsiveness in the altered lateral root formation in response to P availability in *BIG* mutants, we crossed *lpr1-2* with a transgenic plant expressing the *DR5:uidA* construct, which is useful in studying auxin-regulated gene expression in Arabidopsis (Ulmasov et al., 1997). As previously reported (Sabatini et al., 1999), we observed  $\beta$ -glucuronidase (GUS) activity in the columella and quiescent center of primary roots of 10-d-old trans-



**Figure 4.** Effects of P availability on wild-type (Col-0) and *lpr1-1* lateral root development. A, Lateral root stage distribution in 4-, 6-, and 8-d-old primary roots grown on media containing high P. B, Lateral root stage distribution in 4-, 6-, and 8-d-old primary roots grown on media containing low P. C, LRP density at 4, 6, and 8 d after germination in


genic *DR5:uidA* seedlings grown in high P (Fig. 5A). In these conditions, GUS staining in primary root tips of *lpr1-2* seedlings was dramatically reduced (Fig. 5M). In low P conditions, both wild type and *lpr1-2* showed little or no GUS activity (Fig. 5, G and Q). This reduction of the *DR5:uidA* maximum in primary root tips was accompanied by differentiation processes in this region. For instance, long root hairs were formed very close to the root meristem (Fig. 5, G and Q). We further examined GUS expression in LRP at different developmental stages in wild-type and *lpr1-2* seedlings grown in high and low P conditions. For transgenic *DR5:uidA* seedlings, cells of LRP stained for GUS activity when plants were cultivated on high P medium (Fig. 5, B–F). Interestingly, LRP of wild-type plants grown in low P showed increased GUS activity at most developmental stages analyzed (Fig. 5, H–L). In contrast, GUS staining in LRP of *lpr1-2* seedlings grown in high P was undetectable (Fig. 5, N–P). When grown in low P conditions, GUS activity was detected in the LRP of *lpr1-2* seedlings, albeit at a lower level than in wild type (Fig. 5, R–T). Although the number of LRP at advanced developmental stages in *lpr1-2* was partially recovered by low P treatment, only a limited number progressed to emerge from the primary root (Fig. 4B). These primordia clearly showed reduced GUS expression when compared to wild-type primordia at similar developmental stages (Fig. 5, K and T, L and U). These data suggest that low P enhances auxin responsiveness in LRP.

#### Effect of Exogenous Auxin on Root Development in Wild Type and *BIG* Mutants

Exogenous auxin inhibits primary root elongation and stimulates lateral root formation (Casimiro et al., 2001). The observation that the expression of the auxin response marker *DR5:uidA* (Fig. 5G) was not increased in low P conditions suggested that primary root inhibition by low P treatment was not caused by auxin accumulation at the primary root tip region. Instead, it shows that P deprivation negatively affects the auxin response maximum normally present at the root tip under high P conditions. To investigate whether exogenously supplied auxins can restore the auxin maximum and in this way restore primary root growth in low P-grown plants, the effect of the synthetic auxin 2,4-dichlorophenoxyacetic acid (2,4-D) on the growth and development of wild-type and *lpr1-2* root systems was determined at low and high P conditions. Seedlings were grown for 5 d in media with low or high P content and then transferred to media containing the same P level supplemented with varying 2,4-D con-


high or low P conditions. Wild-type (Col-0) and *lpr1-1* seedlings were cleared and the number and stage of LRP recorded according to Zhang et al. (1999). This analysis was repeated twice with similar results.

**Figure 5.** Effect of P availability on auxin-regulated gene expression. Ten-hour GUS staining of *DR5:uidA* primary roots and LRP at various stages of development in wild-type seedlings grown for 10 d in medium with high (A–F) or low (G–L) P content. Ten-hour GUS staining of *DR5:uidA* primary roots and LRP at various stages of development in *lpr1-2* seedlings grown for 10 d in medium with high (M–P) or low (Q–U) P content. Photographs are representative individuals of at least 20 plants stained. Scale bar = 100  $\mu$ m.



centrations. The increase in primary root length was measured after 5 d. As shown in Figure 6A, primary roots of wild-type and *lpr1-2* plants transferred from medium containing low P or high P to similar medium supplemented with 2,4-D all decreased primary root growth with increasing auxin concentrations, although plants grown on high P were more sensitive to the inhibitory effects of 2,4-D on primary root elongation than those grown on low P. These results suggest that, although high concentrations of exogenously supplied auxins and low P conditions both inhibit primary root growth, they likely operate through independent signaling mechanisms. To fur-

ther determine whether exogenous auxin could stimulate lateral root formation in *lpr1-2* mutants, the number of lateral roots was determined 9 d after transfer to medium containing 2,4-D. Interestingly, low P-grown plants significantly increased their lateral root density in the presence of 10 nM 2,4-D when compared to high P-grown plants (Fig. 6B). When exposed to 25 or 50 nM 2,4-D, lateral root density of wild-type plants grown in high P was similar to that of plants grown in low P. Treatment of *lpr1-2* roots grown in low P conditions with 2,4-D gradually increased lateral root density in a dose-dependent manner (Fig. 6B). In contrast, *lpr1-2* seedlings grown in high P and



**Figure 6.** Effects of exogenous auxin on wild-type (Ws) and *lpr1-2* root development in low and high P. A, Mean primary root length of Ws and *lpr1-2* seedlings grown on 2,4-D. Seedlings were transferred at 5 d after germination from auxin-free media to media containing varying concentrations of 2,4-D. Primary root length was recorded 5 d after transfer. B, Mean lateral root density of Ws and *lpr1-2* seedlings grown on 2,4-D. Plants were transferred at 5 d after germination from auxin-free media to media containing varying concentrations of 2,4-D. Lateral root density was recorded 9 d after transfer ( $n = 15$ ).

transferred to 5 or 10 nM 2,4-D did not have a significant effect on lateral root density when compared to the untreated control. Fifty nanomolar 2,4-D allowed normal lateral root formation in the mutant in high P conditions (Fig. 6B). Since auxin rescued the *lpr1-2* lateral root phenotype both in high and low P, our results are consistent with the hypothesis that BIG has an important role in auxin transport. Moreover, as both wild-type and *lpr1-2* plants displayed an amplification of the effect of 2,4-D on lateral root density at low P, it is possible that P limitation increases auxin sensitivity in the Arabidopsis root.

#### Effects of Brefeldin A on Lateral Root Development in High and Low P Conditions


Treatments of Arabidopsis roots with inhibitors that reduce polar auxin transport, such as TIBA and NPA,

inhibit lateral root formation in high and low P conditions (López-Bucio et al., 2002). Recently, it was found that several auxin transport inhibitors are not specific, but rather generally affect the transport of membrane proteins (Geldner et al., 2001). Interestingly, the vesicle-trafficking inhibitor brefeldin A (BFA) was found to mimic most physiological effects of auxin transport inhibitors (Geldner et al., 2001). We therefore tested whether BFA-treated wild type could mimic the lateral root defects of *BIG* mutants by growing wild-type seedlings on vertical agar plates with high or low P content, supplemented with low concentrations of BFA. In high P-grown plants, treatment with 10  $\mu\text{M}$  BFA dramatically altered lateral root formation, whereas 20  $\mu\text{M}$  BFA inhibited primary root growth and altered root gravitropism (Fig. 7, A–E). Similar results were previously reported (Geldner et al., 2001). In P limitation, the most important effect of 20  $\mu\text{M}$  BFA treatment on root system architecture was the inhibition of lateral root formation (Fig. 7, B–F). Further LRP analysis in low P conditions showed that, similarly to mutations in *BIG*, BFA inhibition of lateral root development was mainly caused by a reduction in LRP formation (data not shown).

## DISCUSSION

### The Role of Auxin Transport in Root Architectural Response to Low P Conditions in Arabidopsis

When Arabidopsis plants are grown in limiting P conditions, primary root growth is inhibited and the



**Figure 7.** Effect of BFA on wild-type seedlings grown in high and low P availability. Seedlings were grown on agar plates containing different concentrations of BFA. Twelve-day-old wild-type seedlings grown in media containing high P (A) or low P (B) without BFA. High P-grown wild type treated with 10  $\mu\text{M}$  (C) or 20  $\mu\text{M}$  (E) BFA. Low P-grown wild type treated with 10  $\mu\text{M}$  (D) or 20  $\mu\text{M}$  (F) BFA. Photographs are representative individuals of at least 30 plants analyzed. Scale bar = 1 cm.

lateral root density is increased, resulting in the formation of a shallower and wider root system (Williamson et al., 2001; López-Bucio et al., 2002; Ticconi et al., 2004). Little is known about the physiological and molecular mechanisms by which plants alter their growth and development when exposed to limiting P supply. Auxins are a class of phytohormones required for myriad plant developmental processes. Many plant species respond to the exogenous application of auxins by inhibiting primary root growth and producing large numbers of lateral roots. A regulated, differential distribution of auxin underlies several adaptive processes, including organogenesis, meristem patterning, and postembryonic root development (Bhalerao et al., 2002; Friml, 2003). We previously provided pharmacological data pointing to an important role for auxins in the root architectural changes induced by P limitation (López-Bucio et al., 2002). From these studies, the auxin transport inhibitor TIBA was found to inhibit the formation of lateral roots in P-deprived Arabidopsis plants. In this article, we show that mutations in *BIG* impair lateral root induction in low P conditions and provide genetic support for the requirement of auxin transport in lateral root proliferation during low P stress. Since low P conditions similarly inhibited primary root growth and stimulated root hair growth in *lpr1* mutants and wild type, we conclude that an auxin transport independent pathway is involved in low P stress-induced root architectural alterations in Arabidopsis.

To date, several mutants in *BIG* have been isolated, including *doc1* (Li et al., 1994), *tir3* (Ruegger et al., 1997), and *umb1* (Kanyuka et al., 2003). Although these mutants were isolated in very different screens, complementation tests demonstrated that they are allelic (Gil et al., 2001; Kanyuka et al., 2003). Furthermore, *tir3-1* and *doc1-1* stem segments transport reduced amounts of radioactively labeled indole-3-acetic acid (IAA) compared with wild type. Therefore, it was proposed that *BIG* is required for polar auxin transport in Arabidopsis (Gil et al., 2001). As previously reported for *tir3* (Ruegger et al., 1997), *lpr1* mutants showed reduced primary root growth when compared to wild type grown in high P conditions. This result suggests that auxin transport is required to provide sufficient auxin to promote root growth. However, treatment with 2,4-D showed that excess auxin still results in root growth inhibition (Fig. 5A), suggesting that a precise balance between auxin transport and inactivation is required to sustain root growth.


We also showed that growth of the primary root is dramatically reduced in P-limiting conditions in wild-type and *lpr1* plants, thus suggesting that low P-induced primary root growth inhibition likely operates through a different mechanism than that regulating primary root growth in high P conditions. Therefore, *lpr1* mutants separate lateral root proliferation from primary root growth arrest during P limitation and reveal a key role for *BIG* in lateral root

initiation both in high and low P conditions. No other Arabidopsis mutants with defects in auxin transport/signaling have been found to show clear root developmental alterations to P stress (Williamson et al., 2001; López-Bucio et al., 2002). Since auxin fails to rescue the short-root phenotype of wild-type and *lpr1-2* plants in low P, and *DR5:uidA* expression in primary roots of *lpr1-2* plants is similarly reduced when compared to wild type in low P, we suggest that the inhibition of primary root growth and the stimulation of lateral root emergence triggered by low P operate through at least two independent mechanisms: one triggering primary root growth arrest that is independent of auxin, and one stimulating LRP emergence, a process that requires auxin transport-induced lateral root initiation and normal *BIG* function (Fig. 8).

### Potential Role of *BIG* in Regulating Acropetal Auxin Transport and Lateral Root Emergence in P-Limiting Conditions

Two distinct auxin transport streams have been described in roots, acropetal from the shoot system to the root and basipetal from the root tip to the root base (Rashotte et al., 2000; Bhalerao et al., 2002). As both of these are inhibited by phytohormones, it is difficult to pharmacologically delineate the roles of the two transport streams in regulating lateral root formation in different P supplies.

The removal of primary root tips or cell ablation of root cap cells by expression of a diphtheria toxin stimulate lateral root formation (Torrey, 1950; Tsugeki



**Figure 8.** Root architectural responses to P limitation and their regulation. Low P availability alters root architecture by at least two mechanisms, one that is independent of auxin transport and *BIG* function leads to primary root growth inhibition and root hair growth stimulation, and one where low P promotes root branching interacting with auxin. In low P, lateral root development is a two-step process involving LRP initiation from the pericycle and the further maturation of these LRP into fast-growing lateral roots. In our model, LRP initiation requires auxin transport and normal *BIG* function, whereas LRP maturation is a *BIG* independent process. High P availability represses low P-induced root developmental responses.



and Fedoroff, 1999). Therefore, we propose that inhibition of primary root growth by low P can be considered a physiological decapitation that triggers lateral root development. Since lateral root growth requires auxin, it is possible that elongating LRP during the low P root architectural response become potent new sinks for auxin, reducing the supply of auxin to older root meristems. This premise may explain the drastic reduction of *DR5:uidA* expression in primary root tips of low P-grown *Arabidopsis* plants and the further increase in GUS expression in their LRP (Fig. 5). Our detailed analyses of LRP (Fig. 4) and *DR5:uidA* expression in wild-type and *lpr1* root systems (Fig. 5) provide compelling evidence for the hypothesis that P deprivation accelerates the emergence of LRP and that *BIG* is required mainly for LRP initiation. Since lateral root formation and further lateral root growth that occurs in *BIG* mutants in low P takes place close to the primary root tip and root gravitropism is little or not affected in the mutants in high P conditions (Fig. 1, B and C), we speculate that basipetal transport of auxin is unaltered in *BIG* mutants and that *BIG* may play an essential role in acropetal transport of root auxin. Furthermore, since *doc1* and *tir3* mutants have a 70% reduction in the transport of auxin in inflorescence stems (Gil et al., 2001), we suggest that the *lpr1* alterations in root developmental responses to P availability likely result from reduced acropetal auxin transport.

We also found that roots of *lpr1* mutants may have suboptimal auxin levels or responsiveness, because 2,4-D application restored *lpr1* lateral root formation in high and low P conditions to wild-type levels. This result suggests that, although *BIG* may be an integrator of several hormonal and environmental signals (Kanyuka et al., 2003), the primary root developmental alterations reported for *big* mutants are likely dependent on auxin action.

#### Increased Lateral Root Emergence in Low P Conditions Involves a BFA-Sensitive Auxin Transport Pathway

*BIG* is thought to be required for the correct expression, localization, or stability of the NPA-binding protein, which may affect auxin transport by influencing trafficking of various proteins to the plasma membrane, including the auxin efflux carrier PIN1 (Gil et al., 2001; Muday et al., 2003). Monensin and BFA, inhibitors of Golgi-mediated protein secretion, rapidly perturb the transport activity of plasma membrane-associated efflux carriers for IAA and inhibit polar transport of IAA (Wilkinson and Morris, 1994; Morris and Robinson, 1998). In our experiments, low concentrations of BFA dramatically inhibited lateral root formation in both high and low P conditions. This result suggests that the low P pathway for lateral root induction requires normal cycling of auxin transport proteins and that *BIG* may participate in vesicle transport. Currently, it is unknown how *BIG* could regulate vesicle transport. Gil et al. (2001) reported that *BIG* encodes a protein extraordinary in size (560 kD)

that contains several zinc finger domains. *BIG* is similar to the *Drosophila calossin/pushover* (CalO/PUSH) gene, a member of a gene family also present in *Caenorhabditis elegans* and human genomes. In *Drosophila*, CalO affects synaptic transmission at the neuromuscular junctions, with specific defects in neurotransmitter release evoked by nerve stimulation (Richards et al., 1996). This process is dependent on the synaptic vesicle cycle that involves a cascade of protein-protein interactions (Sudhof, 1995). ADP ribosylation factors-GTP exchange factors are thought to determine the destination of membrane-trafficking vesicles by specific recruitment of protein coats for vesicles that determine their cellular location (Muday et al., 2003). ADP ribosylation factors-GTP exchange factors are also a target of BFA, which blocks vesicle delivery of proteins to the membrane surface. Whatever the target of *BIG*, the signaling cascade by which it controls lateral root development deserves further attention. In this regard, Grebe et al. (2002), while investigating the components of the auxin machinery that regulates root hair polarity, found that the presumptive auxin influx carrier AUX1 contributes to apical-basal cell polarity. The authors showed that AUX1 function is required for polarity changes induced by exogenous application of auxin. Interestingly, similar to *aux1* mutants, treatment with BFA interferes with root hair initiation by inhibiting membrane trafficking of *aux1*. Although our observation that wild-type plants treated with BFA resemble *lpr1* mutants, it is unlikely that the reported lateral root alterations involve AUX1 function. Both *aux1-7* and *eir1* auxin transport mutants respond normally to low P conditions producing large numbers of lateral roots (López-Bucio et al., 2002).

Our data provide evidence that increased lateral root development in low P conditions involves a BFA-sensitive auxin transport pathway. However, we cannot exclude that local auxin synthesis in LRP or increased sensitivity to auxins in LRP of low P-grown plants also accounts for earlier maturation of lateral roots. Further research is needed to provide unequivocal insight into how P limitation regulates the different traits that comprise the final configuration of the *Arabidopsis* root system and which of these developmental alterations involve auxin action.

## MATERIALS AND METHODS

### Plant Material and Growth Conditions

*Arabidopsis* (*Arabidopsis thaliana*) ecotypes Col-0 and Ws were used for all experiments. Seeds were surface sterilized with 95% (v/v) ethanol for 5 min and 20% (v/v) bleach for 7 min. After 5 washes in distilled water, seeds were germinated and grown on agar plates containing 0.1 × Murashige and Skoog medium, pH 5.7, 0.5% (w/v) Suc, and 1% (w/v) agar (López-Bucio et al., 2002). The basic medium contained 2.0 mM NH<sub>4</sub>NO<sub>3</sub>, 1.9 mM KNO<sub>3</sub>, 0.3 mM CaCl<sub>2</sub> 2H<sub>2</sub>O, 0.15 mM MgSO<sub>4</sub> 7H<sub>2</sub>O, 5 μM KI, 25 μM H<sub>3</sub>BO<sub>3</sub>, 0.1 mM MnSO<sub>4</sub> H<sub>2</sub>O, 0.3 mM ZnSO<sub>4</sub> 7H<sub>2</sub>O, 1 μM Na<sub>2</sub>MoO<sub>4</sub> 2H<sub>2</sub>O, 0.1 μM CuSO<sub>4</sub> 5H<sub>2</sub>O, 0.1 μM CoCl<sub>2</sub> 6H<sub>2</sub>O, 0.1 mM FeSO<sub>4</sub> 7H<sub>2</sub>O, 0.1 mM Na<sub>2</sub>EDTA 2H<sub>2</sub>O, inositol (10 mg L<sup>-1</sup>), and Gly (0.2 mg L<sup>-1</sup>).

Plates were placed at an angle of 65° to allow root growth along the agar surface and to allow unimpeded hypocotyl growth into the air. Plants were grown at 22°C to 24°C in a plant growth cabinet (Percival Scientific, Perry, IA), with a photoperiod of 16 h of light, 8 h of darkness, and a light intensity of 300  $\mu\text{mol m}^{-2} \text{s}^{-1}$ . Seeds of transgenic *DR5::uidA* Arabidopsis plants (Ulmasov et al., 1997) were provided by Dr. Tom Guilfoyle, and homozygous *F<sub>3</sub>* progeny of crosses with *lpr1-2* were used for analysis of GUS expression.

To test the effects of 2,4-D and BFA, low (1  $\mu\text{M}$   $\text{NaH}_2\text{PO}_4$ ) and high (1 mM  $\text{NaH}_2\text{PO}_4$ ) P nutrient medium was supplemented with ethanol-dissolved compounds. The compounds were added to cooled (50°C) molten medium and poured into plates. Chemicals were purchased from Sigma (St. Louis).

## Mutant Isolation Procedure

EMS-mutagenized seeds (Col-0) were purchased from Lehle Seeds (Round Rock, TX). T-DNA lines (Ws; Krysan et al., 1999) were provided by the Ohio Arabidopsis Seed Stock Center. Seeds were surface sterilized and plated on low P (1  $\mu\text{M}$   $\text{NaH}_2\text{PO}_4$ ) 0.1  $\times$  Murashige and Skoog medium. A total of approximately 25,000 *M<sub>2</sub>* seedlings descended from EMS-mutagenized seed and 38,000 T-DNA lines were screened for reduced lateral root formation by placing seeds on 100-cm<sup>2</sup> nutrient agar plates (20 seeds/plate). The seeds were distributed in 2 rows on the agar surface at a density of 1 seed/centimeter, stratified at 4°C for 48 h, and then incubated at 22°C. Fourteen days after germination, low P-grown plants have a short primary root and a large number of lateral roots formed close to the root apex. Putative mutants with short primary roots and reduced number of lateral roots were selected, transferred to soil, and allowed to self-fertilize. Homozygous *M<sub>3</sub>* seeds were rescreened for altered lateral root formation in low P and back-crossed four times to wild type (either Col-0 or Ws) to remove unlinked mutations.

## Genetic Analysis of *lpr1* Mutants

The *lpr1-1* allele was isolated in a Col-0 background from an EMS mutant population. The *lpr1-2* allele was isolated in a Ws background from the Sussman and Amasino T-DNA mutant collection. The *lpr1* mutants were backcrossed 4 times with the corresponding wild-type parent. To determine the segregation pattern of the *lpr1* phenotype, 500 *F<sub>2</sub>* seedlings derived from each cross between *lpr1-1*  $\times$  Col-0 and *lpr1-2*  $\times$  Ws were analyzed in solid low P medium. A typical 3:1 recessive segregation was observed for the wild-type/*lpr1* phenotype. For allelism tests, homozygous *lpr1-1*, *lpr1-2*, and *doc1* were crossed to each other and plants from the *F<sub>1</sub>* population were analyzed for the *lpr1* phenotype. Chromosomal localization of mutations corresponding to *lpr1-1* was determined using a combination of cleaved amplified polymorphic sequences (Konieczny and Ausubel, 1993) and simple sequence length polymorphism (Bell and Ecker, 1994) molecular markers. Homozygous *lpr1-1* plants in the Col-0 background were out-crossed to Ws. Genomic DNA for PCR was prepared from *F<sub>2</sub>* plants with *lpr1* mutant phenotypes (Celenza et al., 1995).

## Histochemical Analysis

For histochemical analysis of GUS activity, Arabidopsis seedlings were incubated overnight at 37°C in a GUS reaction buffer (0.5 mg/mL of 5-bromo-4-chloro-3-indolyl- $\beta$ -D-glucuronide in 100 mM sodium phosphate, pH 7), and the stained seedlings were cleared (Malamy and Benfey, 1997). For each marker line and for each treatment, at least 10 transgenic plants were analyzed. A representative plant was chosen for each P or auxin treatment and photographed using the Nomarski optics on a Leica (Wetzlar, Germany) DMR microscope.

## Data Analysis

Arabidopsis root systems were viewed with an AFX-II-A stereomicroscope (Nikon, Tokyo). All lateral roots emerged from the primary one and observed at the 3 $\times$  objective were included in the lateral root number data. Primary root length was determined for each root using a ruler. The stages of LRP were analyzed on cleared roots using a Nikon UW microscope in the 40 $\times$  objective.

For all experiments, the overall data were statistically analyzed in the SPSS 10 program (SPSS, Chicago). Univariate and multivariate analyses with a Tukey's post hoc test were used for testing differences in primary root length, lateral root number, and lateral root density in P treatments in wild-type and mutant plants. Different letters are used to indicate means that differ significantly ( $P < 0.05$ ).

## ACKNOWLEDGMENTS

We are thankful to Drs. Joanne Chory and Tom Guilfoyle for kindly providing us with seeds of mutant and transgenic lines. J.L.-B. gratefully acknowledges the Mexican Academy of Sciences and the Mexico-USA Foundation for Science for financial support to visit the Department of Biochemistry and Cell Biology at Rice University, Houston. We thank Dr. Bethany K. Zolman for her support and technical improvement for *lpr1-1* mapping.

Received July 9, 2004; returned for revision August 18, 2004; accepted November 23, 2004.

## LITERATURE CITED

- Abel S, Ticconi CA, Delatorre CA (2002) Phosphate sensing in higher plants. *Physiol Plant* **115**: 1–8
- Al-Ghazi Y, Muller B, Pinloche S, Tranbarger TJ, Nacry P, Rossignol M, Tardieu F, Doumas P (2003) Temporal responses of Arabidopsis root architecture to phosphate starvation: evidence for the involvement of auxin signaling. *Plant Cell Environ* **26**: 1053–1066
- Bell CJ, Ecker JR (1994) Assignment of 30 microsatellite loci to the linkage map of Arabidopsis. *Genomics* **19**: 137–144
- Bhalerao RP, Eklof J, Ljung K, Marchant A, Bennet M, Sandberg G (2002) Shoot-derived auxin is essential for early lateral root emergence in Arabidopsis seedlings. *Plant J* **29**: 325–332
- Casimiro I, Merchant A, Bhalerao RP, Beekman T, Dhoog S, Swarup R, Graham N, Inze D, Sandberg G, Casero PJ, et al (2001) Auxin transport promotes lateral root initiation. *Plant Cell* **13**: 843–852
- Celenza JL, Grisafi PL, Fink GR (1995) A pathway for lateral root formation in *Arabidopsis thaliana*. *Genes Dev* **9**: 2131–2142
- Friml J (2003) Auxin transport: shaping the plant. *Curr Opin Plant Biol* **6**: 7–12
- Geldner N, Friml J, Dieter Stierhof Y, Jurgens G, Palme K (2001) Auxin transport inhibitors block PIN1 cycling and vesicle trafficking. *Nature* **413**: 425–428
- Gil P, Dewey E, Friml J, Zhao Y, Snowdon KC, Putterill J, Palme K, Estelle M, Chory J (2001) BIG: a calossin-like protein required for polar auxin transport in *Arabidopsis*. *Genes Dev* **15**: 1985–1997
- Gilbert GA, Knight JD, Vance CP, Allan DL (2000) Proteoid root development of phosphorus deficient lupin is mimicked by auxin and phosphonate. *Ann Bot (Lond)* **85**: 921–928
- Grebe M, Friml J, Swarup R, Ljung K, Sandberg G, Terlou M, Palme K, Bennet M, Scheres B (2002) Cell polarity signaling in Arabidopsis involves a BFA-sensitive auxin influx pathway. *Curr Biol* **12**: 329–334
- Johnson JE, Vance CP, Allan DL (1996) Phosphorus deficiency in *Lupinus albus*. Altered lateral root development and enhanced expression of phosphoenolpyruvate carboxylase. *Plant Physiol* **112**: 31–41
- Kanyuka K, Praekelt U, Franklin KA, Billingham OE, Hooley R, Whitlam GC, Halliday KJ (2003) Mutations in the huge *Arabidopsis* gene *BIG* affect a range of hormonal and light responses. *Plant J* **35**: 57–70
- Konieczny A, Ausubel FM (1993) A procedure for mapping Arabidopsis mutations using co-dominant ecotype-specific PCR-based markers. *Plant J* **4**: 403–410
- Krysan PJ, Young JC, Sussman MR (1999) T-DNA as an insertional mutagen in Arabidopsis. *Plant Cell* **12**: 2283–2290
- Li HM, Altschmid L, Chory J (1994) Arabidopsis mutants define downstream branches in the phototransduction pathway. *Genes Dev* **8**: 339–349
- López-Bucio J, Cruz-Ramírez A, Herrera-Estrella L (2003) The role of nutrient availability in regulating root architecture. *Curr Opin Plant Biol* **6**: 280–287
- López-Bucio J, Hernández-Abreu E, Sánchez-Calderón L, Nieto-Jacobo ME, Simpson J, Herrera-Estrella L (2002) Phosphate availability alters architecture and causes changes in hormone sensitivity in the Arabidopsis root system. *Plant Physiol* **129**: 244–256
- López-Bucio J, Nieto-Jacobo ME, Ramírez-Rodríguez V, Herrera-Estrella L (2000) Organic acid metabolism in plants: from adaptive physiology to transgenic varieties for cultivation in extreme soils. *Plant Sci* **160**: 1–13
- Malamy JE, Benfey PN (1997) Organization and cell differentiation in lateral roots of *Arabidopsis thaliana*. *Development* **124**: 33–44

- Morris DA, Robinson JS** (1998) Targeting of auxin carriers to the plasma membrane: differential effects of brefeldin A on the traffic of auxin uptake and efflux carriers. *Planta* **205**: 606–612
- Muday GK, Peer WA, Murphy AS** (2003) Vesicular cycling mechanisms that control auxin transport polarity. *Trends Plant Sci* **8**: 301–304
- Neumann G, Martinoia E** (2002) Cluster roots: an underground adaptation for survival in extreme environments. *Trends Plant Sci* **7**: 162–167
- Raghothama KG** (1999) Phosphate acquisition. *Annu Rev Plant Physiol Plant Mol Biol* **50**: 665–693
- Rashotte AM, Brady SR, Reed RC, Ante SJ, Muday G** (2000) Basipetal auxin transport is required for gravitropism in roots of *Arabidopsis*. *Plant Physiol* **122**: 481–490
- Richards S, Hillman T, Stern M** (1996) Mutations in the *Drosophila pushover* gene confer increased neuronal excitability and spontaneous synaptic vesicle fusion. *Genetics* **142**: 1215–1223
- Ruegger M, Dewey E, Hobbie L, Brown D, Bernasconi P, Turner J, Muday G, Estelle M** (1997) Reduced naphthylphthalamic acid binding in the *tir3* mutant of *Arabidopsis* is associated with a reduction in polar auxin transport and diverse morphological defects. *Plant Cell* **9**: 745–757
- Sabatini S, Beis D, Wolkenfelt H, Murfett J, Guilfoyle T, Malamy J, Benfey P, Leyser O, Bechtold N, Weisbeek P, et al** (1999) An auxin-dependent distal organizer of pattern and polarity in the *Arabidopsis* root. *Cell* **99**: 463–472
- Skene KR, James WM** (2000) A comparison of the effects of auxin on cluster root initiation and development in *Grevillea robusta* Cunn. Ex R. Br. (Proteaceae) and the genus *Lupinus* (Leguminosae). *Plant Soil* **219**: 221–229
- Sudhof TC** (1995) The synaptic vesicle cycle: a cascade of protein-protein interactions. *Nature* **375**: 645–673
- Ticconi CA, Delatorre CA, Lahner B, Salt DE, Abel S** (2004) *Arabidopsis pdr2* reveals a phosphate-sensitive checkpoint in root development. *Plant J* **37**: 801–814
- Torrey JG** (1950) Induction of lateral roots by indoleacetic acid and decapitation. *Am J Bot* **37**: 257–264
- Tsugeki R, Fedoroff NV** (1999) Genetic ablation of root cap cells in *Arabidopsis*. *Proc Natl Acad Sci USA* **96**: 12941–12946
- Ulmasov T, Murfett J, Hagen G, Guilfoyle T** (1997) Aux/IAA proteins repress expression of reporter genes containing natural and highly active synthetic auxin response elements. *Plant Cell* **9**: 1963–1971
- Vance CP, Uhde-Stone C, Allan DL** (2003) Phosphorus acquisition and use: critical adaptations by plants for securing a nonrenewable source. *New Phytol* **157**: 423–447
- Wilkinson S, Morris DA** (1994) Targeting of auxin carriers to the plasma membrane: effects of monensin on transmembrane auxin transport in *Cucurbita pepo* L. tissue. *Planta* **193**: 194–202
- Williamson LC, Ribrioux S, Fitter AH, Leyser O** (2001) Phosphate availability regulates root system architecture in *Arabidopsis*. *Plant Physiol* **126**: 875–882
- Zhang H, Jennings A, Barlow PW, Forde BG** (1999) Dual pathways for regulation of root branching by nitrate. *Proc Natl Acad Sci USA* **96**: 6529–6534