

HKT1;5-Like Cation Transporters Linked to Na⁺ Exclusion Loci in Wheat, *Nax2* and *Kna1*^{1[OA]}

Caitlin S. Byrt*, J. Damien Platten, Wolfgang Spielmeier, Richard A. James, Evans S. Lagudah, Elizabeth S. Dennis, Mark Tester, and Rana Munns

Commonwealth Scientific and Industrial Research Organization Plant Industry, Canberra, Australian Capital Territory 2601, Australia (C.S.B., J.D.P., W.S., R.A.J., E.S.L., E.S.D., R.M.); and Australian Centre for Plant Functional Genomics, Glen Osmond, South Australia 5064, Australia (C.S.B., M.T.)

Bread wheat (*Triticum aestivum*) has a greater ability to exclude Na⁺ from its leaves and is more salt tolerant than durum wheat (*Triticum turgidum* L. subsp. *durum* [Desf.]). A novel durum wheat, Line 149, was found to contain a major gene for Na⁺ exclusion, *Nax2*, which removes Na⁺ from the xylem in the roots and leads to a high K⁺-to-Na⁺ ratio in the leaves. *Nax2* was mapped to the distal region on chromosome 5AL based on linkage to microsatellite markers. The *Nax2* locus on 5AL coincides with the locus for a putative Na⁺ transporter, *HKT1;5* (*HKT8*). The *Nax2* region on 5AL is homoeologous to the region on chromosome 4DL containing the major Na⁺ exclusion locus in bread wheat, *Kna1*. A gene member of the *HKT1;5* family colocalizes to the deletion bin containing *Kna1* on chromosome 4DL. This work provides evidence that *Nax2* and *Kna1* are strongly associated with *HKT1;5* genes.

Increase in salt tolerance of crops is needed to sustain agriculture in regions affected by natural or secondary salinity. Durum wheat (*Triticum turgidum* L. subsp. *durum* [Desf.]) is particularly sensitive to salinity and has a limited ability to exclude sodium, which is an important mechanism of salt tolerance in wheat (Tester and Davenport, 2003; Munns et al., 2006). Exclusion of Na⁺ from the leaves is due to low net Na⁺ uptake by cells in the root cortex and the tight control of net loading of the xylem by parenchyma cells in the stele (Tester and Davenport, 2003). To improve Na⁺ exclusion in durum wheat it is necessary to understand the molecular basis of these Na⁺ transport processes.

Modern durum cultivars do not exclude Na⁺ to the same extent as bread wheat (*Triticum aestivum*); however, a source of sodium exclusion in a novel durum wheat, Line 149, was described by Munns et al. (2000). Line 149 is derived from a cross between a *Triticum monococcum* (accession C68-101; AA) and a durum

cultivar Marrocos (AABB; The, 1973). The *T. monococcum* is the donor of the sodium exclusion trait (James et al., 2006). The sodium exclusion in Line 149 is conferred by two major genes (Munns et al., 2003) named *Nax1* and *Nax2* (for Na⁺ exclusion), inherited from the *T. monococcum*. Line 149 was crossed with the durum cultivar Tamaroi and selected F₂ lines were backcrossed into Tamaroi so that *Nax1* and *Nax2* were separated into two single gene BC₅F₂ families (James et al., 2006). *Nax1* was mapped on chromosome 2AL (Lindsay et al., 2004). *Nax1* is a putative Na⁺ transporter, a member of the HKT (high-affinity K⁺ transporter) family, *T. monococcum* HKT7-A2 (Huang et al., 2006). The *Nax1* gene confers a reduced rate of transport of Na⁺ from root to shoot and retention of Na⁺ in the leaf sheath, thus giving a higher sheath-to-blade Na⁺ concentration ratio (James et al., 2006). The second gene, *Nax2*, also confers a lower rate of transport of Na⁺ from root to shoot and has a higher rate of K⁺ transport, resulting in enhanced K⁺ versus Na⁺ discrimination in the leaf (James et al., 2006).

HKT genes control Na⁺ transport in higher plants, as demonstrated in rice (*Oryza sativa*), barley (*Hordeum vulgare*), and Arabidopsis (*Arabidopsis thaliana*; Mäser et al., 2002a; Berthomieu et al., 2003; Rus et al., 2004; Haro et al., 2005; Ren et al., 2005; Sunarpi et al., 2005). HKT genes have been given a new nomenclature and separated into two groups based on amino acid sequence (Platten et al., 2006). A Gly/Ser residue in the first pore loop of the protein differs between group 1 and group 2 HKT genes.

Group 1 HKT genes have a Ser in the first pore loop; this may make them more selective for Na⁺ (Horie et al., 2001; Mäser et al., 2002b; Garciadeblas et al., 2003; Platten et al., 2006). Group 1 HKT genes, such as *AtHKT1;1* (previous name *AtHKT1*) and *OsHKT1;5*

¹ This work was supported by the Commonwealth Scientific and Industrial Research Organization, Australian Centre for Plant Functional Genomics, and University of Adelaide (scholarship to C.S.B.); and by the New South Wales Agricultural Genomics Centre (J.D.P., E.S.D.); Australian Research Council Federation Fellowship (M.T.); and Grains Research and Development Corporation (R.A.J., M.T., R.M.).

* Corresponding author; e-mail caitlin.byrt@csiro.au; fax 61-2-6246-5399.

The author responsible for distribution of materials integral to the findings presented in this article in accordance with the policy described in the Instructions for Authors (www.plantphysiol.org) is: Caitlin S. Byrt (caitlin.byrt@csiro.au).

[OA] Open Access articles can be viewed online without a subscription.

www.plantphysiol.org/cgi/doi/10.1104/pp.106.093476

(previous name *OsHKT8*), transport Na⁺ only and may be involved in unloading of Na⁺ from the xylem (Uozumi et al., 2000; Garcíadeblas et al., 2003; Ren et al., 2005; Sunarpi et al., 2005). In wheat, group 1 HKT genes are involved in Na⁺ transport and may confer a phenotype of low leaf Na⁺ concentration in leaves (Huang et al., 2006).

For the group 2 genes there is no consensus on the mechanism of action or whether the main function is to transport Na⁺ or K⁺ (Schachtman and Schroeder, 1994; Rubio et al., 1995; Walker et al., 1996; Wang et al., 1998; Horie et al., 2001; Gollidack et al., 2002; Haro et al., 2005). Heterologous expression of *TaHKT2;1* (previous name *TaHKT1*) in yeast (*Saccharomyces cerevisiae*) and *Xenopus laevis* oocytes indicated that this gene is likely to play a role in Na⁺ and K⁺ transport (Rubio et al., 1995; Mäser et al., 2002b). Rodríguez-Navarro and Rubio (2006) provide an explanation for the conflicting results for group 2 HKT genes. They suggest that in plants some HKT mRNA transcripts have alternative initiations of translation and in heterologous systems translation may not occur exactly as in plants, leading to expression of proteins with different kinetic properties. A physiologically relevant function for group 2 transporters might be to transport Na⁺ when there is a limited supply of K⁺ so that the plant may have a monovalent cation for use in osmotic adjustment in the vacuole (Rodríguez-Navarro and Rubio, 2006).

The Na⁺ transporter *OsHKT1;5* localizes to the plasma membrane and is expressed in the xylem tissues (Ren et al., 2005). *OsHKT1;5* is likely to function in transporting Na⁺ out of the xylem into the xylem parenchyma cells (Ren et al., 2005), where it may then be effluxed through the cortex to the epidermis and back into the soil. Evidence is increasing that a major mode of action of *AtHKT1;1* is to increase Na⁺ retrieval from the xylem in mature roots (Sunarpi et al., 2005). *Nax2* functions in a similar way, in transporting Na⁺ out of the xylem. A compartmental loading experiment where ²²Na⁺ was fed only to the lower part of the roots showed that lines with *Nax2* withdrew more of the total ²²Na⁺ into the upper part of the root than lines without *Nax2* (James et al., 2006). The rates of root Na⁺ uptake were identical in the lines with and without *Nax2*, indicating that differences in shoot uptake were due to the net rate of xylem loading in the root (James et al., 2006).

The major Na⁺ exclusion locus in bread wheat, *Kna1*, is located on the D genome, on the distal part of chromosome 4. The phenotype is Na⁺ exclusion from the leaves and discrimination of K⁺ over Na⁺ in leaves, but no difference in Na⁺ concentrations in roots (Gorham et al., 1990). As there is no difference in Na⁺ concentrations in the roots it is likely that *Kna1* controls net xylem loading rather than net Na⁺ uptake (Gorham et al., 1990). Hence, an HKT-like gene is a good candidate for *Kna1*, as well as for *Nax2*.

Kna1 may be homoeologous to *Nax2*, the term homoeologous referring to a gene that used to be ho-

mologous in ancestral wheats before polyploidization of wheats and their related species. If so, one would expect that *Nax2* would be located in the group 4 chromosomes. During the evolution of wheat the distal part of chromosome 4A that is homoeologous to the distal part of chromosome 4D was translocated with chromosome 5A (Nelson et al., 1995). Therefore, if *Kna1* is located on the distal part of chromosome 4D and *Nax2* is homoeologous to *Kna1*, then *Nax2* would be physically located on the distal end of chromosome 5AL.

In the 19 years since the *Kna1* locus was first described (Gorham et al., 1987), *Kna1* has been transferred from bread wheat to durum wheat (from the D genome to the B genome) by homoeologous recombination (Dvořák and Gorham, 1992). *Kna1* has been mapped to the distal portion of the long arm of chromosome 4D and five markers have been identified that are linked to *Kna1*, within 2.2 cM (Dubcovsky et al., 1996; Luo et al., 1996), yet despite the extensive mapping work toward isolating *Kna1* no gene has been identified as a candidate for *Kna1*.

In this study we test whether the two Na⁺ exclusion genes, *Nax2* and *Kna1*, may be homoeologous, and we suggest that a putative Na⁺ transporter gene in wheat, *HKT1;5*, may correspond to both *Nax2* and *Kna1*.

RESULTS

Nax2 Is a Single Major Gene

Line 149 is from a cross between *T. monococcum* (C68-101; AA) and Marrocos (AABB; The, 1973; James et al., 2006). A single gene family segregating for *Nax2* was developed from Line 149, which contains the genes for Na⁺ exclusion *Nax1* and *Nax2*, via backcrossing into the cultivar Tamaroi that lacks the two genes (James et al., 2006). *Nax2* is responsible for a greater than 2-fold difference in leaf Na⁺ concentration. The mean leaf Na⁺ concentration of the BC₄F₂ *Nax2* single gene parent was 473 ± 72 μmol g dry weight⁻¹ compared to Tamaroi, which had a mean leaf Na⁺ concentration of 1,193 ± 48 μmol g dry weight⁻¹. The frequency distribution of leaf Na⁺ concentration from 137 BC₅F₂ plants indicated that *Nax2* was dominant, segregating in a 3:1 (low:high leaf blade Na⁺) ratio (Fig. 1A).

In this study, the 137 BC₅F₂ lines were progeny tested and the segregation of the Na⁺ exclusion trait was confirmed in the F_{2,3} families (773 individuals). The F_{2,3} families fitted the expected ratio for a single major gene (expected 94:31; observed 96:29; χ² = 0.171, P ≥ 0.05; Fig. 1B). The mean leaf Na⁺ of the *Nax2* single gene parent was 462 ± 23 μmol g dry weight⁻¹ in the F_{2,3} generation, compared to 473 ± 72 μmol g dry weight⁻¹ in the F₂ generation.

HKT1;5 Cosegregates with *Nax2*

Publicly available cation transporter sequences from rice were used to screen the GenBank database of




Figure 1. A, Frequency distribution for leaf 3 Na^+ concentrations of the BC_5F_2 lines in the single gene *Nax2* mapping population. Arrows indicate parental means ($n = 6$); Line 149: 278 ± 37 , *Nax2* single gene parent: 473 ± 72 , Tamaroi: $1,193 \pm 48$. Adapted from James et al. (2006). B, Relationship between the F_2 and $\text{F}_{2,3}$ progeny means for Na^+ concentration of leaf 3. Plants were grown at 150 mM NaCl for 10 d. \blacktriangle , Homozygous lacking *Nax2*; \circ , heterozygous for *Nax2*; \blacklozenge , homozygous for *Nax2*.

wheat expressed sequence tags (ESTs) to identify putative cation transporters in wheat. As part of this work, the protein sequence of *OsHKT1;5* was used to search the wheat EST database. The search identified a single closely related partial wheat EST sequence (CK193616). This partial sequence (*TaHKT1;5*) shared 86% identity at the nucleotide level and contained parts of the corresponding sequences of exon 2 and exon 3 of *OsHKT1;5* (Ren et al., 2005).

A probe designed from the partial *HKT1;5* sequence (named *HKT1;5* probe) identified RFLP between parental lines. Polymorphism found between the parents, Line 149 and Tamaroi, was due to the presence of an additional fragment in Line 149 when compared to Tamaroi (Fig. 2). The additional fragment in Line 149 cosegregated with the low Na^+ accumulation phenotype in all of the 137 BC_5F_2 families tested. All of the lines in the mapping family verified as homozygous and heterozygous for *Nax2* had the additional fragment and those lines that were homozygous for the

high leaf Na^+ phenotype lacked the fragment (a selection is shown in Fig. 2).

Wheat *HKT1;5* Gene Homoeologs

The *HKT1;5* probe hybridized to a single restriction fragment in DNA from the diploid *T. monococcum* (C68-101), the putative donor of salt tolerance in Line 149. A fragment of the same size as the *T. monococcum* fragment was present in Line 149 but absent in Tamaroi. Three other fragments were present in both Line 149 and Tamaroi (Fig. 2). The *monococcum* fragment in Line 149 cosegregated with *Nax2* (Fig. 2). Six restriction enzymes were tested and the *HKT1;5* probe always hybridized to at least two fragments in Tamaroi and three fragments in Line 149 (data not shown). *Hind*III produced an additional fragment in both parents (Fig. 2). In summary, Tamaroi contained at least two *HKT1;5*-like genes, while Line 149 had the same two fragments plus one additional gene member that was inherited from *T. monococcum*. These results were confirmed with three additional probes (named *HKT1;5* probe 2, *HKT1;5* probe 3, and *HKT1;5* probe 4) designed on different parts of the open reading frame (ORF) spanning from exon 1 to the 3' untranslated region.

To identify the *HKT1;5* gene family on the A, B, and D genomes of wheat, we analyzed DNA from the durum cultivar Langdon (AABB) carrying individual chromosome substitutions from the hexaploid Chinese




Figure 2. Cosegregation of the A genome *HKT1;5* fragment with *Nax2*. Southern-blot hybridization with the *HKT1;5* probe after *Hind*III digestion of DNA from *T. monococcum* (Mon), Line 149 (L149), Tamaroi (Tam), and plants from the *Nax2* BC_5F_3 mapping population with low leaf 3 Na^+ concentrations and high leaf 3 Na^+ concentrations. The genome origin of each band is shown on the right. In the *Nax2* mapping family the A genome fragment (inherited from the monococcum) cosegregates with *Nax2* in all of the 137 families.




Figure 3. Chromosomal location of the *HKT1;5* fragments. Southern-blot hybridization with the *HKT1;5* probe after *Hind*III restriction digest. A, DNA from *T. monococcum* (Mon), Tamaroi (Tam), Line 149 (L149), CS, Langdon (Lang), and Langdon substitution lines 4D(4A) and 4D(4B). The genome locations of the *HKT1;5* gene members and the approximate sizes (kilo bp) of the fragments are shown on the right. B, DNA from CS; CS chromosome arm deletion lines for chromosome 4B: N4AT4B (1), m4BT4A (2), N4DT4B (3); and CS ditelomic lines Dt4BS (4), Dt4DS (5), and Dt4DL (6); n = nullitetrasonic (no copies); T = tetrasomic (four copies); m = monosomic (one copy); Dt = ditelomic (for Dt4BS the long arm of 4B is missing and the short arm of 4B is present). The genome locations of the *HKT1;5* gene members are shown on the right.

Spring (CS; AABBDD; Joppa 1987). The *HKT1;5* probe hybridized to at least two fragments in Langdon DNA, producing an identical pattern to Tamaroi. Hybridization of the partial *HKT1;5* probe to CS DNA produced a pattern identical to that of Langdon and Tamaroi except for one additional chromosome 4D fragment. A *Hind*III digest of Langdon with chromosome 4A substituted for chromosome 4D of CS retained all three fragments and gained an additional fragment from chromosome 4D of CS. The Langdon substitution line

with chromosome 4B replaced by chromosome 4D of CS had lost all three Langdon fragments but retained the 4D fragment from CS (Fig. 3A). These results indicate that all *HKT1;5* hybridizing fragments in Langdon and Tamaroi are located on chromosome 4B and that Line 149 inherited an additional A genome member from *T. monococcum*.

Analysis of nullitetrasonic CS lines of homoeologous group 4 (refer to "Materials and Methods") confirmed that *HKT1;5* fragments were located on either chromosome 4B or 4D, but not 4A (Fig. 3B). DNA hybridization of the partial *HKT1;5* probe to ditelomic lines of CS, where individual group 4B or 4D chromosome arms have been deleted, positioned these genes on the long arm of chromosome 4B (at least three members), and one member on the long arm of chromosome 4D, where the major Na⁺ exclusion locus in bread wheat, *Kna1*, is located (Fig. 3B).

To study the relationship of *Kna1* to the bread wheat *HKT1;5-D* gene member, we probed DNA from CS and a series of telomeric deletion lines generated from chromosome 4DL in CS (Endo and Gill, 1996). In the telomeric deletion lines the 4DL chromosome has been shortened to different lengths. The *HKT1;5* gene member on chromosome 4DL was absent in the terminal deletion line flow length (Fl) 0.86 with approximately 14% of the chromosome arm deleted (Fig. 4A). The 4DL fragment was also missing in other lines with progressively larger terminal deletions (Fig. 4A). The loss of the chromosomal fragment containing the *TaHKT1;5* gene member on the D genome (*TaHKT1;5-D*) corresponded to an increase in average Na⁺ concentrations in the leaf blade of the deletion lines compared with the euploid salt-tolerant CS, and a decrease in the K⁺-to-Na⁺ ratio from 7.5 to 1.2 (Fig. 4B).

Genetic Map Location of *Nax2*

To position *Nax2* in the durum wheat genome, 34 microsatellite markers previously mapped to chromosome 4A were screened for polymorphism between parental lines and tested for linkage to *Nax2* in the segregating family. None of the markers from chromosome 4A were linked to *Nax2* (C.S. Byrt, unpublished data).

In an ancestor of modern wheats, chromosomes 4AL and 5AL exchanged short terminal segments (Liu et al., 1992). Considering this, we tested microsatellite markers that mapped to the distal segment of chromosome 5AL for linkage to *Nax2*. Microsatellite markers *gwm291*, *gwm410*, and *gpw2181*, previously mapped to the distal region of chromosome 5AL (Roder et al., 1998), cosegregated with *Nax2* in the segregating family of 137 F₃ lines. Therefore, *Nax2* maps to the location that we might expect to find a homoeolog of *Kna1*.

The lack of recombination between the *HKT1;5* gene on the A genome (*HKT1;5-A*) and *gwm291*, *gwm410*, and *gpw218* raised the question whether the introgressed segment carrying *Nax2* from *T. monococcum* was able to recombine with the homologous region in

Tamaroi. To investigate this, we tested other microsatellite markers that were previously positioned on the distal end of chromosome 5AL. Four microsatellite markers, *gwm595*, *gwm179*, *gwm126* (Roder et al., 1998), and *barc232* (<http://wheat.pw.usda.gov>) were polymorphic between Line 149 and Tamaroi but were not retained in the BC₄ parent and therefore failed to segregate in the BC-derived family (C.S. Byrt, unpublished data). These results show that recombination did occur between the *Nax2* gene and *gwm595* and the chromosomal region on 5AL proximal to *gwm2181* was replaced by Tamaroi during the process of five backcrossing steps of transferring *Nax2* into Tamaroi (Fig. 5).

In summary, *Nax2* is located on the ancestral segment of chromosome 4AL that is attached to the distal end of chromosome 5AL. Hence, the tightly linked *TmHKT1;5-A* gene member from *T. monococcum* was also located on chromosome 5AL, consistent with the map locations of other *TaHKT1;5* members on homoeologous chromosomes 4BL and 4DL.

Isolation of Full-Length *HKT1;5* Gene Members

The partial *HKT1;5* probe was used to screen a cDNA library from root tissue of CS. Several positive phagemid clones were isolated and when sequenced revealed identical DNA sequences with insert size varying between 812 and 1,741 bp. The cDNA sequences were identical to genomic sequence derived from a bacterial artificial chromosome clone that was previously isolated from a bacterial artificial chromosome library made from the diploid D genome progenitor species *Aegilops tauschii* Coss (E.S. Lagudah, unpublished data), suggesting that the cDNA sequence isolated from CS was derived from the D genome (*TaHKT1;5-D*). The cDNA was predicted to encode a full-length gene based on the comparison of its predicted amino acid sequence to *OsHKT1;5* (*SKC1*) in rice

(C.S. Byrt, unpublished data). Reverse-transcriptase PCR with primers designed from the 5' and 3' untranslated regions of *TaHKT1;5-D* amplified the corresponding A gene member, *TmHKT1;5-A*, from *T. monococcum* and Line 149. The predicted ORF of *TmHKT1;5-A* is 1,554 bp and the predicted ORF of *TaHKT1;5-D* is 1,551 bp. *OsHKT1;5*, *TmHKT1;5-A*, and *TaHKT1;5-D* each have two introns. The predicted amino acid sequence of *TmHKT1;5-A* and *TaHKT1;5-D* shared 94% identity and were closely related to the rice Na⁺ transporter *OsHKT1;5* (66% identity). The intron and exon structure of the *TmHKT1;5-A*, *TaHKT1;5-D*, and *OsHKT1;5* genes are shown in Figure 6.

Expression of *HKT1;5* Gene Members


Reverse-transcriptase PCR with specific primers for A and D genome members was used to analyze the expression of the *HKT1;5* A and D gene members in *T. monococcum*, Line 149, Tamaroi, and CS. *TmHKT1;5-A* was expressed in the roots of *T. monococcum* and Line 149, but not in the shoots (Fig. 7A). *HKT1;5-A* was not expressed in Tamaroi or CS. The *TaHKT1;5-D* gene member was expressed in CS roots but not shoots (Fig. 7B). The *TaHKT1;5-D* gene member was not expressed in the CS deletion line missing the distal 14% of chromosome 4DL (0.86; Fig. 7B). The expression results for the *HKT1;5-D* gene member are consistent with the mapping results, indicating that *HKT1;5-D* is missing from the CS deletion line 0.86 and is therefore positioned in the same region as *Kna1*.

DISCUSSION

Mapping of *Nax2*

Nax2 is a single dominant gene, as *Nax2* segregated in a 3:1 ratio of low:high leaf blade Na⁺ concentration

Figure 4. A, Southern-blot hybridization with *HKT1;5* probe after *Hind*III restriction digest of DNA from CS and CS chromosome deletion lines 0.86, 0.70, 0.61, and 0.56. The *TaHKT1;5-D* gene member maps to the most distal deletion bin. The genome location of the *TaHKT1;5* gene members are shown on the right. B, CS and CS chromosome deletion lines were grown in salt tanks (see "Materials and Methods"). F1 describes remaining fraction of chromosome 4DL. *Kna1* maps to the same location (Dubcovsky et al., 1996) as *TaHKT1;5-D*. Leaf blade Na⁺ and K⁺ concentrations were recorded after leaf 3 had grown for 10 d in 50 mM NaCl.






Figure 5. Part of chromosome 5AL originated from part of chromosome 4AL due to an ancient reciprocal translocation between the distal ends of chromosomes 4AL and 5AL. Durum 5AL represents the fragment containing *Nax2* that introgressed from Line 149 into the Tamaroi background in the *Nax2* mapping population. Proximal to the introgression is *gwm595*. *Nax2* is linked to *gwm291*, *gwm410*, and *gpw2181*. (See Somers et al. [2004].)

in the BC₅F₂ plants (Fig. 1). *Nax2* is located on the distal part of chromosome 5AL. The *HKT1;5* gene member on the A genome mapped to the same region as *Nax2* and cosegregated with *Nax2*.

HKT genes were considered to be the best candidates for *Nax2* based on the role of other HKT transporters in higher plants. *HKT7* cosegregates with *Nax1*, which confers Na⁺ exclusion from leaves in wheat (Huang et al., 2006), *AtHKT1;1* confers Na⁺ exclusion from leaves in Arabidopsis (Uozumi et al., 2000; Sunarpi et al., 2005), and *OsHKT1;5* (*SKC1*) confers Na⁺ exclusion from leaves in rice (Ren et al., 2005).

There are up to five *HKT1;5* genes in wheat, a partial wheat *HKT1;5* probe detected one gene member on the D genome, two or three on the B genome, and one gene member on the A genome derived from *T. monococcum* (C68-101). The predicted amino acid identity between the wheat *HKT1;5-A* and *HKT1;5-D* gene members is 94%. The most closely related gene in rice, *OsHKT1;5*, shares 66% identity (75% positives) with the predicted wheat *HKT1;5* sequences.

The location of the wheat *HKT1;5* genes and the location of the rice *OsHKT1;5* genes are not syntenic. *OsHKT1;5* is located on chromosome 1S of rice (Lin et al., 2004; Ren et al., 2005). This region is syntenous to chromosome 3 in wheat (Sorrells et al., 2003), however, *TmHKT1;5-A* is located on chromosome 5AL and *TaHKT1;5-D* is located on chromosome 4DL.

There may be a second gene in the *Nax2* region having an effect on leaf Na⁺, but it would have to be closely linked, as every time that the *HKT1;5-A* gene member is lost, leaf Na⁺ increases significantly. This has been demonstrated in the *Nax2* mapping population (Fig. 2) and when *HKT1;5-A* was transferred into other genetic backgrounds including bread wheat. In moderately saline field conditions (100 mM) Tamaroi had an average leaf Na⁺ concentration of 125 μmol g dry weight⁻¹ whereas BC₄F₂-derived lines with *HKT1;5-A* had an average leaf Na⁺ concentration of 25 μmol g dry

weight⁻¹, a 5-fold difference (R. Hare and R. Munns, unpublished data). In a completely different field environment with the same lines *HKT1;5* conferred a 2-fold reduction in leaf Na⁺ concentration (A. Rathjen and R. Munns, unpublished data).


Overall, in field and glasshouse experiments, all the lines with the *HKT1;5-A* gene had at least 2-fold lower leaf Na⁺ than all those without the *HKT1;5-A* gene. When transferred into the bread wheat Westonia, which already contains the Na⁺ exclusion locus *Kna1* on the D genome, *HKT1;5-A* reduced leaf Na⁺ by a further 25% (R. Munns and R. James, unpublished data).

Relationship of *Nax2* to the Major Salt-Tolerant Gene in Hexaploid Wheat, *Kna1*

The *HKT1;5-A* genome member, which is the candidate for *Nax2*, is physically located on the distal part of chromosome 5AL, which ancestrally corresponds to the distal part of chromosome 4AL (Fig. 7). *Kna1* maps to the distal region of chromosome 4DL of wheat (Dubcovsky et al., 1996), raising the possibility that *Nax2* and *Kna1* are homoeologous genes located on ancestral group 4 chromosomes. The *TaHKT1;5-D* genome member maps to the distal 14% of chromosome 4DL in hexaploid bread wheat (Fig. 4A), coinciding with the map location of *Kna1* (Dubcovsky et al., 1996). The locus for *Kna1* on chromosome 4DL is homoeologous to the location for *Nax2* on chromosome 5AL and the *TaHKT1;5-D* genome member may be *Kna1*.

The loss of the region containing the *TaHKT1;5-D* gene member from CS deletion lines corresponded to an increase in average Na⁺ concentrations in the leaf blade and a 6-fold decrease in the K⁺-to-Na⁺ ratio from 7.5 to 1.2 (Fig. 4B), when plants were grown at 50 mM NaCl. These results are consistent with our hypothesis that the *HKT1;5* probe detects not only a candidate gene for *Nax2* in durum wheat, but also a

Figure 6. Gene structures of *TmHKT1;5-A*, *TaHKT1;5-D*, and *OsHKT1;5*. The gray triangles represent the intron regions of the gene. The arrows indicate the primers designed for gene expression analysis (Fig. 7).



candidate gene for *Kna1* in hexaploid bread wheat and that both genes are located in homoeologous regions of the wheat genome. These results are also consistent with other data on the effect of *Kna1* on leaf K^+ and Na^+ concentrations. When the *Kna1* region was transferred from the bread wheat, CS, into the durum wheat cultivar, Langdon, lines with *Kna1* had a greater leaf K^+ -to- Na^+ ratio (Dvořák and Gorham, 1992). Lines of Langdon containing the *Kna1* region had a leaf K^+ -to- Na^+ ratio approximately 8 times higher than those without *Kna1* when plants were grown at 50 mM, and 6 times higher when grown at 150 mM NaCl (Gorham et al., 1987, 1990).

Similarity of Phenotype between Sodium Excluding Genes in Rice and Wheat

There are five phenotypic characteristics in common between *Nax2*, *Kna1*, and *SKC1*: (1) low Na^+ concentration in the leaves; (2) enhanced discrimination of K^+ over Na^+ in transport from the roots to the shoots; (3) regulation of the K^+ -to- Na^+ ratio in the leaves; (4) no effect on root Na^+ concentration; and (5) no effect on the sheath-to-blade Na^+ ratio (Gorham et al., 1990; Davenport et al., 2005; Ren et al., 2005; James et al., 2006). *Nax1*, which cosegregates with *T. monococcum* *HKT7*, has a distinctive mechanism that differs from *Nax2*, *Kna1*, and *SKC1*. The characteristic of *Nax1* is a high sheath-to-blade Na^+ concentration ratio. *Nax1* confers low leaf blade Na^+ and high leaf blade K^+ -to- Na^+ ratio by unloading Na^+ in the leaf sheath and displacing K^+ in the leaf blade (James et al., 2006). In the Tamaroi background, BC_5F_2 lines with *Nax1* (*HKT7-A2*) have a 4-fold lower leaf Na^+ concentration than lines without *Nax1* (Huang et al., 2006).

The mechanism behind the common phenotype for the *Nax2*, *SKC1*, and *Kna1* genes may be unloading of Na^+ from the xylem. We know that *Nax2* unloads Na^+ from the xylem as experiments with $^{22}Na^+$ showed that the rate of unloading of Na^+ from the xylem was double that in lines with *Nax2* than in those without (James et al., 2006). *SKC1* in rice also unloads Na^+ from the xylem (Ren et al., 2005). Under salt stress the shoots and the xylem sap of near-isogenic lines with *SKC1*

contained more K^+ and less Na^+ than those without *SKC1* (Ren et al., 2005). There is also indirect evidence that *Kna1* may unload Na^+ from the xylem. One is the observation that there is no difference in the Na^+ concentration in the roots of lines with and without *Kna1* (Gorham et al., 1990), and the other is that the membrane potential-dependent uptake of Na^+ was no different in vesicles from hexaploid wheat (cv Troy), which has *Kna1*, and tetraploid wheat (cv Langdon), which does not have *Kna1* (Allen et al., 1995). Gorham et al. (1997) suggested that this observation of Allen et al. (1995) would be expected if *Kna1* acts specifically on xylem loading.

A single *HKT* gene may be sufficient to explain the *Nax2* or *Kna1* phenotypes. The *Nax1* gene in wheat and the *SKC1* gene in rice are both *HKT* genes and they both have a strong effect on the leaf K^+ -to- Na^+ ratio and Na^+ concentration. Wheat lines with *Nax1* (*TmHKT7-A2*) have a 4 times greater leaf K^+ -to- Na^+ ratio, and 4 times less leaf Na^+ than lines without *Nax1* (Huang et al., 2006; James et al., 2006). Transformed rice lines with *SKC1* (*OsHKT1;5*) have a 2.5 times greater leaf K^+ -to- Na^+ ratio and 2 times less leaf Na^+ than transformed lines without *SKC1* (Ren et al., 2005). Wheat lines with *Nax2* have a 3.6 times greater leaf K^+ -to- Na^+ ratio and approximately 2.5 times less leaf Na^+ than lines without *Nax2*. One *HKT1;5-A* gene could account for these differences. Likewise, for *Kna1*, one *HKT1;5-D* gene may account for the 6 times greater leaf K^+ -to- Na^+ ratio and the 4 times less leaf Na^+ in CS compared to the CS deletion lines lacking *Kna1*.

Allelic Variation in *HKT* Genes

In rice and Arabidopsis, allelic variation for *OsHKT1;5* and *AtHKT1;1*, respectively, has been linked to variation in function (Ren et al., 2005; Rus et al., 2006). In rice, six nucleotide substitutions in the coding region of *OsHKT1;5* differ between the tolerant allele from Nona Bokra and the sensitive allele from Koshihikari, leading to four amino acid residues differing (Ren et al., 2005). Under salt stress Koshihikari had almost double the leaf Na^+ concentration of near isogenic lines with the Nona Bokra *SKC1* (*OsHKT1;5*) allele.




Figure 7. Expression of *HKT1;5* in wheat analyzed using reverse-transcriptase PCR. A, Expression of *TmHKT1;5-A*. A product of the expected size (442 bp) was observed for the *T. monococcum* and Line 149 root samples. B, Expression of *TaHKT1;5-D*. A product of the expected size (147 bp) was observed for the CS root sample. Mon = *T. monococcum*; L149 = Line 149; Tam = Tamaroi; 0.86 = CS deletion line missing the distal 14% of chromosome 4DL; n = no template control; R = root tissue; L = leaf tissue. Plants were grown in hydroponic solution for 2 weeks before addition of 50 mM NaCl, tissue was harvested 48 h after addition of NaCl. Proton pump was used as a control.

Changes in the coding region, rather than differences in the promoter, are likely to account for the functional variation in the two alleles (Ren et al., 2005). The data that support this include a genetic complementation test, where the Nona Bokra promoter and ORF was transferred into a sensitive line and recovered the *SKC1* phenotype, and an observation that there were no differences in the expression patterns for the different *OshKT1;5* alleles (Ren et al., 2005).

In Arabidopsis, allelic variation in the promoter of *AtHKT1;1*, rather than the coding region, may account for functional differences between *AtHKT1;1* from Columbia-0 and two natural variants, *Tsu-1* and *Ts-1* (Rus et al., 2006). The *AtHKT1;1* alleles from *Tsu-1* and *Ts-1* are linked to higher leaf Na⁺. Sequence differences included 19 single nucleotide polymorphisms in the coding region, leading to seven different amino acid residues (Rus et al., 2006), but a deletion in a tandem repeat sequence in the promoter may be responsible for the high leaf Na⁺ as there was reduced expression of *AtHKT1;1* in the roots (Rus et al., 2006). Curiously, the *Tsu-1* allele linked to higher leaf Na⁺ was also linked to NaCl tolerance, whereas the loss of function mutant *Athkt1;1* has higher leaf Na⁺ than wild type (Columbia-0) but is more NaCl sensitive (Rus et al., 2004), although, as Rus et al. (2006) note, a second gene that influences NaCl tolerance may be segregating with *AtHKT1;1* in the *Tsu-1* population.

In contrast, no allelic variation for *HKT1;5-A* has been identified in wheat. The difference in leaf Na⁺ between lines with and without *HKT1;5-A* seems to be determined by the presence or absence of the gene, rather than allelic variation (Fig. 2). We found a single *HKT1;5* gene member on the A genome of families containing *Nax2* that was absent in those without *Nax2*. This gene member originated in *T. monococcum* (C68-101), was introgressed into Line 149 by crossing *T. monococcum* with the durum cultivar Marrocos (The,

1973), and inherited in all the near-isogenic lines containing *Nax2* that were derived from the cross between Line 149 and cv Tamaroi (James et al., 2006). There is no homoeolog of *HKT1;5* on the A genome of Marrocos or Tamaroi, or in the durum cv Langdon, or the bread wheat CS (Fig. 3A). It is absent in all the modern wheats that we have tested (data not shown). We suggest that the *HKT1;5* gene was not present in the A genome diploid ancestor that gave rise to modern wheat. An extensive screen of diverse genetic material to look for allelic variation for *HKT1;5-A* in wheat ancestors and wild A genome species, including *T. monococcum* and *Triticum urartu*, is under way.

The *HKT1;5-D* allele isolated from CS has the same ORF as the *HKT1;5-D* gene members isolated from three *Triticum tauschii* accessions (data not shown). Screening for allelic variation in *HKT1;5-D* in a diverse range of genetic material is also under way.

Concluding Remarks

The data presented in this work supports the hypothesis that *TmHKT1;5-A* is a candidate for *Nax2* and is a homoeolog of *Kna1*. To test whether *Nax2* is *TmHKT1;5-A* and whether *Kna1* is *TaHKT1;5-D*, functional assays of the *HKT1;5-A* and *HKT1;5-D* transporters are necessary. RNA interference constructs against *TaHKT1;5-D* will be introduced into hexaploid bread wheat to test if silencing *TaHKT1;5-D* results in the same phenotype that we have observed in the CS deletion lines lacking the *Kna1* region. We will also express the wheat *HKT1;5* A, B, and D gene members in yeast and *X. laevis* oocytes to characterize the transport properties of the proteins.

Incorporation of *HKT1;5* into durum wheat breeding programs may provide a mechanism for Na⁺ exclusion for durum wheat of similar potency to that conferred by *Kna1* in bread wheat.

MATERIALS AND METHODS

Plant Material

Parent material for the mapping population was durum wheat (*Triticum turgidum* subsp. *durum* [Desf.] Line 149; Munns et al., 2000) and the cultivar Tamaroi. Line 149 was derived from *Triticum monococcum* (C68-101; low leaf Na⁺) and the durum variety Marrocos (high leaf Na⁺; *T. monococcum*/3*Marrocos; The, 1973). *T. monococcum* C68-101 is the putative donor of *Nax1* and *Nax2* (James et al., 2006). Selected F₂ lines from the cross between Line 149 and Tamaroi were backcrossed into Tamaroi four times. A selected BC₄F₂ individual that carried *Nax2*, but not *Nax1*, was backcrossed once more into Tamaroi to produce a BC₅F_{2,3} family of 137 F₂ individuals (James et al., 2006). This family containing the *Nax2* gene but not the *Nax1* gene formed the basis for this study.

Plant material for the CS deletion line experiment included lines with the following Fls: 0.09, 0.31, 0.38, 0.41, 0.46, 0.53, 0.56, 0.61, 0.70, 0.71, 0.86, and 1.00 (wild-type CS; Endo and Gill, 1996).

Plant material for DNA extraction for Southern-blot hybridization work included CS nullisomic-tetrasomic lines (Sears, 1954) and CS deletion lines (Endo and Gill, 1996) and Langdon substitution lines (Joppa, 1987). The Langdon substitution lines included 4D(4A) and 4D(4B), where chromosome 4D from CS had been substituted for chromosome 4A or 4B of Langdon, respectively.

Phenotyping

Plants were grown in half-strength Hoagland solution in supported hydroponics in a method adapted from Munns et al. (2000). Salt treatment commenced when leaf 2 was half emerged. The NaCl concentration of the hydroponic solution was increased by 25 mM twice daily over 3 d to reach a final concentration of 150 mM. Supplemental calcium [Ca(NO₃)₂] was added to achieve a Na⁺-to-Ca²⁺ ratio of 15:1. Leaf 3 was harvested after 10 d growing in salt and the Na⁺ was extracted with nitric acid (0.5 M). The Na⁺ concentration was measured using inductively coupled plasma analysis. Parental lines were replicated 10 times. Based on the score of F₂ individuals, nine plants from every family predicted to be homozygous low were tested and five plants were tested for every family predicted to be homozygous high or heterozygous. These numbers were based upon recommended population sizes required in biparental populations to obtain at least one target homozygous genotype in later generations for segregating loci (Bonnert et al., 2005).

Genotyping

Plants grown in salt tanks for phenotyping were transplanted into soil and allowed to grow for approximately 4 weeks prior to DNA extraction. One or two plants were retained from each of the F_{2,3} families. For families with a homozygous low Na⁺ accumulation phenotype, the plant with the lowest leaf Na⁺ concentration was used. The plant with the highest leaf Na⁺ was used from families with a homozygous high Na⁺ accumulation phenotype. For half of those families with a heterozygous phenotype, the lowest of the low Na⁺ accumulating plants was used, and for the other half of the families with a heterozygous phenotype, the highest of the high Na⁺ accumulating plant was used. Leaf material from plants was harvested and DNA extracted as per Lagudah et al. (1991).

RFLP Probe Development

A search of the public database identified a wheat EST (CK193616) with strong homology, 86% nucleotide sequence identity, to the rice *SKC1* candidate gene (DQ148410) subsequently named *OsHKT1;5* (Platten et al., 2006). The wheat EST was named *HKT1;5* as recommended by Platten et al. (2006). Primers designed internal to CK193616 were HKT1;5For01 (5'-CATCACCGTCGAGGTTATCAG-3') and HKT1;5Rev01 (5'-TTGAGGTACTCGGCATA-3'). These primers were used to amplify a 332 bp product from *T. monococcum* genomic DNA. The product was cloned into pGEMT-easy vector (Promega Corporation). The PCR-amplified product was radiolabeled with ³²P-CTP using the Megaprime DNA labeling system (Amersham) according to the manufacturer's instructions.

The *HKT1;5* probe spans from exon 2, 1,692 bp into the ORF, to the 3' untranslated region. Three additional probes were developed using the same method. The additional probes were named *HKT1;5* probe 2, *HKT1;5* probe 3,

and *HKT1;5* probe 4. They were used to confirm the mapping and cosegregation data. The sizes of *HKT1;5* probe 2, *HKT1;5* probe 3, and *HKT1;5* probe 4 were 315, 324, and 321 bp in, respectively. Relative to the ORF they start at 935, 1,003, and 1,765 bp, respectively. Together, the four probes span from exon 1 through to the 3' untranslated region.

DNA from the parental lines (Line 149 and Tamaroi) *T. monococcum* (AUS# 90382) and the BC₅F_{2,3} progeny were digested individually with *Hind*III and/or *Eco*RV, *Eco*RI, *Nco*I, *Sac*I, and/or *Xba*I. After gel electrophoresis, the gels were blotted onto a nitrocellulose membrane (Amersham Biosciences Hybond-N⁺) and hybridized with the *HKT1;5* probe. DNA samples from wheat nullisomic-tetrasomic lines and ditelomeric lines in a CS background were also screened with the *HKT1;5* probe as described above. In the nullisomic-tetrasomic lines each pair of homoeologous chromosomes, 4A, 4B, or 4D have been substituted by one of the other pairs, in the ditelomeric lines the short or long chromosome arms have been deleted.

Microsatellite Markers

Microsatellite markers were used to establish the chromosomal location of *Nax2* in the durum Line 149. A group of 470 wheat microsatellite markers were used to screen DNA from the parental lines, Tamaroi and Line 149, for polymorphisms (Roder et al., 1998). Twenty five were found to be polymorphic between Line 149 and Tamaroi (C.S. Byrt, unpublished data). These were tested on bulked DNA samples from the BC₅F_{2,3} lines, two bulks each containing pooled DNA from 10 low Na⁺ lines and two bulks each pooled from 10 high Na⁺ lines. Three wheat microsatellite markers cosegregated with *Nax2*, namely *gwm291*, *gwm410*, and *gpw2181*. These markers had been mapped previously to the distal end of chromosome 5AL (Roder et al., 1998).

To confirm cosegregation of these markers with *Nax2*, the genotype of 19 BC₅F_{2,3} plants with a homozygous low Na⁺ phenotype were tested using the markers *gwm291*, *gwm410*, and *gpw2181*. All 19 had the same genotype as Line 149. In contrast, 11 BC₅F_{2,3} plants that had a homozygous high Na⁺ phenotype all had the same genotypic pattern with *gwm291*, *gwm410*, and *gpw2181* as Tamaroi.

Isolation of *TaHKT1;5-D*

The CS cDNA library was kindly supplied by Professor Timothy J. Close. The library was constructed from drought-stressed root tissue at full tillering. Approximately 120,000 clones of the mass excised phagemid library were plated and screened with the partial *TmHKT1;5* probe according to standard protocols (Sambrook et al., 1989). Eleven colonies hybridized to the probe. These were grown at 37°C overnight in Luria broth (Sambrook et al., 1989) and plasmid DNA was isolated using a Qiaspin miniprep kit according to the manufacturer's instructions (Qiagen). Plasmid DNA was digested with *Eco*RI and *Xho*I to liberate the cloned inserts. The inserts were sequenced using T7 and SP6 universal primers (Invitrogen).

Isolation of RNA, Reverse-Transcriptase PCR, and Isolation of *TmHKT1;5-A*

Plants were grown in hydroponic solution described in "Plant Material." After 2 weeks plants were exposed to 50 mM NaCl. After 48 h leaf and root tissues were harvested separately and snap frozen in liquid nitrogen. RNA was extracted using TRIzol Reagent (Invitrogen) as per the manufacturer's instructions. Reverse-transcriptase PCR to amplify *TmHKT1;5* was undertaken using primers that were external to the coding sequence named 5primeUTRFor (5'-AGAAGTCTCTACACAACCTTACAG-3') and 3primeUTRRev (5'-GAT-CATTGAGAAATATGCAGTCC-3') using a Qiagen OneStep RT-PCR kit as per the manufacturer's instructions. DNA fragments of the appropriate size were amplified from *T. monococcum* and Line 149. These fragments were cut out and purified using a Qiagen gel extraction kit according to the manufacturer's instructions. The fragments were ligated into the pGEM-T vector using the pGEM-T easy vector system 1 kit (Promega).

Reverse-transcriptase PCR to observe presence or absence of expression of the *HKT1;5* A and D gene homoeologs was undertaken using A gene specific primers named ForA1 (5'-GAGTGGGCTCCGACGGCTGAA-3') and RevA1 (5'-CGTCAGGCGTCACTGCCGGCCG-3') and D gene specific primers ForD1 (5'-GCTTGGCCATCTTCATCGCCGTG-3') and RevD1 (5'-GGCCA-CAGCTGTACCCGGTGCTG-3') using a Qiagen OneStep RT-PCR kit as per the manufacturer's instructions. Primer locations are shown in Figure 6. The

PCR was conducted under standard conditions with the following cycling protocol: 50°C, 20 min; 94°C, 15 min; then 35 cycles of 94°C, 30 s; 58°C, 30 s; 68°C, 1 min; and finally 72°C, 2 min. The forward and reverse primers in each primer set were designed in different exons so as to include an intron in between them. Therefore, products that amplified from trace DNA in the RNA samples differed in size from the products amplified from coding DNA. The expected product size for the A gene specific primers was 942 bp from genomic DNA and 442 bp from coding DNA. The expected product size for the D gene specific primers was 322 bp from genomic DNA and 147 bp from coding DNA.

Sequence data from this article can be found in the GenBank/EMBL data libraries under accession numbers DQ646339 (*TmHKT1;5-A*) and DQ646342 (*TaHKT1;5-D*).

ACKNOWLEDGMENTS

We thank Dr. Ray Hare for providing Tamaroi, Line 149, and *T. monococcum* lines, with information of the unusual pedigree of Line 149, Professor Timothy J. Close for the CS cDNA library, Lorraine Mason for Na⁺ and K⁺ analysis by inductively coupled plasma, Karen Glover and Kylie Groom for screening microsatellite markers, Marianne Bloemsa for expert technical assistance, and Dr. Shaobai Huang for scholarly and methodological advice.

Received November 21, 2006; accepted February 3, 2007; published February 23, 2007.

LITERATURE CITED

- Allen GJ, Wyn Jones RG, Leigh RA (1995) Sodium transport measured in plasma membrane vesicles isolated from wheat genotypes with differing K⁺/Na⁺ discrimination traits. *Plant Cell Environ* 18: 105–115
- Berthomieu P, Conéjéro G, Nublát A, Brackenbury WJ, Lambert C, Savio C, Uozumi N, Oiki S, Yamada K, Cellier F (2003) Functional analysis of *AtHKT1* in *Arabidopsis* shows that Na⁺ recirculation by the phloem is crucial for salt tolerance. *EMBO J* 22: 2004–2014
- Bonnett DG, Rebetzke GJ, Spielmeier W (2005) Strategies for efficient implementation of molecular markers in wheat breeding. *Mol Breed* 15: 75–85
- Davenport R, James RA, Zakrisson-Plogander A, Tester M, Munns R (2005) Control of sodium transport in durum wheat. *Plant Physiol* 137: 807–818
- Dubcovsky J, María GS, Epstein E, Luo MC, Dvořák J (1996) Mapping of the K⁺/Na⁺ discrimination locus *Kna1* in wheat. *Theor Appl Genet* 92: 448–454
- Dvořák J, Gorham J (1992) Methodology of gene-transfer by homoeologous recombination into *Triticum-turgidum*—transfer of K⁺/Na⁺ discrimination from *Triticum-aestivum*. *Genome* 35: 639–646
- Endo T, Gill B (1996) The deletion stocks of common wheat. *J Hered* 87: 295–307
- Garciadeblas B, Senn ME, Banuelos MA, Rodriguez-Navarro A (2003) Sodium transport and HKT transporters: the rice model. *Plant J* 34: 788–801
- Golldack D, Su H, Quigley F, Kamasani UR, Munoz-Garay C, Balderas E, Popova OV, Bennett J, Bohnert HJ, Pantoja O (2002) Characterization of a HKT-type transporter in rice as a general alkali cation transporter. *Plant J* 31: 529–542
- Gorham J, Bridges J, Dubcovsky J, Dvorak J, Hollington PA, Luo M-C, Khan JA (1997) Genetic analysis and physiology of a trait for enhanced K⁺/Na⁺ discrimination in wheat. *New Phytol* 137: 109–116
- Gorham J, Hardy C, Wyn Jones RG, Joppa LR, Law CN (1987) Chromosomal location of a K/Na discrimination character in the D genome of wheat. *Theor Appl Genet* 74: 584–588
- Gorham J, Wyn Jones RG, Bristol A (1990) Partial characterization of the trait for enhanced K⁺-Na⁺ discrimination in the D genome of wheat. *Planta* 180: 590–597
- Haro R, Banuelos MA, Senn ME, Barrero-Gil J, Rodriguez-Navarro A (2005) HKT1 mediates sodium uniport in roots: pitfalls in the expression of HKT1 in yeast. *Plant Physiol* 139: 1495–1506
- Horie T, Yoshida K, Nakayama H, Yamada K, Oiki S, Shinmyo A (2001) Two types of HKT transporters with different properties of Na⁺ and K⁺ transport in *Oryza sativa*. *Plant J* 27: 129–138
- Huang S, Spielmeier W, Lagudah ES, James RA, Platten JD, Dennis ES, Munns R (2006) A sodium transporter (HKT7) is a candidate for *Nax1*, a gene for salt tolerance in durum wheat. *Plant Physiol* 142: 1718–1727
- James RA, Davenport RJ, Munns R (2006) Physiological characterisation of two genes for Na⁺ exclusion in durum wheat: *Nax1* and *Nax2*. *Plant Physiol* 142: 1537–1547
- Joppa LR (1987) Aneuploid analysis in tetraploid wheat. In Heyne EG, ed, *Wheat and Wheat Improvement*, Monograph 13. Am Soc Agron, Madison, WI, pp 151–166
- Lagudah ES, Appels R, Brown A, McNeil D (1991) The molecular-genetic analysis of *Triticum tauschii*, the D-genome donor to hexaploid wheat. *Genome* 34: 375–386
- Lin HX, Zhu MZ, Yano M, Gao JP, Liang ZW, Su WA, Hu ZH, Ren ZH, Chao DY (2004) QTLs for Na⁺ and K⁺ uptake of the shoots and roots controlling rice salt tolerance. *Theor Appl Genet* 108: 253–260
- Lindsay MP, Lagudah ES, Hare RA, Munns R (2004) A locus for sodium exclusion (*Nax1*), a trait for salt tolerance, mapped in durum wheat. *Funct Plant Biol* 31: 1105–1114
- Liu CJ, Atkinson MD, Chinoy CN, Devos KM, Gale MD (1992) Non-homoeologous translocations between group 4, 5 and 7 chromosomes within wheat and rye. *Theor Appl Genet* 83: 305–312
- Luo MC, Dubcovsky J, Goyal S, Dvořák J (1996) Engineering of interstitial foreign chromosome segments containing the K⁺/Na⁺ selectivity gene *Kna1* by sequential homoeologous recombination in durum wheat. *Theor Appl Genet* 93: 1180–1184
- Mäser P, Gierth M, Schroeder JI (2002a) Molecular mechanisms of potassium and sodium uptake in plants. *Plant Soil* 247: 43–54
- Mäser P, Hosoo Y, Goshima S, Horie T, Eckelman B, Yamada K, Yoshida K, Bakker EP, Shinmyo A, Oiki S, et al (2002b) Glycine residues in potassium channel-like selectivity filters determine potassium selectivity in four-loop-per-subunit HKT transporters from plants. *Proc Natl Acad Sci USA* 99: 6428–6433
- Munns R, Hare RA, James RA, Rebetzke GJ (2000) Genetic variation for improving the salt tolerance of durum wheat. *Aust J Agric Res* 51: 69–74
- Munns R, James RA, Läuchli A (2006) Approaches to increasing the salt tolerance of wheat and other cereals. *J Exp Bot* 57: 1025–1043
- Munns R, Rebetzke GJ, Husain S, James RA, Hare RA (2003) Genetic control of sodium exclusion in durum wheat. *Aust J Agric Res* 54: 627–635
- Nelson JC, Sorrells ME, Van-Deynze AE, Lu YH, Atkinson M, Bernard M, Leroy P, Faris JD, Anderson JA (1995) Molecular mapping of wheat: major genes and rearrangements in homoeologous groups 4, 5, and 7. *Genetics* 141: 721–731
- Platten JD, Cotsaftis O, Berthomieu P, Bohnert H, Davenport RJ, Fairbairn DJ, Horie T, Leigh RA, Lin HX, Luan S (2006) Nomenclature for HKT transporters, key determinants of plant salinity tolerance. *Trends Plant Sci* 11: 372–374
- Ren ZH, Gao JP, Li LG, Cai XL, Huang W, Chao DY, Zhu MZ, Wang ZY, Luan S, Lin HX (2005) A rice quantitative trait locus for salt tolerance encodes a sodium transporter. *Nat Genet* 37: 1141–1146
- Roder MS, Korzun V, Wendehake K, Plaschke J, Tixier M-H, Leroy P, Ganal MW (1998) A microsatellite map of wheat. *Genetics* 149: 2007–2023
- Rodriguez-Navarro A, Rubio F (2006) High-affinity potassium and sodium transport systems in plants. *J Exp Bot* 57: 1149–1160
- Rubio F, Gassmann W, Schroeder JI (1995) Sodium-driven potassium uptake by the plant potassium transporter HKT1 and mutations conferring salt tolerance. *Science* 270: 1660–1663
- Rus A, Baxter I, Muthukumar B, Gustin J, Lahner B, Yakubova E, Salt DE (2006) Natural variants of *AtHKT1* enhance Na⁺ accumulation in two wild populations of *Arabidopsis*. *PLoS Genetics* 2: 1964–1973
- Rus A, Lee B-h, Munoz-Mayor A, Sharkhuu A, Miura K, Zhu J-K, Bressan RA, Hasegawa PM (2004) *AtHKT1* facilitates Na⁺ homeostasis and K⁺ nutrition in planta. *Plant Physiol* 136: 2500–2511
- Sambrook J, Fritsch EF, Maniatis T (1989) *Molecular Cloning: A Laboratory Manual*, Ed 2. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY
- Schachtman DP, Schroeder JI (1994) Structure and transport mechanism of a high-affinity potassium uptake transporter from higher-plants. *Nature* 370: 655–658
- Sears ER (1954) The aneuploids of common wheat. University of Missouri-Columbia Agricultural Experiment Station Bulletin 572: 1–59

- Somers D, Isaac P, Edwards K** (2004) A high-density microsatellite consensus map for bread wheat (*Triticum aestivum* L.). *Theor Appl Genet* **109**: 1105–1114
- Sorrells ME, La Rota M, Bermudez-Kandianis CE, Greene RA, Kantety R, Munkvold JD, Miftahudin, Mahmoud A, Ma X, Gustafson PJ** (2003) Comparative DNA sequence analysis of wheat and rice genomes. *Genome Res* **13**: 1818–1827
- Sunarpi, Horie T, Motoda J, Kubo M, Yang H, Yoda K, Horie R, Chan WY, Leung HY, Hattori K** (2005) Enhanced salt tolerance mediated by AtHKT1 transporter-induced Na⁺ unloading from xylem vessels to xylem parenchyma cells. *Plant J* **44**: 928–938
- Tester M, Davenport R** (2003) Na⁺ tolerance and Na⁺ transport in higher plants. *Ann Bot (Lond)* **91**: 503–537
- The TT** (1973) Transfer of resistance to stem rust from *Triticum monococcum* L. to hexaploid wheat. PhD thesis. University of Sydney, Sydney
- Uozumi N, Kim EJ, Rubio F, Yamaguchi T, Muto S, Tsuboi A, Bakker EP, Nakamura T, Schroeder JI** (2000) The *Arabidopsis* HKT1 gene homolog mediates inward Na⁺ currents in *Xenopus laevis* oocytes and Na⁺ uptake in *Saccharomyces cerevisiae*. *Plant Physiol* **122**: 1249–1259
- Walker NA, Maathuis FJM, Rubio F, Gassmann W, Schroeder JI** (1996) High-affinity potassium uptake in plants. *Science* **273**: 977–979
- Wang T-B, Gassmann W, Rubio F, Schroeder JI, Glass ADM** (1998) Rapid up-regulation of HKT1, a high-affinity potassium transporter gene, in roots of barley and wheat following withdrawal of potassium. *Plant Physiol* **118**: 651–659