

Legume Transcription Factor Genes: What Makes Legumes So Special?^{1[W]}

Marc Libault, Trupti Joshi, Vagner A. Benedito, Dong Xu, Michael K. Udvardi, and Gary Stacey*

Division of Plant Sciences, National Center for Soybean Biotechnology (M.L., G.S.), and Computer Science Department (T.J., D.X.), University of Missouri, Columbia, Missouri 65211; and Samuel Roberts Noble Foundation, Ardmore, Oklahoma 73401 (V.A.B., M.K.U.)

All eukaryotic organisms have a diversity of transcription factor (TF) gene families, encoding key proteins regulating gene expression. TF families are strongly conserved across eukaryotic organisms, especially plants. The specific function of each of these TF genes is of interest due to their role in controlling plant developmental processes and responses to environmental conditions, including functions of key importance to agronomic performance. In this review, we focus on the role of TF genes in legume species. The review also provides an update on the identification and categorization of TF genes in several eukaryotes, including three partially or completely sequenced legume genomes (soybean [*Glycine max*], *Medicago truncatula*, and *Lotus japonicus*). The role of TF genes in legumes is discussed in an evolutionary context based upon a comprehensive comparison of TF gene distribution and direct experimental data obtained for a significant number of legume TF genes.

TF genes are present in all eukaryotic phyla. They encode regulatory proteins that interact with genomic DNA promoter and enhancer sequences. These interactions facilitate the transcriptional activation or repression of proximal genes and enable cells to respond to changes in their environment (e.g. biotic and abiotic stresses), to regulate the cell cycle, and, in the case of the most complex organisms, to control cell fate. As mentioned by Carroll (2001), the expansion of regulatory protein numbers and interactions, as well as changes to their spatial and temporal expression, is part of the evolutionary process leading to increasingly complex organisms. Therefore, determining the repertoire of TF genes in genomes, the regulation of their expression, and their biochemical properties (i.e. DNA- and protein-binding affinities) is important to the understanding of TF regulatory networks and organism evolution.

The immobile nature of plants represents a major disadvantage compared with animals, which can flee many environmental assaults. The ability to face environmental challenges implies that plants must possess complex regulatory systems to respond appropriately. This sometimes involves changing developmental programs, which is facilitated by the fact that plants maintain active stem cells, called meristems, which can differentiate and develop into various organs depending on environmental and endogenous cues. As regulators of transcription, TFs play important roles in helping plants meet and master environmental challenges. Therefore, it is not surprising that plants have more TF genes than animals (Riechmann et al., 2000; this study).

Most of the extant knowledge of plant TF genes was obtained from studies of the major genetic model in plant biology, *Arabidopsis* (*Arabidopsis thaliana*). However, while *Arabidopsis* is a useful model for many developmental and other processes common to all higher plants, it lacks certain traits that are of immense value to agriculture, such as the ability to form nitrogen-fixing symbioses with rhizobia and soil nutrient-scavenging symbioses with mycorrhizal fungi. Legumes, on the other hand, are able to establish such beneficial symbioses and, as a result, have been mainstays for sustainable agriculture for thousands of years. The legume family includes important food plants such as common bean (*Phaseolus vulgaris*), soybean, and pea (*Pisum sativum*) and important forage species such as alfalfa (*Medicago sativa*) and clover (*Trifolium* spp.). There is also growing interest in the use of legumes as a source of biomass for biofuel production. Although mycorrhizal symbioses are widespread among plant families, occurring in approximately 90% of all species, symbiotic nitrogen fixation (SNF) is restricted to legumes and a few nonlegume families. This makes legumes special, but just how SNF evolved in legumes remains largely unknown. Answers to this question may emerge from comparative analysis of the genomes of legumes and nonlegumes.

Genome sequencing of three legume species, *L. japonicus* (<http://www.kazusa.or.jp/lotus>), soybean (<http://www.phytozome.net/soybean>), and *M. truncatula* (<http://www.medicago.org/genome>), is nearing completion, and the genome sequences of several nonlegumes, including *Arabidopsis* (*Arabidopsis* Ge-

¹ This work was supported by the National Science Foundation Plant Genome Program (grant no. DBI-0421620).

* Corresponding author; e-mail staceyg@missouri.edu.

The author responsible for distribution of materials integral to the findings presented in this article in accordance with the policy described in the Instructions for Authors (www.plantphysiol.org) is: Gary Stacey (staceyg@missouri.edu).

^[W] The online version of this article contains Web-only data.

www.plantphysiol.org/cgi/doi/10.1104/pp.109.144105

nome Initiative, 2000), *Vitis vinifera* (Jaillon et al., 2007), *Sorghum bicolor* (Paterson et al., 2009), *Physcomitrella patens* (Rensing et al., 2008), *Chlamydomonas reinhardtii* (Merchant et al., 2007), *Oryza sativa* (Yu et al., 2002), and *Populus trichocarpa* (Tuskan et al., 2006), are known (Table I).

Given the central role of TFs in regulating plant gene expression and consequently development, differentiation, and responses to the environment, as well as their key roles in evolution (Ramalingam et al., 2003; Miller et al., 2006; Francia et al., 2007), the set of genes encoding TFs is an obvious place to start looking for evolutionary innovations that define key legume traits, such as SNF. More generally, the study of legume TFs will shed light on their roles in plant processes common to all plant species.

When comparing legumes with other plant phyla, a key question is, "What traits define a legume?" This question was raised several years ago by Doyle and Luckow (2003) and by Zhu et al. (2005). Morphogenesis and life styles of different plant species and more generally of eukaryotes depend heavily on the control of gene expression. Therefore, it can be assumed that the repertoire of TF genes within a plant species, their expression pattern, and their function largely determine the unique aspects of the species. Unfortunately, as discussed below, the functions of only a few legume TF genes are known. Therefore, a tremendous effort is required to characterize the role of TF genes in legumes. The emerging high-throughput genome-based technologies that build upon the sequencing of several legume genomes open new fields of investigation, including a complete identification and classification of legume TF genes.

In this review, we summarize knowledge of legume TFs while updating the classification of legume, nonlegume, and nonplant eukaryotic TF genes. Legume TF function and TF gene expression are also discussed in the light of recently published studies. Altogether, these data will be discussed in an evolutionary context to better understand the impact of TF genes on legume development.

LEGUME TF GENES: WHAT IS KNOWN?

The last comprehensive review of legume TFs predated the completion of any of the legume genome sequencing projects but noted that more than 99% of predicted legume TFs remained to be characterized functionally (Udvardi et al., 2007). In the interim, the genome sequences of three legumes have been completed, or nearly so, enabling both comparative and functional genomics studies, and the roles of new legume TFs have been determined.

Recent studies have elucidated the roles of several TFs in legume development. Initially, several legume TF genes were identified based upon similarity of mutant phenotypes in legumes and *Arabidopsis* and sequence homology between legume genes and the *Arabidopsis* genes known to confer specific mutant phenotypes. Legume TF genes discovered in this way include those involved in the control of floral meristems, such as *LjFLO*, *PsFLO/LFY*, *PsPEAM4*, and *MtPIM* (Hofer et al., 1997; Berbel et al., 2001; Taylor et al., 2002; Dong et al., 2005; Benlloch et al., 2006). In the same way, three *L. japonicus* MYB TFs with homology to *Arabidopsis* *TRANSPARENT TESTA2* (*AtTT2*) were shown to regulate proanthocyanidin biosynthesis after their transient expression in *Arabidopsis* leaves (Yoshida et al., 2008). More recently, the role of legume TF genes was investigated using reverse-genetic approaches. For example, expression of the soybean *GmWRKY13* gene in *Arabidopsis* positively affected lateral root formation (Zhou et al., 2008). RNA interference knockdown of *MtSERF1*, which encodes an AP2-EREBP TF, altered the production of somatic embryos (Mantiri et al., 2008). Legume TF genes were also recently implicated in responses to environmental challenges. For instance, after identifying TF genes regulated in response to phosphate deprivation in common bean (Hernández et al., 2007), Valdés-López et al. (2008) characterized the role of one MYB TF, *PvPHR1*, as a key protein in phosphorus uptake. Similarly, after quantifying the expression levels of

Table I. Classification of main plant species based on their class, order, family, and genus

The genome size, ploidy level of each genome, number of predicted protein-coding genes, and number of TF genes identified are also indicated.

Plant Type	Organism	Organism Genome Size	Chromosome No. and Ploidy	Predicted Total Gene No.	Predicted TF Gene No.	Predicted TF Gene Percentage
Alga	<i>Chlamydomonas reinhardtii</i>	120 Mb	<i>n</i> = 17 (diploid)	16,709	349	2.09
Moss	<i>Physcomitrella patens</i>	511 Mb	<i>n</i> = 27 (diploid)	35,938	1,316	3.66
Monocotyledon	<i>Oryza sativa</i>	430 Mb	<i>n</i> = 12 (diploid)	67,393	4,432	6.58
Monocotyledon	<i>Zea mays</i>	2,400 Mb	<i>n</i> = 10 (diploid)	125,435	5,383	4.29
Monocotyledon	<i>Sorghum bicolor</i>	730 Mb	<i>n</i> = 10 (tetraploid)	36,338	2,464	6.78
Dicotyledon, rosid, Malvidae	<i>Arabidopsis thaliana</i>	115 Mb	<i>n</i> = 5 (diploid)	32,825	2,269	6.91
Dicotyledon, rosid, Fabidae, Fabales	<i>Medicago truncatula</i>	500–550 Mb	<i>n</i> = 8 (diploid)	38,835	1,473	3.79
Dicotyledon, rosid, Fabidae, Fabales	<i>Lotus japonicus</i>	450 Mb	<i>n</i> = 6 (diploid)	42,395	1,637	3.86
Dicotyledon, rosid, Fabidae, Fabales	<i>Glycine max</i>	1,115 Mb	<i>n</i> = 20 (tetraploid)	66,153	5,557	8.40
Dicotyledon, rosid, Fabidae, Malpighiales	<i>Ricinus communis</i>	425 Mb	<i>n</i> = 10 (diploid)	31,221	1,543	4.94
Dicotyledon, rosid, Fabidae, Malpighiales	<i>Populus trichocarpa</i>	550 Mb	<i>n</i> = 19 (diploid)	45,555	2,758	6.05
Dicotyledon, rosid, Vitaceae	<i>Vitis vinifera</i>	504.6 Mb	<i>n</i> = 19 (diploid)	30,434	1,675	5.50

soybean *Dof* genes in various organs, including flowers and pods, the overexpression of *GmDof4* and *GmDof11* TF genes in Arabidopsis was shown to increase seed lipid content (Wang et al., 2007).

A large number of studies also highlighted the role of legume TF genes in plant responses to abiotic stresses. For example, a common method for examining gene function is to overexpress the gene of interest from a strong, constitutive promoter (e.g. cauliflower mosaic virus 35S) and then to gauge the response of the resulting transgenic plants to a variety of treatments (e.g. abiotic stress). For example, the overexpression of two *TFIIIA-related* TF genes (*MtZPT2-1* and *MtZPT2-2*) in *M. truncatula* led to an increase in the size of the plant root system under salt stress (de Lorenzo et al., 2007). *GmWRKY54* and *GmDREB2* increased tolerance to both salt and drought when overexpressed in Arabidopsis (Chen et al., 2007; Zhou et al., 2008). In the same study, Zhou et al. (2008) also implicated two other *WRKY* TF genes, *GmWRKY21* and *GmWRKY13*, in tolerance to cold, salt, and mannitol stresses by overexpressing these genes in Arabidopsis plants. A similar approach established a role for *GmDREB3* in Arabidopsis tolerance to cold, drought, and high-salt stresses (Chen et al., 2009). Zhu et al. (2006) observed a higher tolerance of soybean plants to high temperature following overexpression of the *GmHSFA1* TF gene. Interestingly, Zhang et al. (2008) identified two *GmAP2-EREBP* TF genes involved not only in abiotic stress response but also in plant-pathogen defense mechanisms. Overexpression of *GmAP2-EREBPs* in tobacco (*Nicotiana tabacum*) enhanced plant resistance to drought, salt stress, and pathogen infection. This study suggested that some TF genes are central to the general plant stress response. With regard to pathogen response, Park et al. (2007) reported that *GmZF-HD1* and *GmZF-HD2* were induced in response to *Pseudomonas syringae* infection.

LEGUME TF GENES AND NODULATION

In addition to the activation of plant defense systems through the activation of TF genes, legume TFs are also involved in the control of mutualistic interactions between plant root and soil microorganisms. For example, nodulation involves the interaction between root and soil bacteria leading to SNF. This complex interaction is mainly restricted to legumes and, for this reason, makes legumes special. The infection of plant roots by symbiotic bacteria begins by the invasion of root hair cells by the symbiont through the newly formed infection thread. This infection is dependent on several genes. Genes that are specifically expressed during nodulation are termed nodulins. Several years ago, protein factors that bound to the AT-rich promoter sequences of nodulin genes were identified (Jensen et al., 1988; Metz et al., 1988; Forde et al., 1990; Laursen et al., 1994; Hansen et al., 1999). More recently, TF genes specifically involved in the rhizobial infection process have been identified. Among them,

the *M. truncatula* *NSP1* and *NSP2* genes, which encode two GRAS TFs, are essential to root hair infection. The root hairs of *Mtnsp1* and *Mtnsp2* mutants are not infected by the symbiont (Catoira et al., 2000; Oldroyd and Long, 2003). More recently, based on a mutant screen for plants lacking infection threads upon inoculation with rhizobia, the *L. japonicus* *NSP1* and *NSP2* genes were identified (Heckmann et al., 2006).

The first TF shown to have an essential role in nodulation was the *L. japonicus* *NIN* gene (Schauser et al., 1999). Mutations in this gene abolish both infection and the induction of nodule primordia. However, *NIN* appears to act downstream of the initial steps in symbiont recognition by the plant (Oldroyd and Downie, 2008). By screening fast-neutron and *Tnt1* transposon-tagged mutagenized populations, Marsh et al. (2007) characterized the *M. truncatula* ortholog of *LjNIN* (Catoira et al., 2001). The *PsSym35* locus defines the pea *NIN* ortholog (Borisov et al., 2003). The likely soybean ortholog of *NIN* is clearly evident in the soybean genomic sequence, and this genomic region shows significant microsynteny to the *NIN*-encoding regions of *M. truncatula* and *L. japonicus* (M. Libault, X.C. Zhang, and G. Stacey, unpublished data).

Additional TF genes critical to the nodulation process were also identified by direct screening for nodulation-defective mutants. For example, this approach led to the identification of the *L. japonicus* *ASTRAY* gene encoding a bZIP TF (Nishimura et al., 2002). *L. japonicus* *astray* mutants exhibit a hypernodulation phenotype (i.e. increased nodule numbers). Similarly, Middleton et al. (2007) identified the *MtERN* gene, encoding an AP2-EREBP TF, as necessary for invasion of plant cells by symbiotic bacteria. Recently, an additional AP2-EREBP TF gene was identified as a positive regulator of nodulation in *L. japonicus* (Asamizu et al., 2008). Other strategies also allowed the identification of important TF genes involved in nodulation. For example, based on its interaction with *LjSymRK* (a receptor kinase critical for rhizobial infection; Stracke et al., 2002), Zhu et al. (2008) identified *LjSIP1*, a protein previously described for its role during nodulation that encodes an ARID TF. The authors proposed that *LjSIP1* controlled the expression of the Nod factor-induced *NIN* gene. El Yahyaoui et al. (2004) utilized DNA microarray analysis to identify several genes regulated during *M. truncatula* nodulation, including an AP2-EREBP TF gene named *MtEFD*. Based on the phenotype of an *edf1* null mutant, as well as plants either silenced for *EFD* expression or overexpressing the gene, Vernié et al. (2008) defined *MtEFD* as an important regulator controlling *Sinorhizobium meliloti* infection of *M. truncatula*.

In summary, recent analyses highlight the roles of a diverse group of TF genes in a wide variety of legume biological processes. Overexpression of legume TF genes in nonlegume plants often produces phenotypes consistent with strong conservation of TF function among higher plant species. Despite the redundancy of TF gene families across the plant kingdom, it is

interesting that legumes nodulate while most other plant species do not. Although specific TF genes have been implicated in nodule development, these belong to families common to nonlegumes. This suggests that neofunctionalization of TF genes was important in the evolution of SNF, rather than the invention of novel TF genes/families.

EVOLUTION OF TF GENES AND FAMILIES IN THE EUKARYOTIC KINGDOM

To better understand the evolution of TFs, comparative studies of TF gene families of several eukaryote organisms, including *Arabidopsis*, have been performed (Riechmann et al., 2000). However, limited genome sequences from plant species and underrepresentation of TFs in cDNA and EST databases have hampered such studies until recently. Now, however, genome sequences of a large diversity of organisms, including several plant species, have been released or will be soon (Table I). The release of genomic sequences combined with the use of improved gene prediction software (e.g. FGENESH) and protein function prediction software (e.g. Pfam [Finn et al., 2008], InterProScan [Hunter et al., 2009], and UniProt [UniProt Consortium, 2009]) allows a more complete identification of the gene content in each genome, including TF genes. Furthermore, prediction tools are now well supported by the detection of gene transcripts via ultra-high-throughput sequencing methods. Based on such characteristics, it is now possible to characterize the entire TF gene population of a few plant species, at least as far as known TF families are concerned. Highlighting similarities and differences in TF gene populations among eukaryotes, and more specifically among plants, may help to answer the question, "What makes a legume a legume?" (Doyle and Luckow, 2003).

We investigated TF gene evolution in eukaryotic phyla based on their family membership. By mining protein sequence databases of 19 major eukaryotes, we identified signature Pfam domains conserved in the different TF families (e value $< e^{-3}$; Table II) and categorized them according to their family membership (Fig. 1). The percentage of identified TF genes compared with the total number of protein-encoding genes analyzed fluctuates between 2% and 9% among the 19 organisms studied. Not surprisingly, the smallest populations of TF genes were identified in the most primitive organisms (e.g. *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, and *C. reinhardtii*, where TF genes represent 2%–4% of the genes annotated). In higher eukaryotes, the higher complexity of form and function likely dictates the need for an increased number of TF genes (7.36% in *Drosophila melanogaster*, 9.12% in *Rattus norvegicus*, 7.65% in *Mus musculus*, and 8.15% in *Homo sapiens*; an average of 5.7% of plant genes are TF genes; Table II).

Considering the distribution of genes among the 94 TF families identified, 22 families are specific to plants

and 20 families are specific to animals (Table II). Eight families are found exclusively in yeast, while one TF family is represented in *D. melanogaster* only. Forty-three families were common to all phyla represented in this study. No legume-specific TF gene family was found, which may reflect the paucity of knowledge about legume TFs but more likely indicates conservation of TF families among plants. Consistent with this notion, the distribution of legume genes among the various TF families is similar to that of other plant species. Overall, based on this census of genes in the various TF families, there does not appear to be any enrichment of known TF families in legumes.

CAN THE KNOWLEDGE ACQUIRED ON TF GENES IN ONE PLANT BE USED IN LEGUMES?

The absence of significant differences in TF gene distribution across TF families between legume and nonlegume plants suggests that legume-specific traits are likely dependent on TF gene expression patterns and TF protein function. So far, most of the plant TFs characterized belong to *Arabidopsis*. Therefore, an attractive strategy is to apply the knowledge established in *Arabidopsis* to legume TFs (Hofer et al., 1997; Berbel et al., 2001; Taylor et al., 2002; Dong et al., 2005; Benlloch et al., 2006). The expression of legume TF genes in *Arabidopsis* or tobacco plants also supports a conservation of functions for TF homologs across plant species (e.g. in plant development [*GmWRKY13*; Zhou et al., 2008], metabolite biosynthesis [*LjMYB* TFs homologous to *AtTT2*; Yoshida et al., 2008], plant resistance to pathogens [*GmAP2-EREBP* TFs; Zhang et al. 2008], and plant resistance to abiotic stresses [*GmWRKY* TFs; Zhou et al. 2008; *GmDREB3*; Chen et al., 2009]). However, complete functional redundancy of TF proteins between plants is not always found. For example, the *AtPAP1* gene encoding a *MYB* TF gene controls anthocyanin synthesis in *Arabidopsis* but it does not activate this pathway when expressed in *M. truncatula* (Peel et al., 2009).

Based upon these few examples, TF function appears highly but not absolutely conserved across plant species. This conclusion is also supported by the strong conservation of TF signature protein domains and the tertiary structure of TFs (Dr. Jianlin Cheng, personal communication). However, one difficulty in applying this strategy is the accurate identification of true orthologs between plant species. As described below, due to the evolutionary distance existing between plants, syntenic relationships are difficult to define. For example, in the rosoid clade, *Arabidopsis* and legumes fall in the Malvidae (Eurosidae II) and the Fabidae (Eurosidae I) subclades, respectively (Table I). These two subclades diverged approximately 115 to 93 millions years ago (Mya). According to Wang et al. (2009), a diversification of rosoids occurred suddenly (in less than 15 millions years) shortly after the divergence between Fabidae and Malvidae. Such diversifi-

Table II. Number of TF genes in 19 of the main eukaryotic organisms

TF proteins were identified for 19 model eukaryotes: *Saccharomyces cerevisiae* (Sc), *Schizosaccharomyces pombe* (Sp), *Caenorhabditis elegans* (Ce), *Drosophila melanogaster* (Dm), *Mus musculus* (Mm), *Rattus norvegicus* (Rn), *Homo sapiens* (Hs), *Chlamydomonas reinhardtii* (Cr), *Physcomitrella patens* (Pp), *Oryza sativa* (Os), *Zea mays* (Zm), *Sorghum bicolor* (Sb), *Arabidopsis thaliana* (At), *Ricinus communis* (Rc), *Populus trichocarpa* (Pt), *Medicago truncatula* (Mt), *Lotus japonicus* (Lj), *Glycine max* (Gm), and *Vitis vinifera* (Vv). The TF genes were distributed across 94 families based on the identification of the TF family signature protein domain (Kakar et al. [2008] and the DBD database [http://dbd.mrc-lmb.cam.ac.uk/DBD/index.cgi?Home; Wilson et al., 2008] were used as references for this classification). The color code used is based on phylogenetic relationships: yeast (black), animal (blue; nematode, arthropod, and mammal in dark blue, blue, and light blue, respectively), algae (dark green), moss (green), monocotyledon (light green), dicotyledon (warm colors; Malvidae, Fabidae, and Vitaceae in red, yellow/orange (Fabales/Malpighiales), and pink, respectively).

Transcription Factor families	Sc	Sp	Ce	Dm	Mm	Rn	Hs	Cr	Pp	Os	Zm	Sb	At	Mt	Lj	Gm	Rc	Pt	Vv
ABI3/VP1	0	0	0	0	0	0	0	2	37	70	97	58	71	85	34	78	35	103	21
AF-4	0	0	0	1	6	3	5	0	0	0	0	0	0	0	0	0	0	0	0
AFT	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
AP-2 TF	0	0	4	2	6	4	7	0	0	0	0	0	0	0	0	0	0	0	0
AP2-EREBP	0	0	0	0	0	0	0	15	154	199	338	161	146	109	159	381	114	211	127
APSES	4	4	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
ARID	2	2	4	8	14	9	22	3	6	8	19	5	10	12	9	22	8	12	9
AUX-IAA-ARF	0	0	0	0	0	0	0	0	15	99	191	59	51	24	36	129	37	70	43
BED-type (Zn)	1	0	8	6	2	2	7	0	0	138	176	28	5	15	8	6	0	30	5
BESS	0	0	0	21	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
bHLH	6	4	48	67	113	77	138	8	91	174	309	151	172	71	84	393	96	145	97
BRCT	0	0	0	0	0	1	10	0	0	0	0	0	0	0	0	0	0	0	0
BROMODOMAIN	0	0	8	5	13	14	43	1	0	1	68	21	29	13	13	57	2	1	1
BTB/POZ	1	3	100	131	142	51	109	34	34	142	103	125	98	22	31	145	27	39	28
BZIP	12	6	39	48	56	38	78	8	38	133	245	103	78	44	38	176	49	83	45
BZIP-MAF	0	0	1	2	11	6	10	0	0	0	0	0	0	0	0	0	0	0	0
C2C2 (Zn) CO-like	0	0	11	14	77	33	125	1	26	42	66	32	34	15	21	72	22	38	24
C2HC (Zn)	0	0	1	2	5	3	4	0	0	0	0	0	0	0	0	0	0	0	0
C3H-type1(Zn)	5	7	32	20	54	24	61	14	33	91	193	55	69	41	50	147	43	72	46
C4 (Zn)	0	0	327	50	64	47	112	0	0	0	0	0	0	0	0	0	0	0	0
C2C2 (Zn) Dof	0	0	0	0	0	0	0	1	21	37	54	29	36	21	22	82	23	42	26
C2C2 (Zn) GATA	10	5	16	9	12	9	15	12	15	35	63	33	29	29	16	62	19	38	20
C2C2 (Zn) YABBY	0	0	0	0	0	0	0	0	0	15	41	8	6	6	4	18	6	13	7
C ₂ H ₂ (Zn)	36	18	185	429	843	300	1000	6	42	118	162	88	173	52	56	395	59	73	57
CAMTA	0	0	2	4	2	1	2	0	1	7	14	7	2	6	4	15	5	7	4
CCAAT	5	7	9	11	11	8	9	6	20	64	99	38	38	22	25	106	31	51	27
CCHC (Zn)	10	7	14	17	31	14	24	5	49	1448	347	206	66	141	105	144	31	99	121
Churchill	0	0	0	0	1	1	1	0	0	0	0	0	0	0	0	0	0	0	0
Copper fist DNA binding protein	3	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
CP2	0	0	1	9	8	4	12	0	0	0	0	0	0	0	0	0	0	0	0
CSD	0	0	6	6	16	7	10	1	3	2	6	1	4	0	2	7	4	5	3
DDT	1	0	4	5	2	1	2	0	0	8	18	6	6	2	4	15	3	3	5
DHHC (Zn)	7	5	19	33	27	13	33	10	21	51	86	30	24	18	15	47	23	37	25
DM TF	0	0	13	6	8	5	8	0	0	0	0	0	0	0	0	0	0	0	0
E2F/DP	0	0	7	6	11	9	12	3	11	12	29	10	8	6	7	14	6	10	6
EIL	0	0	0	0	0	0	0	0	2	11	13	7	3	7	6	13	4	6	4
Ets	0	0	13	18	36	19	40	0	0	0	0	0	0	0	0	0	0	0	0
Fez1	0	0	0	0	5	2	4	0	0	0	0	0	0	0	0	0	0	0	0
FHA	15	9	39	56	83	49	107	10	15	22	28	14	5	9	10	33	15	19	13
Fungal specific TF	25	22	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
GCM	0	0	0	3	2	2	0	0	0	0	0	0	0	0	0	0	0	0	0
GRAS	0	0	0	0	0	0	0	0	39	64	105	74	33	36	43	130	45	98	45
GRF	1	1	1	1	8	5	5	4	1	51	12	29	2	14	15	6	8	7	3
HD-ZIP	0	0	0	0	0	0	0	0	2	4	1	1	2	1	7	3	5	0	0
HMG	7	7	22	43	85	38	101	7	9	17	33	14	12	2	9	31	9	12	7
Homeodomain/HOMEBOX	7	2	108	173	317	133	290	0	37	121	211	74	112	46	55	318	59	104	69
HSF	5	2	1	4	6	4	11	2	8	42	58	25	24	16	18	11	19	31	19
HTH11	0	0	0	0	0	0	0	2	0	0	0	0	0	0	0	0	0	0	0
HTH-ARAC	0	0	0	0	0	0	0	20	0	0	0	0	0	1	13	26	6	0	0
HTH-FIS	0	0	0	0	0	0	0	0	0	0	0	0	14	0	0	7	2	5	0
HTHpsq	0	0	1	10	2	0	1	0	3	0	0	0	0	0	0	0	0	0	0
IRF	0	0	0	0	9	7	17	0	0	0	0	0	0	0	0	0	0	0	0
JMONJI	3	5	15	12	29	7	27	5	12	17	46	21	21	13	12	77	15	19	17
KRAB	0	0	0	0	77	3	44	0	0	0	0	0	0	0	0	2	0	0	0
LFY	0	0	0	0	0	0	0	0	2	1	4	1	1	1	2	4	1	1	1
LIM	4	4	48	89	93	48	92	0	11	16	59	9	13	9	12	42	9	20	14
MADS	4	3	2	7	8	2	5	0	22	95	147	82	109	60	83	212	39	111	60
MAT Alpha1	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
MBF1	0	0	0	0	0	0	0	0	1	1	7	0	3	1	3	4	11	3	0
MYB	14	12	21	33	40	17	46	29	147	298	564	262	303	171	191	791	189	378	242
NAC	3	2	3	8	12	2	12	2	40	162	252	126	114	64	101	208	94	179	83
NDT80_PhoG	1	0	4	0	2	1	4	0	0	0	0	0	0	0	0	0	0	0	0
Opi1 TF	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
P53	0	0	0	2	3	3	3	0	0	0	0	0	0	0	0	0	0	0	0
p53-like	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	5	0	0	0
PAX	0	0	7	9	5	2	15	0	0	0	0	0	0	0	0	0	0	0	0
PHD	10	17	23	45	60	30	77	28	68	87	202	74	55	45	47	222	58	90	58
PLATZ	0	0	0	0	0	0	0	3	13	22	22	17	3	10	9	25	11	20	10
R3H	2	3	2	10	13	7	11	0	3	3	5	2	7	1	0	14	3	3	2
RFX DNA binding	1	1	4	3	10	3	11	0	0	0	0	0	0	0	0	0	0	0	0
RHD	0	0	0	13	14	5	22	0	0	0	0	0	0	0	0	0	0	0	0
Runt1	0	0	1	6	3	2	7	0	0	0	0	0	0	0	0	0	0	0	0
RWP-RK	0	0	0	0	0	0	0	15	9	14	37	13	14	5	12	23	10	18	8
S1Fa-like	0	0	0	0	0	0	0	1	2	3	2	0	3	1	4	1	2	2	2
SAND	0	0	4	2	4	2	6	0	0	0	0	0	0	0	0	0	0	0	0
SBP	0	0	0	0	0	0	0	23	14	29	70	18	16	11	15	48	15	29	18
SNF2	14	14	23	26	32	22	39	15	31	42	85	34	33	17	23	69	25	41	28
SSB	1	1	1	2	2	1	1	0	4	9	10	4	3	4	7	6	8	4	4
STE TF	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
TAZ	0	0	9	1	2	1	3	3	5	11	17	5	10	5	2	1	5	7	4
T-box	0	0	21	11	22	13	27	0	0	0	0	0	0	0	0	0	0	0	0
TCP	0	0	0	0	0	0	0	6	24	59	26	6	11	21	65	21	34	19	0
TEA	1	0	6	7	5	1	6	0	0	0	0	0	0	0	0	0	0	0	0
TPR	22	25	48	75	124	76	154	63	98	163	258	112	65	52	86	318	100	142	98
TUB	0	0	2	3	8	3	6	2	6	26	43	20	2	12	8	24	7	11	8
U1-type (Zn)	1	2	2	5	10	5	12	2	4	3	9	4	7	2	2	6	2	3	1




Figure 1. Comparison of TF gene distribution based on family membership across 19 different eukaryotic species. For each species, the representation of each TF family in the total pool of TF genes identified was expressed as a percentage. The color scale indicates the degree of correlation (red, low correlation; black, strong correlation). The heat map was generated using the Euclidean distances between each organism using the MultiExperiment Viewer (<http://www.tm4.org/mev.html>).

cation might explain the vast heterogeneity in rosid habitats, morphology, and development. This evolutionary distance between crop plants and Arabidopsis is also highlighted by the strong changes in the size and organization of their genomes (Table I), such as the ancient genome duplication in legumes occurring around 45 to 55 Mya (Pfeil et al., 2005; Cannon et al., 2006) followed by a more recent duplication in soybean around 10 to 15 Mya (Schlueter et al., 2004, 2007). Despite the evolutionary distances existing between legumes and Arabidopsis, microsynteny is still found in some regions (Grant et al., 2000; Yan et al., 2003). For example, 14% of analyzed contig groups showed microsynteny between Arabidopsis and soybean (Zhu et al., 2003; Yan et al., 2004; Mudge et al., 2005). Likewise, a comparison between soybean, *Medicago*, and Arabidopsis identified two blocks with strong microsynteny (Shultz et al., 2007). Conversely, macrosynteny between Arabidopsis and legumes is difficult to identify due to genome duplication, recombination, and gene loss (Kevei et al., 2005; Schlueter et al., 2008).

Extensive macrosyntentic relationships exist between legume species (Choi et al., 2004; Cannon et al., 2006; for review, see Young and Udvardi, 2009). The establishment of genome-wide colinearity

between legumes will be an important advantage for transferring information to soybean, an economically important plant, from the knowledge established in the legume models *M. truncatula* and *L. japonicus*. The fact that all four major legume genetic models (*L. japonicus*, *M. truncatula*, soybean, and common bean) fall in the Papilionoideae group, one of the three groups of legumes (i.e. legumes are divided into three groups named Caesalpinioideae, Mimosoideae, and Papilionoideae), supports the strong macrosynteny found in the available legume genome sequences. The Papilionoideae group diverged from the two other legume clades approximately 50 Mya, while the *Lotus* and *Medicago* lineages diverged from one another around 40 Mya (Wojciechowski, 2003). This recent divergence among major legumes makes it easier to identify orthologs by direct genome comparison. For example, this approach allowed the identification of key TF gene orthologs involved in the nodulation process and in floral meristem development (e.g. *LjNIN* and *PsSym35*; *LjNSP1* and *MtNSP1*; *LjNSP2* and *MtNSP2*; and *LjFLO* and *PsFLO* [Hofer et al., 1997; Schauser et al., 1999; Catoira et al., 2000; Borisov et al., 2003; Oldroyd and Long, 2003; Dong et al., 2005; Kalo et al., 2005; Smit et al., 2005; Heckmann et al., 2006]).

SOME LEGUME TF GENES ARE INDUCED SPECIFICALLY DURING NODULATION

As described above, legume TF distribution across families and their basic functions appear to be conserved compared with other plant families. Consequently, special legume traits may derive from unique TF gene expression patterns. During the last decade, the availability of cDNA and oligonucleotide arrays allowed the quantification of gene expression patterns in a large number of organisms in different tissues and under differing environmental conditions. However, several studies clearly demonstrated the limit of this technology to accurately quantify the expression of low-abundance transcripts, such as TF genes (Czechowski et al., 2004; Libault et al., 2007). To better characterize the expression of TF genes, large-scale quantitative reverse transcription-PCR platforms were developed to quantify TF gene expression in Arabidopsis, rice, *M. truncatula*, soybean, and *L. japonicus* (Czechowski et al., 2004; Caldana et al., 2007; Kakar et al., 2008; Libault et al., 2009; O. Montanari and M.K. Udvardi, unpublished data). Several studies used these resources to quantify TF gene expression in different tissues and in response to different treatments (McGrath et al., 2005; Libault et al., 2007; Gruber et al., 2009; Libault et al., 2009). Complementary to the use of large-scale quantitative reverse transcription-PCR platforms, ultra-high-throughput sequencing technologies, such as the 454 Life Sciences (Margulies et al., 2005) and Illumina Solexa (Bennett et al., 2005) platforms, allow an accurate quantification of low-abundance transcripts. These resources allow the char-

acterization of the TF gene transcriptome. However, the use of such technology is fully informative only if the genome of the organism of interest is sequenced and accurately annotated.

To assess the potential of divergent gene expression patterns as a potential reason for legume-specific attributes, we investigated the expression of members of the *NIN*-like gene family. This family was selected based upon the involvement of some of its members in root hair infection and nodule development (Schauer et al., 1999; Catoira et al., 2001; Borisov et al., 2003; Marsh et al., 2007). Two recent studies place *NIN* gene

function just downstream to *NSP1* and *NSP2* GRAS in the signaling cascade regulating legume nodulation. For example, *LjNIN* gene expression is under the control of *LjNSP2* (Murakami et al., 2006). Moreover, *MtNSP1* and *MtNSP2* interact with the *MtNIN* gene promoter to regulate its expression (Hirsch et al., 2009). In the same study, Hirsch et al. (2009) showed that the molecular interaction of *MtNSP1* and *MtNSP2* was also required to activate *MtENOD11* gene expression during nodulation. In addition, Gonzalez-Rizzo et al. (2006) showed that *MtNIN* gene expression was under the control of cytokinin, a hormone previously




Figure 2. Arabidopsis, soybean, *L. japonicus*, and *M. truncatula* RWP-RK gene expression in various plant tissues. Relative expression levels of Arabidopsis (A), soybean (B), *L. japonicus* (C), and *M. truncatula* (D) RWP-RK genes (y axis) are reported for five different tissues (nodule [blue], root [purple], leaf [yellow], flower [green], and pod [red]). For each plant species, the identity of the RWP-RK genes is reported on the x axis. Details are available in Supplemental Table S1. Genes encoding proteins carrying a PB1 domain (*NIN*-like genes) are highlighted by arrows. *LjNIN* (Schauer et al., 1999) and *MtNIN* (Marsh et al., 2007) involved in nodulation are highlighted. *, *GmRWP-RK1* was also strongly expressed in soybean apical meristem and root tip; consequently, *GmRWP-RK1* is not considered a nodule-specific RWP-RK.

described to influence legume nodulation (Murray et al., 2007; Frugier et al., 2008).

Based on a bioinformatic analysis and similarly to Schauser et al. (2005), we identified two distinctive domains in the plant NIN-like amino acid sequences: an RWP-RK domain and a Phox and Bem1p (PB1) domain (the PB1 domain is found in a large number of eukaryotic cytoplasmic signaling proteins; Sumimoto et al., 2007; Fig. 2). In addition, TF proteins carrying only the RWP-RK domain were also identified in different plant genomes. The *RWP-RK* gene TF family is a small one, with 14 members in Arabidopsis (0.62% of all identified TF genes), five in *M. truncatula* (0.34%), 12 in *L. japonicus* (0.73%), and 23 in soybean (0.41%; Fig. 2). Among them, we identified nine, three, four, and 14 *NIN-like* genes in Arabidopsis, *M. truncatula*, *L. japonicus*, and soybean, respectively. The expression of 11 *RWP-RK* genes Arabidopsis, three in *M. truncatula*, 12 in *L. japonicus*, and 23 in soybean was quantified in a large variety of tissues using Affymetrix arrays or ultra-high-throughput sequencing (Schmid et al., 2005; Benedito et al., 2008; Høglund et al., 2009; M. Libault, A. Farmer, G.D. May, and G. Stacey, unpublished data; Fig. 2). *RWP-RK* genes were expressed at least 10-fold higher in nodules than in other organs analyzed in the case of one gene in *M. truncatula*, two in *L. japonicus*, and seven in soybean (Fig. 2). All of these legume *RWP-RK* genes expressed specifically in nodules share the PB1 domain and, consequently, are *NIN-like* genes. Interestingly, *M. truncatula* and soybean nodule-specific *NIN-like* genes are not expressed (or expressed at very low levels) in other tissues, indicating specialization of these genes for nodulation and SNF. Because the proportion of *NIN-like* genes in the Arabidopsis and soybean genomes is similar, this analysis supports the hypothesis that legumes coopted a subset of pre-existing *NIN-like* genes for use in symbiotic establishment by modifying their expression patterns.

FUTURE DIRECTIONS IN LEGUME TF RESEARCH

An analysis of TF genes among the sequenced plant genomes does not reveal a legume-specific family, nor does it identify a specific TF family that appears to have been preferentially expanded in legumes. However, legumes have clearly succeeded in developing specific traits by diverting some TF genes for a more specialized function. In the case of *NIN-like* TFs, this likely involved changes in the sequences of promoters that allowed activation of gene expression during nodule development.

What is now needed is a clearer picture of the legume TF gene transcriptome, interactome (protein-protein interaction), and elucidation of the regulon controlled by each TF. The legume TF transcriptome should be elucidated soon with the emergence of ultra-high-throughput sequencing approaches, making the establishment of a TF interactome the next important challenge. Such an interactome will encompass TF-TF

interactions as well as TF-DNA interactions. Identifying TF-TF interactions will highlight the complexity of legume gene regulation. Studies focusing on bacteria and yeast cells have already established TF-TF interaction networks, giving the first insights into the complexity of these systems (Babu et al., 2004; Luscombe et al., 2004; Ye et al., 2009).

The in vivo identification of cis-regulatory regions of TF genes is now possible by combining chromatin immunoprecipitation (ChIP) methods with the use of tiling arrays developed after sequencing of the whole genome (Gregory et al., 2008). The most recent approach utilizes ChIP coupled with ultra-high-throughput sequencing technologies (ChIP-Seq; Barski and Zhao, 2009). These approaches allow the identification of accessible promoter elements that interact with TF proteins with respect to chromatin remodeling. In fact, chromatin rearrangements are an important tool for regulation of gene expression at the epigenetic level by controlling access to TF-binding sites (Barrera and Ren, 2006). Analyzing TF interactome (TF-TF and TF-DNA) data in the context of chromatin structure will provide a clearer picture of legume gene regulation.

CONCLUSION

Legumes are a fascinating family of plants due, in part, to their ability to develop unique organs, called nodules, which harbor nitrogen-fixing rhizobia. Among the genes involved in the nodulation process, several TFs have been characterized. The similar distribution of TF genes in the various known TF families in legumes and nonlegumes, as well as the conserved nature of their basic biochemical functions, cannot readily explain how legume-specific traits such as SNF evolved. Based upon analysis of the *NIN-like* TF family in soybean, *L. japonicus*, *M. truncatula*, and Arabidopsis, we assume that legumes coopted existing TF genes for use in legume-specific processes by modifying their expression patterns. Analysis of synteny between legume and nonlegume plants and subsequent dissection of the expression patterns of orthologous TF genes will help to test this hypothesis. Furthermore, identification of DNA-binding sequences of TF proteins and detailed analysis of TF gene expression will provide a means to understand the impact of TF activity on the legume transcriptome.

Supplemental Data

The following materials are available in the online version of this article.

Supplemental Table S1. Expression levels of Arabidopsis, soybean, *L. japonicus*, and *M. truncatula* *RWP-RK* genes used to create Figure 2.

Received July 1, 2009; accepted August 26, 2009; published September 2, 2009.

LITERATURE CITED

Arabidopsis Genome Initiative (2000) Analysis of the genome sequence of the flowering plant *Arabidopsis thaliana*. *Nature* **408**: 796–815

- Asamizu E, Shimoda Y, Kouchi H, Tabata S, Sato S (2008) A positive regulatory role for *LjERF1* in the nodulation process is revealed by systematic analysis of nodule-associated transcription factors of *Lotus japonicus*. *Plant Physiol* **147**: 2030–2040
- Babu MM, Luscombe NM, Aravind L, Gerstein M, Teichmann SA (2004) Structure and evolution of transcriptional regulatory networks. *Curr Opin Struct Biol* **14**: 283–291
- Barrera LO, Ren B (2006) The transcriptional regulatory code of eukaryotic cells: insights from genome-wide analysis of chromatin organization and transcription factor binding. *Curr Opin Cell Biol* **18**: 291–298
- Barski A, Zhao K (2009) Genomic location analysis by ChIP-Seq. *J Cell Biochem* **107**: 11–18
- Benedito VA, Torres-Jerez I, Murray JD, Andriankaja A, Allen S, Kakar K, Wandrey M, Verdier J, Zuber H, Ott T, et al (2008) A gene expression atlas of the model legume *Medicago truncatula*. *Plant J* **55**: 504–513
- Benlloch R, d'Erforth I, Ferrandiz C, Cosson V, Beltran JP, Canas LA, Kondorosi A, Madueno F, Ratet P (2006) Isolation of *mtpm* proves *Tnt1* a useful reverse genetics tool in *Medicago truncatula* and uncovers new aspects of AP1-like functions in legumes. *Plant Physiol* **142**: 972–983
- Bennett ST, Barnes C, Cox A, Davies L, Brown C (2005) Toward the 1,000 dollars human genome. *Pharmacogenomics* **6**: 373–382
- Berbel A, Navarro C, Ferrandiz C, Canas LA, Madueno F, Beltran JP (2001) Analysis of *PEAM4*, the pea *AP1* functional homologue, supports a model for *AP1-like* genes controlling both floral meristem and floral organ identity in different plant species. *Plant J* **25**: 441–451
- Borisov AY, Madsen LH, Tsyganov VE, Umehara Y, Voroshilova VA, Batagov AO, Sandal N, Mortensen A, Schausler L, Ellis N, et al (2003) The *Sym35* gene required for root nodule development in pea is an ortholog of *Nin* from *Lotus japonicus*. *Plant Physiol* **131**: 1009–1017
- Caldana C, Scheible WR, Mueller-Roeber B, Ruzicic S (2007) A quantitative RT-PCR platform for high-throughput expression profiling of 2500 rice transcription factors. *Plant Methods* **3**: 7
- Cannon SB, Sterck L, Rombauts S, Sato S, Cheung F, Gouzy J, Wang X, Mudge J, Vasdewani J, Schiex T, et al (2006) Legume genome evolution viewed through the *Medicago truncatula* and *Lotus japonicus* genomes. *Proc Natl Acad Sci USA* **103**: 14959–14964
- Carroll SB (2001) Chance and necessity: the evolution of morphological complexity and diversity. *Nature* **409**: 1102–1109
- Catoira R, Galera C, de Billy F, Penmetsa RV, Journet EP, Maillet F, Rosenberg C, Cook D, Gough C, Denarie J (2000) Four genes of *Medicago truncatula* controlling components of a nod factor transduction pathway. *Plant Cell* **12**: 1647–1666
- Catoira R, Timmers AC, Maillet F, Galera C, Penmetsa RV, Cook D, Denarie J, Gough C (2001) The *HCL* gene of *Medicago truncatula* controls Rhizobium-induced root hair curling. *Development* **128**: 1507–1518
- Chen M, Wang QY, Cheng XG, Xu ZS, Li LC, Ye XG, Xia LQ, Ma YZ (2007) GmDREB2, a soybean DRE-binding transcription factor, conferred drought and high-salt tolerance in transgenic plants. *Biochem Biophys Res Commun* **353**: 299–305
- Chen M, Xu Z, Xia L, Li L, Cheng X, Dong J, Wang Q, Ma Y (2009) Cold-induced modulation and functional analyses of the DRE-binding transcription factor gene, *GmDREB3*, in soybean (*Glycine max* L.). *J Exp Bot* **60**: 121–135
- Choi HK, Mun JH, Kim DJ, Zhu H, Baek JM, Mudge J, Roe B, Ellis N, Doyle J, Kiss GB, et al (2004) Estimating genome conservation between crop and model legume species. *Proc Natl Acad Sci USA* **101**: 15289–15294
- Czechowski T, Bari RP, Stitt M, Scheible WR, Udvardi MK (2004) Real-time RT-PCR profiling of over 1400 Arabidopsis transcription factors: unprecedented sensitivity reveals novel root- and shoot-specific genes. *Plant J* **38**: 366–379
- de Lorenzo L, Merchan F, Blanchet S, Megias M, Frugier F, Crespi M, Sousa C (2007) Differential expression of the TFIIIA regulatory pathway in response to salt stress between *Medicago truncatula* genotypes. *Plant Physiol* **145**: 1521–1532
- Dong ZC, Zhao Z, Liu CW, Luo JH, Yang J, Huang WH, Hu XH, Wang TL, Luo D (2005) Floral patterning in *Lotus japonicus*. *Plant Physiol* **137**: 1272–1282
- Doyle JJ, Luckow MA (2003) The rest of the iceberg: legume diversity and evolution in a phylogenetic context. *Plant Physiol* **131**: 900–910
- El Yahyaoui F, Kuster H, Ben Amor B, Hohnjec N, Puhler A, Becker A, Gouzy J, Vernie T, Gough C, Niebel A, et al (2004) Expression profiling in *Medicago truncatula* identifies more than 750 genes differentially expressed during nodulation, including many potential regulators of the symbiotic program. *Plant Physiol* **136**: 3159–3176
- Finn RD, Tate J, Mistry J, Coghill PC, Sammut SJ, Hotz HR, Ceric G, Forslund K, Eddy SR, Sonnhammer EL, et al (2008) The Pfam protein families database. *Nucleic Acids Res* **36**: D281–D288
- Forde BG, Freeman J, Oliver JE, Pineda M (1990) Nuclear factors interact with conserved A/T-rich elements upstream of a nodule-enhanced glutamine synthetase gene from French bean. *Plant Cell* **2**: 925–939
- Francia E, Barabaschi D, Tondelli A, Laido G, Rizza F, Stanca AM, Busconi M, Fogher C, Stockinger EJ, Pecchioni N (2007) Fine mapping of a HvCBF gene cluster at the frost resistance locus Fr-H2 in barley. *Theor Appl Genet* **115**: 1083–1091
- Frugier F, Kosuta S, Murray JD, Crespi M, Szczyglowski K (2008) Cytokinin: secret agent of symbiosis. *Trends Plant Sci* **13**: 115–120
- Gonzalez-Rizzo S, Crespi M, Frugier F (2006) The *Medicago truncatula* CRE1 cytokinin receptor regulates lateral root development and early symbiotic interaction with *Sinorhizobium meliloti*. *Plant Cell* **18**: 2680–2693
- Grant D, Cregan P, Shoemaker RC (2000) Genome organization in dicots: genome duplication in Arabidopsis and synteny between soybean and Arabidopsis. *Proc Natl Acad Sci USA* **97**: 4168–4173
- Gregory BD, Yazaki J, Ecker JR (2008) Utilizing tiling microarrays for whole-genome analysis in plants. *Plant J* **53**: 636–644
- Gruber V, Blanchet S, Diet A, Zahaf O, Boualem A, Kakar K, Alunni B, Udvardi M, Frugier F, Crespi M (2009) Identification of transcription factors involved in root apex responses to salt stress in *Medicago truncatula*. *Mol Genet Genomics* **281**: 55–66
- Hansen AC, Busk H, Marcker A, Marcker KA, Jensen EO (1999) *VsENBP1* regulates the expression of the early nodulin *PsENOD12B*. *Plant Mol Biol* **40**: 495–506
- Heckmann AB, Lombardo F, Miwa H, Perry JA, Bunnewell S, Parniske M, Wang TL, Downie JA (2006) *Lotus japonicus* nodulation requires two GRAS domain regulators, one of which is functionally conserved in a non-legume. *Plant Physiol* **142**: 1739–1750
- Hernandez G, Ramirez M, Valdes-Lopez O, Tesfaye M, Graham MA, Czechowski T, Schlereth A, Wandrey M, Erban A, Cheung F, et al (2007) Phosphorus stress in common bean: root transcript and metabolic responses. *Plant Physiol* **144**: 752–767
- Hirsch S, Kim J, Munoz A, Heckmann AB, Downie JA, Oldroyd GE (2009) GRAS proteins form a DNA binding complex to induce gene expression during nodulation signaling in *Medicago truncatula*. *Plant Cell* **21**: 545–557
- Hofer J, Turner L, Hellens R, Ambrose M, Matthews P, Michael A, Ellis N (1997) *UNIFOLIATA* regulates leaf and flower morphogenesis in pea. *Curr Biol* **7**: 581–587
- Hogslund N, Radutoiu S, Krusell L, Voroshilova V, Hannah MA, Goffard N, Sanchez DH, Lippold F, Ott T, Sato S, et al (2009) Dissection of symbiosis and organ development by integrated transcriptome analysis of *lotus japonicus* mutant and wild-type plants. *PLoS One* **4**: e6556
- Hunter S, Apweiler R, Attwood TK, Bairoch A, Bateman A, Binns D, Bork P, Das U, Daugherty L, Duquenne L, et al (2009) InterPro: the integrative protein signature database. *Nucleic Acids Res* **37**: D211–D215
- Jaillon O, Aury JM, Noel B, Policriti A, Clepet C, Casagrande A, Choise N, Aubourg S, Vitulo N, Jubin C, et al (2007) The grapevine genome sequence suggests ancestral hexaploidization in major angiosperm phyla. *Nature* **449**: 463–467
- Jensen EO, Marcker KA, Schell J, Bruijn FJ (1988) Interaction of a nodule specific, trans-acting factor with distinct DNA elements in the soybean leghaemoglobin *lbc(3)* 5' upstream region. *EMBO J* **7**: 1265–1271
- Kakar K, Wandrey M, Czechowski T, Gaertner T, Scheible WR, Stitt M, Torres-Jerez I, Xiao Y, Redman JC, Wu HC, et al (2008) A community resource for high-throughput quantitative RT-PCR analysis of transcription factor gene expression in *Medicago truncatula*. *Plant Methods* **4**: 18
- Kalo P, Gleason C, Edwards A, Marsh J, Mitra RM, Hirsch S, Jakab J, Sims S, Long SR, Rogers J, et al (2005) Nodulation signaling in legumes requires NSP2, a member of the GRAS family of transcriptional regulators. *Science* **308**: 1786–1789
- Kevei Z, Seres A, Kereszt A, Kalo P, Kiss P, Toth G, Endre G, Kiss GB (2005) Significant microsynteny with new evolutionary highlights is detected between Arabidopsis and legume model plants despite the lack of macrosynteny. *Mol Genet Genomics* **274**: 644–657
- Laursen NB, Larsen K, Knudsen JY, Hoffmann HJ, Poulsen C, Marcker

- KA, Jensen EO (1994) A protein binding AT-rich sequence in the soybean leghemoglobin c3 promoter is a general cis element that requires proximal DNA elements to stimulate transcription. *Plant Cell* 6: 659–668
- Libault M, Joshi T, Takahashi K, Hurley-Sommer A, Puricelli K, Blake S, Xu D, Nguyen HT, Stacey G (2009) Large-scale analysis of putative soybean regulatory gene expression identifies a *Myb* gene involved in soybean nodule development. *Plant Physiol* 151: 1207–1220
- Libault M, Wan J, Czechowski T, Udvardi M, Stacey G (2007) Identification of 118 Arabidopsis transcription factor and 30 ubiquitin-ligase genes responding to chitin, a plant-defense elicitor. *Mol Plant Microbe Interact* 20: 900–911
- Luscombe NM, Babu MM, Yu H, Snyder M, Teichmann SA, Gerstein M (2004) Genomic analysis of regulatory network dynamics reveals large topological changes. *Nature* 431: 308–312
- Mantiri FR, Kurdyukov S, Lohar DP, Sharopova N, Saeed NA, Wang XD, Vandenbosch KA, Rose RJ (2008) The transcription factor *MtSERF1* of the *ERF* subfamily identified by transcriptional profiling is required for somatic embryogenesis induced by auxin plus cytokinin in *Medicago truncatula*. *Plant Physiol* 146: 1622–1636
- Margulies M, Egholm M, Altman WE, Attiya S, Bader JS, Bemben LA, Berka J, Braverman MS, Chen YJ, Chen Z, et al (2005) Genome sequencing in microfabricated high-density picolitre reactors. *Nature* 437: 376–380
- Marsh JE, Rakocevic A, Mitra RM, Brocard L, Sun J, Eschstruth A, Long SR, Schultze M, Ratet P, Oldroyd GE (2007) *Medicago truncatula* *NIN* is essential for rhizobial-independent nodule organogenesis induced by autoactive calcium/calmodulin-dependent protein kinase. *Plant Physiol* 144: 324–335
- McGrath KC, Dombrecht B, Manners JM, Schenk PM, Edgar CI, Maclean DJ, Scheible WR, Udvardi MK, Kazan K (2005) Repressor- and activator-type ethylene response factors functioning in jasmonate signaling and disease resistance identified via a genome-wide screen of Arabidopsis transcription factor gene expression. *Plant Physiol* 139: 949–959
- Merchant SS, Prochnik SE, Vallon O, Harris EH, Karpowicz SJ, Witman GB, Terry A, Salamov A, Fritz-Laylin LK, Marechal-Drouard L, et al (2007) The Chlamydomonas genome reveals the evolution of key animal and plant functions. *Science* 318: 245–250
- Metz BA, Welters P, Hoffmann HJ, Jensen EO, Schell J, de Bruijn FJ (1988) Primary structure and promoter analysis of leghemoglobin genes of the stem-nodulated tropical legume *Sesbania rostrata*: conserved coding sequences, cis-elements and trans-acting factors. *Mol Gen Genet* 214: 181–191
- Middleton PH, Jakab J, Penmetts RV, Starker CG, Doll J, Kalo P, Prabhu R, Marsh JE, Mitra RM, Kereszt A, et al (2007) An ERF transcription factor in *Medicago truncatula* that is essential for Nod factor signal transduction. *Plant Cell* 19: 1221–1234
- Miller AK, Galiba G, Dubcovsky J (2006) A cluster of 11 CBF transcription factors is located at the frost tolerance locus Fr-Am2 in Triticum monococcum. *Mol Genet Genomics* 275: 193–203
- Mudge J, Cannon SB, Kalo P, Oldroyd GE, Roe BA, Town CD, Young ND (2005) Highly syntenic regions in the genomes of soybean, *Medicago truncatula*, and *Arabidopsis thaliana*. *BMC Plant Biol* 5: 15
- Murakami Y, Miwa H, Imaizumi-Anraku H, Kouchi H, Downie JA, Kawaguchi M, Kawasaki S (2006) Positional cloning identifies Lotus japonicus NSP2, a putative transcription factor of the GRAS family, required for *NIN* and *ENOD40* gene expression in nodule initiation. *DNA Res* 13: 255–265
- Murray JD, Karas BJ, Sato S, Tabata S, Amyot L, Szczylowski K (2007) A cytokinin perception mutant colonized by Rhizobium in the absence of nodule organogenesis. *Science* 315: 101–104
- Nishimura R, Ohmori M, Kawaguchi M (2002) The novel symbiotic phenotype of enhanced-nodulating mutant of Lotus japonicus: astray mutant is an early nodulating mutant with wider nodulation zone. *Plant Cell Physiol* 43: 853–859
- Oldroyd GE, Downie JA (2008) Coordinating nodule morphogenesis with rhizobial infection in legumes. *Annu Rev Plant Biol* 59: 519–546
- Oldroyd GE, Long SR (2003) Identification and characterization of nodulation-signaling pathway 2, a gene of *Medicago truncatula* involved in Nod factor signaling. *Plant Physiol* 131: 1027–1032
- Park HC, Kim ML, Lee SM, Bahk JD, Yun DJ, Lim CO, Hong JC, Lee SY, Cho MJ, Chung WS (2007) Pathogen-induced binding of the soybean zinc finger homeodomain proteins GmZF-HD1 and GmZF-HD2 to two repeats of ATTA homeodomain binding site in the calmodulin isoform 4 (GmCaM4) promoter. *Nucleic Acids Res* 35: 3612–3623
- Paterson AH, Bowers JE, Bruggmann R, Dubchak I, Grimwood J, Gundlach H, Haberer G, Hellsten U, Mitros T, Poliakov A, et al (2009) The *Sorghum bicolor* genome and the diversification of grasses. *Nature* 457: 551–556
- Peel GJ, Pang Y, Modolo LV, Dixon RA (2009) The LAP1 MYB transcription factor orchestrates anthocyanidin biosynthesis and glycosylation in *Medicago*. *Plant J* 59: 136–149
- Pfeil BE, Schlueter JA, Shoemaker RC, Doyle JJ (2005) Placing paleopolyploidy in relation to taxon divergence: a phylogenetic analysis in legumes using 39 gene families. *Syst Biol* 54: 441–454
- Ramalingam J, Vera Cruz CM, Kukreja K, Chittoor JM, Wu JL, Lee SW, Baraoidan M, George ML, Cohen MB, Hulbert SH, et al (2003) Candidate defense genes from rice, barley, and maize and their association with qualitative and quantitative resistance in rice. *Mol Plant Microbe Interact* 16: 14–24
- Rensing SA, Lang D, Zimmer AD, Terry A, Salamov A, Shapiro H, Nishiyama T, Perroud PF, Lindquist EA, Kamisugi Y, et al (2008) The Physcomitrella genome reveals evolutionary insights into the conquest of land by plants. *Science* 319: 64–69
- Riechmann JL, Heard J, Martin G, Reuber L, Jiang C, Keddie J, Adam L, Pineda O, Ratcliffe OJ, Samaha RR, et al (2000) Arabidopsis transcription factors: genome-wide comparative analysis among eukaryotes. *Science* 290: 2105–2110
- Schauser L, Roussis A, Stiller J, Stougaard J (1999) A plant regulator controlling development of symbiotic root nodules. *Nature* 402: 191–195
- Schauser L, Wieloch W, Stougaard J (2005) Evolution of NIN-like proteins in Arabidopsis, rice, and *Lotus japonicus*. *J Mol Evol* 60: 229–237
- Schlueter JA, Dixon P, Granger C, Grant D, Clark L, Doyle JJ, Shoemaker RC (2004) Mining EST databases to resolve evolutionary events in major crop species. *Genome* 47: 868–876
- Schlueter JA, Lin JY, Schlueter SD, Vasylenko-Sanders IF, Deshpande S, Yi J, O'Bleness M, Roe BA, Nelson RT, Scheffler BE, et al (2007) Gene duplication and paleopolyploidy in soybean and the implications for whole genome sequencing. *BMC Genomics* 8: 330
- Schlueter JA, Scheffler BE, Jackson S, Shoemaker RC (2008) Fractionation of synteny in a genomic region containing tandemly duplicated genes across *Glycine max*, *Medicago truncatula*, and *Arabidopsis thaliana*. *J Hered* 99: 390–395
- Schmid M, Davison TS, Henz SR, Pape UJ, Demar M, Vingron M, Scholkopf B, Weigel D, Lohmann JU (2005) A gene expression map of *Arabidopsis thaliana* development. *Nat Genet* 37: 501–506
- Shultz JL, Ray JD, Lightfoot DA (2007) A sequence based synteny map between soybean and *Arabidopsis thaliana*. *BMC Genomics* 8: 8
- Smit P, Raedts J, Portyanko V, Debelle F, Gough C, Bisseling T, Geurts R (2005) NSP1 of the GRAS protein family is essential for rhizobial Nod factor-induced transcription. *Science* 308: 1789–1791
- Stracker S, Kistner C, Yoshida S, Mulder L, Sato S, Kaneko T, Tabata S, Sandal N, Stougaard J, Szczylowski K, et al (2002) A plant receptor-like kinase required for both bacterial and fungal symbiosis. *Nature* 417: 959–962
- Sumimoto H, Kamakura S, Ito T (2007) Structure and function of the PB1 domain, a protein interaction module conserved in animals, fungi, amoebas, and plants. *Sci STKE* 2007: re6
- Taylor SA, Hofer JM, Murfet IC, Sollinger JD, Singer SR, Knox MR, Ellis TH (2002) *PROLIFERATING INFLORESCENCE MERISTEM*, a MADS-box gene that regulates floral meristem identity in pea. *Plant Physiol* 129: 1150–1159
- Tuskan GA, Difazio S, Jansson S, Bohlmann J, Grigoriev I, Hellsten U, Putnam N, Ralph S, Rombauts S, Salamov A, et al (2006) The genome of black cottonwood, *Populus trichocarpa* (Torr. & Gray). *Science* 313: 1596–1604
- Udvardi MK, Kakar K, Wandrey M, Montanari O, Murray J, Andriankaja A, Zhang JY, Benedito V, Hofer JM, Chueng F, et al (2007) Legume transcription factors: global regulators of plant development and response to the environment. *Plant Physiol* 144: 538–549
- UniProt Consortium (2009) The Universal Protein Resource (UniProt) 2009. *Nucleic Acids Res* 37: D169–D174
- Valdés-López O, Arenas-Huertero C, Ramírez M, Girard L, Sánchez F, Vance CP, Reyes JL, Hernández G (2008) Essential role of MYB transcription factor: PvPHR1 and microRNA: PvmiR399 in phosphorus-deficiency signaling in common bean roots. *Plant Cell Environ* 31: 1834–1843

- Vernié T, Moreau S, de Billy F, Plet J, Combiér JP, Rogers C, Oldroyd G, Frugier F, Niebel A, Gamas P (2008) *EFD* is an *ERF* transcription factor involved in the control of nodule number and differentiation in *Medicago truncatula*. *Plant Cell* **20**: 2696–2713
- Wang H, Moore MJ, Soltis PS, Bell CD, Brockington SF, Alexandre R, Davis CC, Latvis M, Manchester SR, Soltis DE (2009) Rosid radiation and the rapid rise of angiosperm-dominated forests. *Proc Natl Acad Sci USA* **106**: 3853–3858
- Wang HW, Zhang B, Hao YJ, Huang J, Tian AG, Liao Y, Zhang JS, Chen SY (2007) The soybean *Dof-type* transcription factor genes, *GmDof4* and *GmDof11*, enhance lipid content in the seeds of transgenic Arabidopsis plants. *Plant J* **52**: 716–729
- Wilson D, Charoensawan V, Kummerfeld SK, Teichmann SA (2008) DBD: taxonomically broad transcription factor predictions. New content and functionality. *Nucleic Acids Res* **36**: D88–D92
- Wojciechowski MF (2003) Reconstructing the phylogeny of legumes (Fabaceae): an early 21st century perspective. In BB Klitgaard, A Bruneau, eds, *Advances in Legume Systematics. Part 10. Higher Level Systematics*. Royal Botanic Garden, Kew, UK, pp 5–35
- Yan HH, Mudge J, Kim DJ, Larsen D, Shoemaker RC, Cook DR, Young ND (2003) Estimates of conserved microsynteny among the genomes of *Glycine max*, *Medicago truncatula* and *Arabidopsis thaliana*. *Theor Appl Genet* **106**: 1256–1265
- Yan HH, Mudge J, Kim DJ, Shoemaker RC, Cook DR, Young ND (2004) Comparative physical mapping reveals features of microsynteny between *Glycine max*, *Medicago truncatula*, and *Arabidopsis thaliana*. *Genome* **47**: 141–155
- Ye C, Galbraith SJ, Liao JC, Eskin E (2009) Using network component analysis to dissect regulatory networks mediated by transcription factors in yeast. *PLOS Comput Biol* **5**: e1000311
- Yoshida K, Iwasaka R, Kaneko T, Sato S, Tabata S, Sakuta M (2008) Functional differentiation of *Lotus japonicus* TT2s, R2R3-MYB transcription factors comprising a multigene family. *Plant Cell Physiol* **49**: 157–169
- Young ND, Udvardi M (2009) Translating *Medicago truncatula* genomics to crop legumes. *Curr Opin Plant Biol* **12**: 193–201
- Yu J, Hu S, Wang J, Wong GK, Li S, Liu B, Deng Y, Dai L, Zhou Y, Zhang X, et al (2002) A draft sequence of the rice genome (*Oryza sativa* L. ssp. *indica*). *Science* **296**: 79–92
- Zhang G, Chen M, Chen X, Xu Z, Guan S, Li LC, Li A, Guo J, Mao L, Ma Y (2008) Phylogeny, gene structures, and expression patterns of the *ERF* gene family in soybean (*Glycine max* L.). *J Exp Bot* **59**: 4095–4107
- Zhou QY, Tian AG, Zou HF, Xie ZM, Lei G, Huang J, Wang CM, Wang HW, Zhang JS, Chen SY (2008) Soybean *WRKY-type* transcription factor genes, *GmWRKY13*, *GmWRKY21*, and *GmWRKY54*, confer differential tolerance to abiotic stresses in transgenic Arabidopsis plants. *Plant Biotechnol J* **6**: 486–503
- Zhu B, Ye C, Lu H, Chen X, Chai G, Chen J, Wang C (2006) Identification and characterization of a novel heat shock transcription factor gene, *GmHsfA1*, in soybeans (*Glycine max*). *J Plant Res* **119**: 247–256
- Zhu H, Chen T, Zhu M, Fang Q, Kang H, Hong Z, Zhang Z (2008) A novel ARID DNA-binding protein interacts with SymRK and is expressed during early nodule development in *Lotus japonicus*. *Plant Physiol* **148**: 337–347
- Zhu H, Choi HK, Cook DR, Shoemaker RC (2005) Bridging model and crop legumes through comparative genomics. *Plant Physiol* **137**: 1189–1196
- Zhu H, Kim DJ, Baek JM, Choi HK, Ellis LC, Kuester H, McCombie WR, Peng HM, Cook DR (2003) Syntenic relationships between *Medicago truncatula* and Arabidopsis reveal extensive divergence of genome organization. *Plant Physiol* **131**: 1018–1026