

# EXORDIUM-LIKE1 Promotes Growth during Low Carbon Availability in Arabidopsis<sup>1[C][W]</sup>

Florian Schröder, Janina Lisso, and Carsten Müssig\*

Universität Potsdam, Max Planck Institute of Molecular Plant Physiology, Department Lothar Willmitzer, 14476 Golm, Germany (F.S., J.L.); and GoFORSYS, Universität Potsdam, 14476 Golm, Germany (C.M.)

Little is known about genes that control growth and development under low carbon (C) availability. The Arabidopsis (*Arabidopsis thaliana*) EXORDIUM-LIKE1 (*EXL1*) gene (At1g35140) was identified as a brassinosteroid-regulated gene in a previous study. We show here that the EXL1 protein is required for adaptation to C- and energy-limiting growth conditions. In-depth analysis of *EXL1* transcript levels under various environmental conditions indicated that *EXL1* expression is controlled by the C and energy status. Sugar starvation, extended night, and anoxia stress induced *EXL1* gene expression. The C status also determined EXL1 protein levels. These results suggested that EXL1 is involved in the C-starvation response. Phenotypic changes of an *exl1* loss-of-function mutant became evident only under corresponding experimental conditions. The mutant showed diminished biomass production in a short-day/low-light growth regime, impaired survival during extended night, and impaired survival of anoxia stress. Basic metabolic processes and signaling pathways are presumed to be barely impaired in *exl1*, because the mutant showed wild-type levels of major sugars, and transcript levels of only a few genes such as *QUAQUINE STARCH* were altered. Our data suggest that EXL1 is part of a regulatory pathway that controls growth and development when C and energy supply is poor.

The coordination of carbon (C) supply and growth in varying environmental conditions is a major challenge for plants. The rate of CO<sub>2</sub> assimilation, the accumulation of storage carbohydrates during the day, the use of assimilates for growth, and the rate of carbohydrate degradation at night represent processes that require coordinated regulation (Paul and Pellny, 2003; Smith and Stitt, 2007). The diurnal C balance is of pivotal importance for the maintenance of growth, because even transient periods of carbohydrate deficiency cause growth inhibition by temporary inhibition of carbohydrate utilization (Gibon et al., 2004). Short-term undesirable conditions such as moisture deficit and cold temperatures reduce photosynthetic C supply, and biotic damage to leaves or defoliation can interrupt photosynthetic activity.

Plants evolved mechanisms that ensure the coordination of growth with metabolism. These mechanisms imply both long-distance and tissue- or cell type-specific signaling mechanisms (Rolland et al., 2006). Glc, Fru, and Suc are the main products of photosynthesis. Sugar-dependent feedback mechanisms control

photosynthesis (Paul and Pellny, 2003). Sink tissues modify local and long-distance sugar signaling by the level of sugar import and the action of Suc-cleaving enzymes (Koch, 2004). The C and energy status controls metabolic pathways by posttranslational regulation of enzymes (Huber and Hardin, 2004). In addition, sugar availability controls transcription, translation, and protein stability of transcription factors (Smeekens et al., 2010) and modulates global gene expression patterns (Ho et al., 2001; Contento et al., 2004; Lee et al., 2004; Thimm et al., 2004; Bläsing et al., 2005; Usadel et al., 2008).

Although sugar regulation is necessarily far more complex in multicellular organisms, yeast is a useful model for sugar sensing and signaling at the cellular level (Rolland et al., 2006). Three major Glc sensing and signaling pathways have been identified in yeast. Glc is the preferred C source for yeast. The metabolism of other C substrates is repressed if Glc is available. This phenomenon is referred to as "Glc repression." It requires the activity of the glycolytic enzyme and sensor Hexokinase2 (Hxk2). In response to Glc, Hxk2 is transported into the nucleus, where it interacts with the transcription factor Multicopy inhibitor of GAL expression1 (Mig1) and generates a repressor complex. This complex is responsible for the repression of genes involved in the transport and metabolism of alternative C substrates, gluconeogenesis, and respiration (Moreno et al., 2005). The activity of Mig1 is also controlled by a key protein kinase, Sucrose nonfermenting1 (Snf1). Upon Glc depletion, Snf1 is activated and controls many processes through the phosphorylation of transcription factors and metabolic enzymes. The general effect of the Snf1 kinase in yeast is to

<sup>1</sup> This work was supported by the Deutsche Forschungsgemeinschaft (grant nos. MU 1738/4-1 and MU 1738/4-3).

\* Corresponding author; e-mail muessig@uni-potsdam.de.

The author responsible for distribution of materials integral to the findings presented in this article in accordance with the policy described in the Instructions for Authors ([www.plantphysiol.org](http://www.plantphysiol.org)) is: Carsten Müssig (muessig@uni-potsdam.de).

<sup>[C]</sup> Some figures in this article are displayed in color online but in black and white in the print edition.

<sup>[W]</sup> The online version of this article contains Web-only data.

[www.plantphysiol.org/cgi/doi/10.1104/pp.111.177204](http://www.plantphysiol.org/cgi/doi/10.1104/pp.111.177204)

switch the cellular metabolism from fermentation to respiration (Zhang et al., 2010). Phosphorylation of Mig1 by Snf1 is essential for the nuclear export of Mig1 and the derepression of genes that are repressed in the presence of high Glc concentrations (Ahuatzi et al., 2007). Fermentative Glc metabolism requires a high metabolic flux through glycolysis. The expression of hexose transporters (Hxts) is up-regulated through the action of the Hxt homologs Snf3 and Restores glucose transport2 (Rgt2; Rolland et al., 2006). Snf3 and Rgt2 function as sensors for extracellular Glc (Gancedo, 2008). The third Glc signaling pathway in yeast is based on Glc activation of cAMP synthesis. Activation of cAMP synthesis by the adenylate cyclase Cyr1 requires Glc phosphorylation by either Glucokinase1 (Glk1) or one of the two hexokinases (Hxk1 or Hxk2) and Glc activation of the G-protein-coupled receptor Gpr1. The cAMP-dependent protein kinases A (PKAs; Tpk1–Tpk3) control a large number of proteins, including transcriptional regulators such as Rgt1, Msn2, and Msn4. The PKAs stimulate glycolytic activity, the mobilization of reserve carbohydrates, and ribosome biogenesis. Furthermore, the cAMP-PKA pathway mediates cell growth and represses the expression of stress resistance genes (Rolland et al., 2006; Gancedo, 2008).

Experimental evidence suggests the presence of at least three systems for Glc sensing and additional systems for Suc sensing in plants. The AtHXK1/GIN2 hexokinase has Glc signaling functions that are uncoupled from metabolic activities (Jang et al., 1997; Zhou et al., 1998; Moore et al., 2003). It is mainly associated with mitochondria, possibly as part of a glycolytic metabolon, but also found in the nucleus (Yanagisawa et al., 2003). The Arabidopsis (*Arabidopsis thaliana*) genome encodes two further hexokinases and three hexokinase-like (HKL) proteins. The HKL1 protein lacks glucokinase activity and plays a specific role (Karve and Moore, 2009). In contrast to yeast and animals, G-protein signaling is poorly characterized in plants. AtRGS1, an unusual hybrid seven-transmembrane domain protein with a C-terminal RGS box with GTPase-accelerating activity (Chen et al., 2003), potentially acts as an extracellular receptor for Glc and mediates Glc regulation of a limited set of genes (Grigston et al., 2008). The Suc transporter-like protein SUT2 resembles the yeast Snf3 and Rgt2 proteins (Barker et al., 2000), but a role as extracellular Glc or Suc sensor has not been demonstrated. Evidence for additional sugar sensing mechanisms in plants comes from experiments with Glc and Suc analogs and mutants with defects in sugar responses.

Evolutionarily conserved protein kinases termed Snf1 in yeast, AMPK in mammals, and Snf1-Related Protein Kinase1 (SnRK1) in plants are major components of energy homeostasis maintenance in eukaryotes (Baena-González and Sheen, 2008; Zhang et al., 2010). As mentioned above, Snf1 in yeast is activated upon Glc depletion and controls many processes through the regulation of transcription factors and

metabolic enzymes. SnRK1 in plants represents a master regulator during C and energy deprivation (Baena-González and Sheen, 2008; Halford and Hey, 2009; Jossier et al., 2009). SnRK1.1/KIN10-mediated gene expression is positively correlated with that regulated by C deprivation and negatively correlated with that controlled by sugar (Baena-González et al., 2007). Plant SnRK1 kinases are also involved in the posttranslational regulation of key enzymes and thus may directly regulate major biosynthetic pathways (Sugden et al., 1999). The activity of SnRK1 is inhibited by proteins such as PRL1 (Bhalerao et al., 1999) and the sugar-induced signaling molecule trehalose-6-phosphate (Zhang et al., 2009). AtSnAK1/GRIK2 and AtSnAK2/GRIK1 act as upstream kinases and activate SnRK1 (Hey et al., 2007; Shen et al., 2009). The calcineurin B-like-interacting protein kinase CIPK15 is an upstream regulator of SnRK1A in rice (*Oryza sativa*) and links oxygen-deficiency signals to the SnRK1-dependent sugar sensing cascade (Lu et al., 2007; Lee et al., 2009).

Several mutants impaired in phytohormone biosynthesis or signaling also show altered sugar responses. For example, abscisic acid-deficient or abscisic acid-insensitive mutants are Glc insensitive (Arenas-Huertero et al., 2000; Huijser et al., 2000; Laby et al., 2000; Cheng et al., 2002). Ethylene overproduction mutants and mutants with constitutive ethylene signaling show reduced responses to high levels of Glc or Suc, whereas ethylene-insensitive mutants are hypersensitive to sugar (Zhou et al., 1998; Cheng et al., 2002).

In this study, we characterize the Arabidopsis gene *EXORDIUM-LIKE1* (*EXL1*). *EXL1* was identified as a BR-induced gene (Schröder et al., 2009). *EXL1* encodes a putative protein of 309 amino acids that is the closest homolog to the previously characterized EXO protein (67% identity and 79% similarity; Farrar et al., 2003; Coll-Garcia et al., 2004; Schröder et al., 2009). Structure prediction tools such as PSIPRED (McGuffin et al., 2000) suggest that the positions of helices and strands are highly conserved in EXO and *EXL1*. The major structural feature of the proteins is the so-called PHOSPHATE-INDUCED PROTEIN1 conserved region (see Pfam entry PF04674 [Finn et al., 2008]). Proteins harboring this region were identified in evolutionarily distant green plants (Schröder et al., 2009). The molecular function of the region is unknown. The EXO and *EXL1* primary sequences do not show similarities to other known protein domains. In silico analyses and experimental evidence indicate an extracellular localization of the proteins. EXO, *EXL1*, and other members of the protein family carry an N-terminal signal peptide. Sequence analysis tools such as TargetP (Emanuelsson et al., 2000) and WoLF PSORT (Horton et al., 2007) predict that the proteins enter the secretory pathway (Schröder et al., 2009). Several proteomics approaches identified EXO, *EXL1*, and other members of the protein family as part of the cell wall proteome (Borderies et al., 2003; Bayer et al., 2006; Feiz et al., 2006; Jamet et al., 2006), and EXO:HA (for hemagglu-

tinin) and EXO:GFP fusion proteins were detected in the apoplast (Schröder et al., 2009).

Here, we show that *EXL1* plays an essential role under specific environmental conditions. *EXL1* is largely irrelevant under favorable light conditions and nonlimiting C supply but controls growth and development under low C and energy availability. Our evidence suggests that *EXL1* is part of a regulatory mechanism that serves to balance the C status with development and growth.

## RESULTS

### Identification of Mutants

An *exl1* knockout mutant (GABI-Kat line 592A10; Rosso et al., 2003) was identified. The mutant carries a T-DNA insertion in the coding region (126 bp upstream of the stop codon). Northern-blot and reverse transcription (RT)-PCR analyses did not detect *EXL1* transcript (data not shown). Introduction of a 35S::*EXL1* construct into the *exl1* mutant (*exl1/35S::EXL1*) resulted in strong *EXL1* expression in the mutant and normalized the phenotypic changes (see below). Thus, the *exl1* phenotype was caused by the T-DNA insertion into the *EXL1* coding sequence. Furthermore, a knock-on mutant termed *exl1-D* (SALK line 027172; Alonso et al., 2003) was identified. The *exl1-D* mutant carries a T-DNA insertion at position -467 upstream of the translation start codon and showed a several-fold increase in *EXL1* transcript levels in comparison with the wild type.

### *EXL1* Expression Is Induced by C Starvation

Expression of *EXL1* and *EXO* is induced by BR (Schröder et al., 2009), and the encoded proteins are highly similar. Thus, we assumed that *EXL1* and *EXO* play similar roles in the control of vegetative growth and tested biomass production and development of the *exl1* mutant in a greenhouse. In contrast to *EXO*, the loss of *EXL1* was largely irrelevant for growth and development under standard conditions.


Therefore, we used Web-based platforms such as Genevestigator (Hruz et al., 2008) and the Arabidopsis eFP Browser (Winter et al., 2007) to identify experimental conditions that have a strong impact on *EXL1* expression. The data indicated that *EXL1* expression is controlled by the C status. Relevant Affymetrix ATH1 microarray profiles were downloaded from the Gene Expression Omnibus (Edgar et al., 2002) and analyzed using the ROBIN software that is based on functions of the R/BioConductor project (Lohse et al., 2010). In addition, plants were grown in synthetic medium supplemented with different sugar levels or in soil under different light regimes. *EXL1* transcript and protein levels were analyzed by means of quantitative RT-PCR and western-blot analysis, respectively.

The expression of *EXL1* showed the following profile. First, *EXL1* transcript levels decreased slightly

during the light period and increased slightly during the night in wild-type plants (Fig. 1). This diurnal cycling was intensified in the *pgm* mutant (Fig. 1A, top right panel). The *pgm* mutant does not form transitory starch (Caspar et al., 1985) and is unable to maintain appropriate carbohydrate and energy levels during the night. Consequently, *pgm* plants show poor growth and symptoms of C starvation in the dark period (Gibson et al., 2006). Second, wild-type plants showed strong *EXL1* expression in extended night (Fig. 1, A, bottom left panel, and C). Third, *EXL1* mRNA levels decreased in wild-type seedlings grown in liquid culture upon sugar supply (Fig. 1A, bottom right panel). Fourth, wild-type plants grown in an atmosphere with 50  $\mu\text{L L}^{-1}$   $\text{CO}_2$  showed higher *EXL1* expression than plants grown in 350  $\mu\text{L L}^{-1}$   $\text{CO}_2$  (Fig. 1A, bottom right panel). Fifth, addition of increasing levels of Suc to synthetic growth medium suppressed *EXL1* expression (data not shown). Sixth, overexpression of SnRK1.1 (also known as KIN10 or AKIN10) resulted in elevated *EXL1* mRNA levels (Baena-González et al., 2007). In summary, low C availability and SnRK1.1/KIN10 action result in elevated *EXL1* transcript levels.

In addition to mRNA synthesis and stability, translation efficiency and protein stability could determine the *EXL1* protein level. *EXL1:HA* transcript and protein levels were assessed in plants carrying the pEXL1::*EXL1:HA* and 35S::*EXL1:HA* constructs. The pEXL1::*EXL1:HA* construct conferred expression of an HA-tagged *EXL1* protein driven by a genomic *EXL1* fragment that is located upstream of the *EXL1* translation start codon (positions -1 to -928). The sequence of several full-length cDNAs and promoter analysis tools such as PPDB (Yamamoto and Obokata, 2008) indicate a length for the 5' untranslated region of *EXL1* of about 92 bp and a putative TATA box at positions -110 to -121. Thus, the cloned promoter sequence in the pEXL1::*EXL1:HA* construct spans about 830 bp. The 35S::*EXL1:HA* construct resulted in the expression of an identical *EXL1:HA* protein under control of the cauliflower mosaic virus 35S promoter.

High levels of *EXL1:HA* transcript were detected in shoots of the pEXL1::*EXL1:HA* transgenic lines when plants were grown in the presence of 0.2% (w/v) Suc. Accordingly, decreasing transcript levels were detected when plants were grown in the presence of 0.5%, 1%, 3%, or 5% Suc (Fig. 2A). *EXL1:HA* expression was less repressed by sugar in roots (Fig. 2B). *EXL1:HA* expression driven by the *EXL1* promoter resulted in corresponding *EXL1:HA* protein levels (Fig. 3). The 35S promoter caused strong *EXL1:HA* expression under all conditions (Fig. 2) and largely prevented the modulation of *EXL1:HA* transcript (Fig. 2) and *EXL1:HA* protein levels (data not shown) by the sugar supply and light regime. These data indicate that transcriptional regulation of *EXL1* confers an adjustment of *EXL1* protein levels to the C status.



**Figure 1.** *EXL1* expression in response to C availability. A, Expression profiles from publicly available microarray data (Bläsing et al., 2005; Gibon et al., 2006; Osuna et al., 2007; Usadel et al., 2008). The data were normalized using RMA and  $\log_2$  transformed. The means of two or three replicates are shown. The  $\pm$  SE is shown if three replicates were available. Expression of *EXO* is shown for comparison. FN, Full nutrition; C-starv, C starvation. B, Quantitative RT-PCR analysis of *EXL1* transcript levels in wild-type (Col-0) shoots. Plants were grown in soil under the low-total-irradiance regime (4 h of light [ $60 \mu\text{mol m}^{-2} \text{s}^{-1}$ ]/20 h of dark). RNA was extracted from shoots of 28-d-old plants. *elf1 $\alpha$*  cycle threshold values were subtracted from respective cycle threshold values of the gene of interest. Subsequently, differences were subtracted from an arbitrary value (i.e. 40). Higher numbers indicate higher transcript levels. A difference of 1 unit indicates a 2-fold change. Error bars indicate  $\pm$  SE of the gene of interest in three technical replicates. The data shown are from one experiment representative of three independent biological replicates. eN, End of night; eD, end of day. C, Quantitative RT-PCR analysis of *EXL1* transcript levels in wild-type (Col-0) shoots. Plants were grown in soil under short-day conditions (8 h of light [ $140 \mu\text{mol m}^{-2} \text{s}^{-1}$ ]/16 h of dark). RNA was extracted from 35-d-old plants. Technical details are as given for B. [See online article for color version of this figure.]


### EXL1 Controls Growth and Development during Low C Availability

Plants were grown in half-concentrated Murashige and Skoog (MS) medium supplemented with different sugar quantities. Sugar responses were tested both under standard growth conditions (16 h of light [ $145 \mu\text{mol m}^{-2} \text{s}^{-1}$ ]/8 h of dark) and in continuous darkness.

The *exl1* mutant developed like the wild type when grown in the presence of 1% (w/v) Suc in a diurnal cycle. Biomass production and leaf size were similar. Next, plants were grown under standard conditions in the presence of high sugar levels (3%–6% [w/v] Suc). High sugar levels similarly impaired the development of the mutant and the wild type. No differences

between the genotypes were observed. In contrast, loss of *EXL1* caused slightly reduced leaf growth and a significant 27% reduction in biomass in the presence of 0.2% Suc (controls in Figs. 4 and 5).

Arabidopsis seedlings can develop leaf- and flower-like organs in the dark, but skotomorphogenesis critically depends on the sugar availability (Roldán et al., 1999; Baier et al., 2004; Li et al., 2007). Similar to photomorphogenic development, skotomorphogenic development of *exl1* was identical with the wild type when grown in medium supplemented with 1% Suc or 1% Glc. Nearly all plants developed true leaves after 3 weeks. Sporadically, plants developed another pair of true leaves (data not shown). There were also no clear differences between the genotypes in the presence of 0.05% Suc (Fig. 6A), presumably because very low



**Figure 2.** *EXL1:HA* expression in response to different Suc levels and light regimes. A, Relative *EXL1:HA* transcript levels in shoots. Plants carrying the pEXL1::EXL:HA and 35S::EXL1:HA constructs were grown in synthetic medium supplemented with 0.2%, 0.5%, 1%, 3%, and 5% (w/v) Suc. RNA was extracted from 14-d-old plants. Technical details are as given for Figure 1B. The data shown are from one experiment representative of three independent biological replicates. B, Relative *EXL1:HA* transcript levels in roots. RNA was extracted from the same plants as in A. C, Relative *EXL1:HA* transcript levels in shoots. Plants were grown in soil under light-limited conditions (4 h of light [ $60 \mu\text{mol m}^{-2} \text{s}^{-1}$ ]/20 h of dark) and subjected to extended night. RNA was extracted from 35-d-old plants. eD, End of day; eN, end of night. D, Relative *EXL1:HA* transcript levels in shoots. Plants were grown in soil under long-day conditions (16 h of light/8 h of dark) and subjected to extended night. RNA was extracted from 28-d-old plants.

sugar levels prohibited growth. Supply of 0.15% Suc or Glc, however, caused accelerated development of the *exl1* mutant. Wild-type plants typically developed elongated hypocotyls and opened cotyledons but hardly produced true leaves. In contrast, approximately 15% of the *exl1* plants developed an internodium and true leaves (Fig. 6). Introduction of the 35S::EXL1 construct into the *exl1* mutant suppressed true leaf formation (Fig. 6A). Thus, the loss of *EXL1* resulted in enhanced dark development in the presence of low sugar levels.

#### EXL1 Limits BR-Promoted Growth during Low C Availability


BR treatments were used to enforce growth. Growth of wild-type and *exl1* plants was tested in half-concentrated MS medium supplemented with 0.2% or 1% (w/v) Suc and 0 nM BR (control), 50 nM castasterone, or 50 nM brassinolide. Exogenous BR stimu-

lated growth at both Suc levels, but the relative BR effect on biomass production was more pronounced at 0.2% Suc. Application of 50 nM castasterone or 50 nM brassinolide resulted in a 1.7- to 2-fold increase of biomass in wild-type plants and a 2.8- to 3.3-fold increase of biomass in *exl1*, respectively (Fig. 4). In contrast, the biomass of *exl1* was similar to that of the wild type at 1% Suc (Fig. 4).


Several studies have shown that BR promotes leaf growth. For example, enhancement of BR responses by means of *DWF4*, *BR11*, or *BAK1* overexpression or application of synthetic BR resulted in leaves that are characterized by longer petioles and an increased leaf index (Choe et al., 2001; Wang et al., 2001; Nam and Li, 2002). To investigate the influence of *EXL1* on leaf growth, plants were grown in synthetic medium supplemented with 0.2% Suc and 0 nM BR (control), 50 nM 24-epibrassinolide, or 50 nM brassinolide. Leaf length and leaf width were determined after 16 d. In the absence of exogenous BR, leaf size of *exl1* plants was slightly smaller in comparison with the wild type, but leaf indices did not differ significantly (Fig. 5). Application of 50 nM 24-epibrassinolide or 50 nM brassinolide resulted in a significant increase of the leaf index in all genotypes. The BR-promoted growth response was more pronounced in plants with impaired *EXL1* expression. For example, the presence of 50 nM brassinolide caused a 1.7-fold increase of leaf length in the wild type and a 2.4-fold increase in *exl1* (Fig. 5).

#### EXL1 Is Required for Growth under Light-Limited Conditions

In order to examine the role of *EXL1* in the adaptation to low C under physiological conditions, plants were grown in soil under short-day conditions with low light (4 h of light [ $60 \mu\text{mol m}^{-2} \text{s}^{-1}$ ]/20 h of darkness). The low photosynthetically active radiation and long dark period resulted in a low total irradiance



**Figure 3.** Western-blot analysis of EXL1:HA protein levels. Plants carrying the pEXL1::EXL:HA construct were grown under different conditions. The EXL1:HA fusion protein was detected using a monoclonal anti-HA antibody and an enhanced chemiluminescence detection system. Similar results were obtained for independent transgenic lines. A, Plants were grown in synthetic medium supplemented with 0.2%, 0.5%, 1%, 3%, and 5% (w/v) Suc. Protein was extracted from the same shoot material as used for transcript analysis (Fig. 2A). B, Plants were grown in long-day conditions as described in the legend of Figure 2D. eD, End of day; eN, end of night. C, Plants were grown under light-limited conditions as described in the legend of Figure 2C.



**Figure 4.** Biomass production in response to Suc and BR. Plants were grown in synthetic medium supplemented with 0.2% or 1% (w/v) Suc and brassinosteroids (50 nM castasterone [CS] or 50 nM brassinolide [BL]). Fresh weight was determined after 16 d. Results are means  $\pm$  SE ( $n = 5$  plates per genotype and treatment containing 10 plants each). Biomass of *exl1* plants was significantly reduced in the control treatment with 0.2% Suc (denoted with an asterisk;  $t$  test,  $P < 0.01$ ). Percentage differences in BR treatments with 0.2% Suc supply were significantly larger in *exl1* in comparison with the wild type ( $t$  test,  $P < 0.05$ ).


that limited growth. Several independent experiments revealed that the *exl1* mutant showed less growth in comparison with the wild type (Fig. 7A). Fresh weight and dry weight of *exl1* were reduced by approximately 50% (Fig. 7B). Thus, *EXL1* is necessary for growth under light-limited conditions. The *exl1-D* mutant did not show improved growth and biomass production, suggesting that wild-type *EXL1* expression is sufficient under these conditions.

Reduced growth of *exl1* could be due to defects in the primary metabolism of major carbohydrates. Altered levels of starch, Glc, Fru, or Suc are indicative of such defects. Starch levels of wild-type, *exl1-D*, and *exl1* plants were measured based on the enzymatic hydrolysis and photometric determination of Glc. Plant material of three independent short-day/low-light experiments was analyzed. No significant differences were detected between the genotypes. Differences


between independent experiments actually exceeded differences between the genotypes (wild-type, 2.5–3.8  $\mu\text{mol hexose equivalents mg}^{-1}$  chlorophyll [means, end of night] and 8.9–12.9  $\mu\text{mol hexose equivalents mg}^{-1}$  chlorophyll [means, end of day]; *exl1-D*, 2.6–3.7 and 8.8–10.5  $\mu\text{mol mg}^{-1}$  chlorophyll; *exl1*, 2.2–3.5 and 11.5–11.6  $\mu\text{mol mg}^{-1}$  chlorophyll; similar results were obtained when starch levels were related to dry weight). Similarly, Glc, Fru, and Suc levels did not differ significantly (data not shown). This suggests that major metabolic pathways are functional in *exl1*, although it cannot be excluded that the plasticity of metabolic networks compensates for perturbations.

Wild-type and *exl1* plants were also analyzed by means of gene expression profiling. Eight profiles were established using Affymetrix ATH1 microarrays. First, wild-type and *exl1* plants were grown in an 8-h-day (140  $\mu\text{mol m}^{-2} \text{s}^{-1}$ )/16-h-night regime. These conditions allowed largely normal growth of *exl1*. Second, wild-type and *exl1* plants were grown under the low-total-irradiance regime mentioned above (4 h of light [60  $\mu\text{mol m}^{-2} \text{s}^{-1}$ ]/20 h of darkness). These conditions caused significantly less growth of *exl1* in comparison with the wild-type. For each light regime and genotype, plant material was harvested at the beginning and the end of the light period.

The activity of most genes was virtually identical under the 8-h-light/16-h-dark growth regime. Only nine and 14 genes showed at least a 2-fold change in the comparison of *exl1* versus the wild type at the end of the night and the end of the day, respectively (Supplemental File S1). A single gene, *QUA-QUINE STARCH (QQS; At3g30720)*, showed consistent up-regulation in *exl1* (7.3- and 5.9-fold change; Supplemental File S1). The wild-type and *exl1* expression profiles were more different under the low-total-irradiance regime. Twenty-six and 40 genes showed at least a 2-fold change in *exl1* compared with the wild type at the end of the night and the end of the day, respectively. The *QQS* gene again showed stronger expression in *exl1* (2.8- and 3.4-fold change; Supplemental File S1). *QQS*



**Figure 5.** Leaf growth in response to Suc and BR. Plants were grown in synthetic medium supplemented with 0.2% (w/v) Suc and brassinosteroids (50 nM brassinolide [BL] or 50 nM 24-epibrassinolide [EBL]). Length and width of primary leaves were determined after 16 d. Results are means  $\pm$  SE ( $n = 5$  plates per genotype and treatment containing 10 plants each). Mutant values denoted with an asterisk are significantly different from those of their wild type ( $t$  test,  $P < 0.05$ ). Percentage differences of leaf length in BR treatments were significantly larger in *exl1* in comparison with the wild type ( $t$  test,  $P < 0.05$ ).



**Figure 6.** Dark development phenotypes. A, Seedlings were grown in half-concentrated MS medium supplemented with 0.05% or 0.15% (w/v) Suc for 3 weeks in the dark. Only the largest seedlings of a representative experiment are shown. B, Seedlings were grown in half-concentrated MS medium supplemented with 0.15% (w/v) Glc for 3 weeks in the dark. Development of plants was grouped into three categories: elongated hypocotyls with closed cotyledons (black bars), opened cotyledons (gray bars), and formation of true leaves (dark gray bars). The experiment was carried out independently three times. The observed responses from one representative experiment are shown.

transcript levels were studied in more detail by quantitative RT-PCR under light-limited conditions (4 h of low light; Fig. 8A), in a short-day diurnal cycle (8 h of light/16 h of dark), and in extended night (more than 16 h of dark; Fig. 8B). The diurnal regulation and induction in extended night of *QQS* in wild-type plants were unaltered in *exl1*, but *QQS* transcript levels were higher at all times in *exl1* compared with the wild type. *QQS* encodes a protein of 59 amino acids. The precise function is unknown, but it was suggested to play a role in the control of starch metabolism (Li et al., 2009).

Transcript levels of sugar-responsive genes (Price et al., 2004; Osuna et al., 2007), of genes that show diurnal changes (Schaffer et al., 2001; Bläsing et al., 2005; Michael et al., 2008), and of phytohormone-responsive genes (Goda et al., 2008) were barely changed in *exl1* in comparison with the wild type. The marginal changes in the expression profiles indicate that *exl1* does not suffer from a severe disturbance of sugar responses, from incorrect phasing of transcription under circadian conditions, or from defects in the phytohormone signaling network.

### EXL1 Warrants the Survival of Extended Night Periods and Confers Anoxia Tolerance

To examine the role of EXL1 in prolonged darkness, wild-type and *exl1* plants were grown in soil for 3 weeks in a controlled-growth chamber. Subsequently, the light was switched off. Survival rates were determined by transferring 15 plants per genotype and point in time into an illuminated greenhouse after different periods of extended night. Five independent experiments were performed. Survival rates of the *exl1* mutant were significantly lower in comparison with the wild type after 11 to 15 d in darkness (Fig. 9A).


Several experiments in the Genevestigator Stimulus database (Hruz et al., 2008) indicated that *EXL1* tran-

script levels were elevated under oxygen deficiency in comparison with normoxia. The most obvious effect of an oxygen shortage is an energy deficit, and genes that are under the control of the oxygen level may mediate the adaptation to an altered energy status. Anoxia tolerance of wild-type, *exl1*, and *exl1-D* plants was tested as described by Licausi et al. (2010). Plants were established in half-concentrated MS medium supplemented with 1% (w/v) Suc and subjected to temporary anoxia stress. Survival rates were negatively correlated with the length of the anoxic treatment and positively correlated with *EXL1* expression. Short anoxic treatments (6 h) allowed the survival of all plants (data not shown). However, 8- and 10-h anoxic treatments revealed differences between the genotypes. The *exl1* mutant showed significantly reduced survival rates, and the *exl1-D* mutant showed significantly higher survival rates in comparison with the wild type (Fig. 9B). The anoxia tolerance of *exl1-D* seedlings was not due to stronger growth (Supplemental File S2). Neither genotype survived longer anoxic treatments (more than 11 h).


## DISCUSSION

### EXL1 Transcript Accumulates during Low C and Low Energy Availability

Several lines of evidence indicate that *EXL1* expression is induced under low C and low energy availability (Figs. 1 and 2). For example, extended night, sugar starvation in synthetic medium, and diminished oxygen supply promoted *EXL1* expression. Diurnal changes of *EXL1* expression were strongly pronounced in the starch-deficient *pgm* mutant (Fig. 1A), and *EXL1* expression was under the control of SnRK1.1/KIN10 (Baena-González et al., 2007). The increase of the *EXL1*



**Figure 7.** Phenotype of the *exl1* mutant under growth-limiting light conditions. A, Wild-type, *exl1*, *exl1-D*, and complemented *exl1* plants were raised for 2 weeks in standard conditions and then grown for 2 weeks under light-limited conditions (4 h of light [ $60 \mu\text{mol s}^{-1} \text{m}^{-2}$ ]/20 h of dark). B, Fresh weight (FW) and dry weight (DW) of plants shown in A. Results are means  $\pm$  SE ( $n = 6$  individuals per genotype). Mutant values denoted with an asterisk are significantly different from those of their wild type ( $t$  test,  $P < 0.01$ ). [See online article for color version of this figure.]



**Figure 8.** QQS transcript levels in wild-type and *exl1* plants. A, Wild-type and *exl1* plants were grown in soil under the low-total-irradiance regime (4 h of light [ $60 \mu\text{mol m}^{-2} \text{s}^{-1}$ ]/20 h of dark). For the wild type, the same cDNAs were used as for EXL1 analysis (Fig. 1B). eN, End of night; eD, end of day. B, QQS expression in a diurnal cycle (end of night to 16 h of dark) and in extended night in wild-type and *exl1* plants. eN corresponds to 16 h of dark. The same wild-type cDNAs were used as for EXL1 analysis (Fig. 1C).

mRNA level was associated with an increase in the protein level (Fig. 3).

#### EXL1 Adapts Growth and Development to Low C Supply

Plants are able to avoid C deficiency under fluctuating environmental conditions because growth and development are synchronized with the supply and utilization of C (Gibon et al., 2004; Hummel et al., 2010). C signaling may ultimately determine the activity of biosynthetic and catabolic pathways (Stitt et al., 2007).

Sugar responses were assessed by plant cultivation on medium supplemented with different levels of sugar. Growth and development of *exl1* under a diurnal light regime and in constant darkness were similar to the wild type in the presence, for example, of 1% (w/v) Suc. Phenotypic changes only became apparent when C supply was suboptimal. The mutant produced less biomass and slightly smaller leaves under a diurnal light regime in medium supplemented with 0.2% Suc (Figs. 4 and 5). Skotomorphogenesis is a sensitive indicator of the effects of sugar on development, because C availability determines activity of the shoot apical meristem (Roldán et al., 1999; Baier et al., 2004). Dark development of *exl1* was accelerated in medium supplemented with 0.15% Glc or Suc (Fig. 6). The molecular basis of this result is unknown and could be manifold. Altered sugar transport, enhanced sugar responses, and cell wall changes were demonstrated to accelerate leaf and internodium formation in darkness (Baier et al., 2004; Li et al., 2007).


The conditional phenotype suggests that EXL1 controls growth and development only during low C availability. This hypothesis was tested further by the application of synthetic BR. The primary function of BR is growth promotion of nearly all plant organs (Müssig, 2005). BR-promoted growth depends on the C supply (Fig. 4; data not shown). The *exl1* mutant

showed a wild-type response to Suc and BR in the presence of 1% Suc. In contrast, BR-dependent biomass production and leaf growth were much stronger in *exl1* at 0.2% Suc (Figs. 4 and 5). The EXL1 protein could limit BR-dependent growth under low C availability. Alternatively, enhanced growth of the *exl1* mutant could be due to higher sensitivity to BR in low-C conditions. The molecular processes that link sugar responses and BR signaling have not been resolved so far. BR presumably coordinates and integrates diverse processes required for growth, partly via interactions with other phytohormones and signaling pathways. BR has been implicated in the control of C metabolism and C allocation. For example, BR modifies enzyme activities (e.g. acid invertases, Rubisco, and cytosolic  $\beta$ -amylase) and controls sink strength and source efficiency (Goetz et al., 2000; Schlüter et al., 2002; Lisso et al., 2006; Wu et al., 2008).

#### EXL1 Is Essential for Growth during Low Total Irradiance, Survival of Extended Night, and Survival of Anoxia Stress

The physiological role of EXL1 was tested in three additional experimental setups that are characterized by low C and low energy availability.

First, plants were grown in soil under short-day conditions with low light (4 h of light [ $60 \mu\text{mol m}^{-2} \text{s}^{-1}$ ]/20 h of dark). Plants can adapt to limited light, although this adaptation requires comprehensive reprogramming of the central metabolism and decrease of the growth rate (Gibon et al., 2004, 2009). Growth of the *exl1* mutant was clearly impaired under



**Figure 9.** Survival rates after extended night and anoxic stress. A, Plants were raised under short-day conditions (8 h of light/16 h of dark) and subjected to different periods of continuous darkness. After retransfer to a greenhouse with long-day conditions, plants were grown for 1 week prior to determining survival rates. Five independent experiments were analyzed. Results are mean survival rates  $\pm$  SE ( $n = 15$  plants per genotype, time, and experiment). Mutant values denoted with an asterisk are significantly different from those of their wild type ( $t$  test,  $P < 0.01$ ). B, Seven-day-old plants were subjected to anoxia for 8 and 10 h. After the treatment, plants were transferred to ambient air, and survival rates were determined after a 10-d recovery period. Results are mean survival rates  $\pm$  SE ( $n = 4$  plates containing at least 50 seedlings each). Mutant values denoted with an asterisk are significantly different from those of their wild type ( $t$  test,  $P < 0.01$ ).

these conditions in comparison with the wild type, resulting in approximately 50% less biomass production (Fig. 7). Therefore, the adaptation to light-limited conditions depends on EXL1. In contrast, growth was largely normal under short-day (8 h of light [ $140 \mu\text{mol m}^{-2} \text{s}^{-1}$ ]/16 h of dark), long-day (16 h of light [ $180 \mu\text{mol m}^{-2} \text{s}^{-1}$ ]/8 h of dark), and high-light (16 h of light [ $700 \mu\text{mol m}^{-2} \text{s}^{-1}$ ]/8 h of dark) conditions (data not shown).

Second, plants were subjected to extended night. Plants switch each day between high C availability in the light and a negative C balance at night. Starch turnover is regulated in wild-type plants, and only a small amount of starch is left at the end of the night (Smith et al., 2004; Usadel et al., 2008). Abrupt periods of extended night cause massive changes in global gene expression and of metabolite levels and eventually provoke an acute C limitation (Smith and Stitt, 2007). The *exl1* mutant is impaired in the adaptation to prolonged darkness, because survival rates were significantly smaller in comparison with the wild type (Fig. 9A).

Third, plants were subjected to anoxia stress. Plant responses to too little oxygen serve to reduce oxygen consumption in order to maintain a minimal internal oxygen concentration that is required for oxidative phosphorylation (i.e. activity of cytochrome oxidase) and sustainment of several metabolic pathways that require molecular oxygen (Geigenberger, 2003). Decreasing internal oxygen is sensed in plants and leads to an inhibition of respiration, a decrease of the ATP/ADP ratio, and the prioritization of metabolic pathways that conserve energy. Several components that control anaerobic responses have been identified (Licausi et al., 2010; Mustroph et al., 2010). EXL1 represents another factor that is essential for survival of anoxia (Fig. 9B).

## CONCLUSION

The phenotypic changes of *exl1* in soil and in synthetic medium only became evident when C and energy supply was suboptimal. Low sugar supply in synthetic medium, limited light and elongated dark periods in soil, and oxygen deficiency require comprehensive adaptations of metabolism and growth. EXL1 is essential for these adaptations. The mode of action of EXL1 is as yet unknown. The presumed extracellular localization of EXL1 suggests a specific role that is distinct from known sugar signaling pathways.

## MATERIALS AND METHODS

### Screen for Mutants and Establishment of Transgenic Lines

The SALK 027172 line (Alonso et al., 2003) carries a T-DNA insertion in the *EXL1* promoter and was named *exl1-D*. The GABI-Kat line GK 592A10 (Rosso et al., 2003) carries a T-DNA insertion in the *EXL1* coding sequence and was named *exl1*. DNA insertion sites were confirmed by sequencing. Homozygosity of both T-DNA insertion lines was confirmed by PCR on genomic DNA using T-DNA border-specific and gene-specific primers.

Several constructs were established using Gateway-compatible vectors. The *EXL1* coding sequence was amplified using the primers EXL1ga\_fw

(5'-CACCATGGCTTCTTTGTGATGGG-3') and EXL1ga\_rev (5'-AAACA-GAGTCGAGCAAGAATCTG-3'). The PCR fragments were cloned into the pENTR/D-TOPO (Invitrogen) entry vector. Sequencing confirmed full sequence identity of the PCR product with the ecotype Columbia (Col-0) genomic sequence. The *EXL1* coding sequence was inserted into the pH7WG2 vector for expression under control of the 35S promoter (Karimi et al., 2002) and into the pGWB14 vector (Nakagawa et al., 2007) for fusion of three copies of the HA epitope to the C terminus of EXL1. The resulting constructs were termed 35S::EXL1 and 35S::EXL1:HA, respectively. A genomic *EXL1* fragment comprising 928 bp upstream of the translation start codon and the coding sequence was amplified using the primers pEXL1ga\_fw (5'-CACCTT-GATTCGGTTTTTCGGATT-3') and EXL1ga\_rev. Cloning into the pGWB13 vector (Nakagawa et al., 2007) resulted in the pEXL1::EXL1:HA construct. All constructs were transformed into *Arabidopsis* (*Arabidopsis thaliana*) plants using the floral dip method.

## Growth Conditions

Seeds for growth experiments were derived from plants grown in parallel in a greenhouse. Plants were grown in half-concentrated MS medium supplemented with different Glc or Suc concentrations and solidified with 0.8% (w/v) agar. After 2 to 3 d in a cold room (4°C), plants were transferred into a growth chamber with a long-day light regime (16 h of light [ $140 \mu\text{mol m}^{-2} \text{s}^{-1}$ , 22°C]/8 h of dark [22°C]) and grown in a randomized manner. Alternatively, plants were established in soil. Seeds were allowed to germinate and grow for 2 weeks in controlled-growth chambers (7 d of 16 h of light [ $140 \mu\text{mol m}^{-2} \text{s}^{-1}$ , 20°C, 75% relative humidity]/8 h of dark [6°C, 75% relative humidity]; thereafter, 7 d of 8 h of light [ $140 \mu\text{mol m}^{-2} \text{s}^{-1}$ , 20°C, 60% relative humidity]/16 h of dark [16°C, 75% relative humidity]). Subsequently, plants were either left in short-day conditions or transferred to long-day conditions (Schröder et al., 2009), or plants were grown in a controlled-growth chamber under low total irradiance (4 h of light [ $60 \mu\text{mol m}^{-2} \text{s}^{-1}$ , 20°C, 60% relative humidity]/20 h of dark [16°C, 75% relative humidity]). All genotypes were grown in the same chamber at the same time in a randomized manner. Anaerobic responses were tested as described by Licausi et al. (2010).

## Gene Expression Analysis and Protein Extraction

Gene expression analyses were performed as described (Schröder et al., 2009). Primer sequences for quantitative RT-PCR were as follows: EXL1\_fw (5'-TGGATCGGCTTGTACTGGAG-3') and EXL1\_rev (5'-GGGTCAAACA-AAGCCGGTAA-3';  $E = 1.982 \pm 0.003$  [mean  $\pm$  se]); EXL1:HA\_fw (identical to EXL1\_fw) and EXL1:HA\_rev (5'-TCCTGCATAGTCCGGGACGTC-3';  $E = 1.983 \pm 0.005$ ); QQS\_fw (5'-AGCCATTGAAGAAACCTCTTTCG-3') and QQS\_rev (5'-ATGGCTGACCGTGTGAGTCTTG-3';  $E = 1.977 \pm 0.002$ ); and eIF1 $\alpha$ \_fw (5'-TTGACAGGCGTCTGTGTAAGG-3') and eIF1 $\alpha$ \_rev (5'-CAGCGTCACCATTCTTCAAAA-3'; At5g60390). Average PCR efficiencies ( $E$ ) were computed for each primer pair across all analyzed samples with the LinRegPCR software (Ramakers et al., 2003). Protein for western-blot analysis was isolated using ice-cold extraction buffer (125 mM Tris [pH 8.8], 1% SDS, 10% glycerol, 50 mM  $\text{Na}_2\text{SO}_4$ , and Complete Protease Inhibitor Cocktail [Roche]).

Sequence data from this article can be found in the GenBank/EMBL data libraries under accession numbers AY084831.1 (EXL1) and EU805808.1 (QQS).

## Supplemental Data

The following materials are available in the online version of this article.

**Supplemental File S1.** Affymetrix\_ATH1\_profiles\_SUPPLEMENT.xls.

**Supplemental File S2.** Biomass\_exl1-D\_SUPPLEMENT.xls.

## ACKNOWLEDGMENTS

We thank Justyna Sikocinska for implementation of sterile culture experiments and analyses of sugar levels. We thank Francesco Licausi and Joost van Dongen for assistance with the anoxia treatments.

Received April 4, 2011; accepted May 1, 2011; published May 4, 2011.

## LITERATURE CITED

- Ahuatzi D, Riera A, Peláez R, Herrero P, Moreno F (2007) Hxk2 regulates the phosphorylation state of Mig1 and therefore its nucleocytoplasmic distribution. *J Biol Chem* **282**: 4485–4493
- Alonso JM, Stepanova AN, Leisse TJ, Kim CJ, Chen H, Shinn P, Stevenson DK, Zimmerman J, Barajas P, Cheuk R, et al (2003) Genome-wide insertional mutagenesis of *Arabidopsis thaliana*. *Science* **301**: 653–657
- Arenas-Huertero F, Arroyo A, Zhou L, Sheen J, León P (2000) Analysis of *Arabidopsis* glucose insensitive mutants, *gin5* and *gin6*, reveals a central role of the plant hormone ABA in the regulation of plant vegetative development by sugar. *Genes Dev* **14**: 2085–2096
- Baena-González E, Rolland F, Thevelein JM, Sheen J (2007) A central integrator of transcription networks in plant stress and energy signaling. *Nature* **448**: 938–942
- Baena-González E, Sheen J (2008) Convergent energy and stress signaling. *Trends Plant Sci* **13**: 474–482
- Baier M, Hemmann G, Holman R, Corke F, Card R, Smith C, Rook F, Bevan MW (2004) Characterization of mutants in *Arabidopsis* showing increased sugar-specific gene expression, growth, and developmental responses. *Plant Physiol* **134**: 81–91
- Barker L, Kühn C, Weise A, Schulz A, Gebhardt C, Hirner B, Hellmann H, Schulze W, Ward JM, Frommer WB (2000) SUT2, a putative sucrose sensor in sieve elements. *Plant Cell* **12**: 1153–1164
- Bayer EM, Bottrill AR, Walshaw J, Vigouroux M, Naldrett MJ, Thomas CL, Maule AJ (2006) *Arabidopsis* cell wall proteome defined using multidimensional protein identification technology. *Proteomics* **6**: 301–311
- Bhalerao RP, Salchert K, Bakó L, Okrész L, Szabados L, Muranaka T, Machida Y, Schell J, Koncz C (1999) Regulatory interaction of PRL1 WD protein with *Arabidopsis* SNF1-like protein kinases. *Proc Natl Acad Sci USA* **96**: 5322–5327
- Bläsing OE, Gibon Y, Günther M, Höhne M, Morcuende R, Osuna D, Thimm O, Usadel B, Scheible W-R, Stitt M (2005) Sugars and circadian regulation make major contributions to the global regulation of diurnal gene expression in *Arabidopsis*. *Plant Cell* **17**: 3257–3281
- Borderies G, Jamet E, Lafitte C, Rossignol M, Jauneau A, Boudart G, Monsarrat B, Esquerré-Tugayé M-T, Boudet A, Pont-Lezica R (2003) Proteomics of loosely bound cell wall proteins of *Arabidopsis thaliana* cell suspension cultures: a critical analysis. *Electrophoresis* **24**: 3421–3432
- Caspar T, Huber SC, Somerville C (1985) Alterations in growth, photosynthesis, and respiration in a starchless mutant of *Arabidopsis thaliana* (L.) deficient in chloroplast phosphoglucomutase activity. *Plant Physiol* **79**: 11–17
- Chen J-G, Willard FS, Huang J, Liang J, Chasse SA, Jones AM, Siderovski DP (2003) A seven-transmembrane RGS protein that modulates plant cell proliferation. *Science* **301**: 1728–1731
- Cheng W-H, Endo A, Zhou L, Penney J, Chen H-C, Arroyo A, Leon P, Nambara E, Asami T, Seo M, et al (2002) A unique short-chain dehydrogenase/reductase in *Arabidopsis* glucose signaling and abscisic acid biosynthesis and functions. *Plant Cell* **14**: 2723–2743
- Choe S, Fujioka S, Noguchi T, Takatsuto S, Yoshida S, Feldmann KA (2001) Overexpression of *DWARF4* in the brassinosteroid biosynthetic pathway results in increased vegetative growth and seed yield in *Arabidopsis*. *Plant J* **26**: 573–582
- Coll-García D, Mazuch J, Altmann T, Müssig C (2004) EXORDIUM regulates brassinosteroid-responsive genes. *FEBS Lett* **563**: 82–86
- Contento AL, Kim S-J, Bassham DC (2004) Transcriptome profiling of the response of *Arabidopsis* suspension culture cells to Suc starvation. *Plant Physiol* **135**: 2330–2347
- Edgar R, Domrachev M, Lash AE (2002) Gene Expression Omnibus: NCBI gene expression and hybridization array data repository. *Nucleic Acids Res* **30**: 207–210
- Emanuelsson O, Nielsen H, Brunak S, von Heijne G (2000) Predicting subcellular localization of proteins based on their N-terminal amino acid sequence. *J Mol Biol* **300**: 1005–1016
- Farrar K, Evans IM, Topping JE, Souter MA, Nielsen JE, Lindsey K (2003) EXORDIUM: a gene expressed in proliferating cells and with a role in meristem function, identified by promoter trapping in *Arabidopsis*. *Plant J* **33**: 61–73
- Feiz L, Irshad M, Pont-Lezica RF, Canut H, Jamet E (2006) Evaluation of cell wall preparations for proteomics: a new procedure for purifying cell walls from *Arabidopsis* hypocotyls. *Plant Methods* **2**: 10
- Finn RD, Tate J, Mistry J, Coggill PC, Sammut SJ, Hotz H-R, Ceric G, Forslund K, Eddy SR, Sonnhammer ELL, et al (2008) The Pfam protein families database. *Nucleic Acids Res (Database issue)* **36**: D281–D288
- Gancedo JM (2008) The early steps of glucose signalling in yeast. *FEMS Microbiol Rev* **32**: 673–704
- Geigenberger P (2003) Response of plant metabolism to too little oxygen. *Curr Opin Plant Biol* **6**: 247–256
- Gibon Y, Bläsing OE, Palacios-Rojas N, Pankovic D, Hendriks JHM, Fisahn J, Höhne M, Günther M, Stitt M (2004) Adjustment of diurnal starch turnover to short days: depletion of sugar during the night leads to a temporary inhibition of carbohydrate utilization, accumulation of sugars and post-translational activation of ADP-glucose pyrophosphorylase in the following light period. *Plant J* **39**: 847–862
- Gibon Y, Pyl E-T, Sulpice R, Lunn JE, Höhne M, Günther M, Stitt M (2009) Adjustment of growth, starch turnover, protein content and central metabolism to a decrease of the carbon supply when *Arabidopsis* is grown in very short photoperiods. *Plant Cell Environ* **32**: 859–874
- Gibon Y, Usadel B, Bläsing OE, Kamlage B, Hoehne M, Trethewey R, Stitt M (2006) Integration of metabolite with transcript and enzyme activity profiling during diurnal cycles in *Arabidopsis* rosettes. *Genome Biol* **7**: R76
- Goda H, Sasaki E, Akiyama K, Maruyama-Nakashita A, Nakabayashi K, Li W, Ogawa M, Yamauchi Y, Preston J, Aoki K, et al (2008) The AtGenExpress hormone and chemical treatment data set: experimental design, data evaluation, model data analysis and data access. *Plant J* **55**: 526–542
- Goetz M, Godt DE, Roitsch T (2000) Tissue-specific induction of the mRNA for an extracellular invertase isoenzyme of tomato by brassinosteroids suggests a role for steroid hormones in assimilate partitioning. *Plant J* **22**: 515–522
- Grigston JC, Osuna D, Scheible W-R, Liu C, Stitt M, Jones AM (2008) D-Glucose sensing by a plasma membrane regulator of G signaling protein, AtRGS1. *FEBS Lett* **582**: 3577–3584
- Halford NG, Hey SJ (2009) Snf1-related protein kinases (SnRKs) act within an intricate network that links metabolic and stress signalling in plants. *Biochem J* **419**: 247–259
- Hey S, Mayerhofer H, Halford NG, Dickinson JR (2007) DNA sequences from *Arabidopsis*, which encode protein kinases and function as upstream regulators of Snf1 in yeast. *J Biol Chem* **282**: 10472–10479
- Ho S-L, Chao Y-C, Tong W-F, Yu S-M (2001) Sugar coordinately and differentially regulates growth- and stress-related gene expression via a complex signal transduction network and multiple control mechanisms. *Plant Physiol* **125**: 877–890
- Horton P, Park K-J, Obayashi T, Fujita N, Harada H, Adams-Collier CJ, Nakai K (2007) WoLF PSORT: protein localization predictor. *Nucleic Acids Res (Web Server issue)* **35**: W585–W587
- Huber SC, Hardin SC (2004) Numerous posttranslational modifications provide opportunities for the intricate regulation of metabolic enzymes at multiple levels. *Curr Opin Plant Biol* **7**: 318–322
- Huijser C, Kortstee A, Pego J, Weisbeek P, Wisman E, Smeekens S (2000) The *Arabidopsis* SUCROSE UNCOUPLED-6 gene is identical to *ABSCISIC ACID INSENSITIVE-4*: involvement of abscisic acid in sugar responses. *Plant J* **23**: 577–585
- Hummel I, Pantin F, Sulpice R, Piques M, Rolland G, Dauzat M, Christophe A, Pervent M, Bouteillé M, Stitt M, et al (2010) *Arabidopsis* plants acclimate to water deficit at low cost through changes of carbon usage: an integrated perspective using growth, metabolite, enzyme, and gene expression analysis. *Plant Physiol* **154**: 357–372
- Hruz T, Laule O, Szabo G, Wessendorp F, Bleuler S, Oertle L, Widmayer P, Gruissem W, Zimmermann P (2008) Genevestigator V3: a reference expression database for the meta-analysis of transcriptomics. *Adv Bioinform* **2008**: 420747
- Jamet E, Canut H, Boudart G, Pont-Lezica RF (2006) Cell wall proteins: a new insight through proteomics. *Trends Plant Sci* **11**: 33–39
- Jang J-C, León P, Zhou L, Sheen J (1997) Hexokinase as a sugar sensor in higher plants. *Plant Cell* **9**: 5–19
- Jossier M, Bouly J-P, Meimoun P, Arjmand A, Lessard P, Hawley S, Grahame Hardie D, Thomas M (2009) SnRK1 (SNF1-related kinase 1) has a central role in sugar and ABA signalling in *Arabidopsis thaliana*. *Plant J* **59**: 316–328
- Karimi M, Inzé D, Depicker A (2002) Gateway vectors for *Agrobacterium*-mediated plant transformation. *Trends Plant Sci* **7**: 193–195

- Karve A, Moore BD** (2009) Function of Arabidopsis hexokinase-like1 as a negative regulator of plant growth. *J Exp Bot* **60**: 4137–4149
- Koch K** (2004) Sucrose metabolism: regulatory mechanisms and pivotal roles in sugar sensing and plant development. *Curr Opin Plant Biol* **7**: 235–246
- Laby RJ, Kincaid MS, Kim D, Gibson SI** (2000) The Arabidopsis sugar-insensitive mutants *sis4* and *sis5* are defective in abscisic acid synthesis and response. *Plant J* **23**: 587–596
- Lee E-J, Koizumi N, Sano H** (2004) Identification of genes that are up-regulated in concert during sugar depletion in Arabidopsis. *Plant Cell Environ* **27**: 337–345
- Lee K-W, Chen P-W, Lu C-A, Chen S, Ho T-H, Yu S-M** (2009) Coordinated responses to oxygen and sugar deficiency allow rice seedlings to tolerate flooding. *Sci Signal* **2**: ra61
- Li L, Foster CM, Gan Q, Nettleton D, James MG, Myers AM, Wurtele ES** (2009) Identification of the novel protein QQS as a component of the starch metabolic network in Arabidopsis leaves. *Plant J* **58**: 485–498
- Li Y, Smith C, Corke F, Zheng L, Merali Z, Ryden P, Derbyshire P, Waldron K, Bevan MW** (2007) Signaling from an altered cell wall to the nucleus mediates sugar-responsive growth and development in *Arabidopsis thaliana*. *Plant Cell* **19**: 2500–2515
- Licausi F, van Dongen JT, Giuntoli B, Novi G, Santaniello A, Geigenberger P, Perata P** (2010) HRE1 and HRE2, two hypoxia-inducible ethylene response factors, affect anaerobic responses in *Arabidopsis thaliana*. *Plant J* **62**: 302–315
- Lisso J, Altmann T, Müssig C** (2006) Metabolic changes in fruits of the tomato *d<sup>r</sup>* mutant. *Phytochemistry* **67**: 2232–2238
- Lohse M, Nunes-Nesi A, Krüger P, Nagel A, Hannemann J, Giorgi FM, Childs L, Osorio S, Walther D, Selbig J, et al** (2010) Robin: an intuitive wizard application for R-based expression microarray quality assessment and analysis. *Plant Physiol* **153**: 642–651
- Lu C-A, Lin C-C, Lee K-W, Chen J-L, Huang L-F, Ho S-L, Liu H-J, Hsing Y-I, Yu S-M** (2007) The SnRK1A protein kinase plays a key role in sugar signaling during germination and seedling growth of rice. *Plant Cell* **19**: 2484–2499
- McGuffin LJ, Bryson K, Jones DT** (2000) The PSIPRED protein structure prediction server. *Bioinformatics* **16**: 404–405
- Michael TP, Mockler TC, Breton G, McEntee C, Byer A, Trout JD, Hazen SP, Shen R, Priest HD, Sullivan CM, et al** (2008) Network discovery pipeline elucidates conserved time-of-day-specific cis-regulatory modules. *PLoS Genet* **4**: e14
- Moore B, Zhou L, Rolland F, Hall Q, Cheng W-H, Liu Y-X, Hwang I, Jones T, Sheen J** (2003) Role of the Arabidopsis glucose sensor HXK1 in nutrient, light, and hormonal signaling. *Science* **300**: 332–336
- Moreno E, Ahuatzli D, Riera A, Palomino CA, Herrero P** (2005) Glucose sensing through the Hxk2-dependent signalling pathway. *Biochem Soc Trans* **33**: 265–268
- Müssig C** (2005) Brassinosteroid-promoted growth. *Plant Biol (Stuttg)* **7**: 110–117
- Mustroph A, Lee SC, Oosumi T, Zanetti ME, Yang H, Ma K, Yaghoubi-Masihi A, Fukao T, Bailey-Serres J** (2010) Cross-kingdom comparison of transcriptomic adjustments to low-oxygen stress highlights conserved and plant-specific responses. *Plant Physiol* **152**: 1484–1500
- Nakagawa T, Kurose T, Hino T, Tanaka K, Kawamukai M, Niwa Y, Toyooka K, Matsuoka K, Jinbo T, Kimura T** (2007) Development of series of Gateway binary vectors, pGWBs, for realizing efficient construction of fusion genes for plant transformation. *J Biosci Bioeng* **104**: 34–41
- Nam HK, Li J** (2002) BRI1/BAK1, a receptor kinase pair mediating brassinosteroid signalling. *Cell* **26**: 203–212
- Osuna D, Usadel B, Morcuende R, Gibon Y, Bläsing OE, Höhne M, Günter M, Kamlage B, Trethewey R, Scheible W-R, et al** (2007) Temporal responses of transcripts, enzyme activities and metabolites after adding sucrose to carbon-deprived Arabidopsis seedlings. *Plant J* **49**: 463–491
- Paul MJ, Pellny TK** (2003) Carbon metabolite feedback regulation of leaf photosynthesis and development. *J Exp Bot* **54**: 539–547
- Price J, Laxmi A, St Martin SK, Jang J-C** (2004) Global transcription profiling reveals multiple sugar signal transduction mechanisms in *Arabidopsis*. *Plant Cell* **16**: 2128–2150
- Ramakers C, Ruijter JM, Deprez RHL, Moorman AFM** (2003) Assumption-free analysis of quantitative real-time polymerase chain reaction (PCR) data. *Neurosci Lett* **339**: 62–66
- Roldán M, Gómez-Mena C, Ruiz-García L, Salinas J, Martínez-Zapater JM** (1999) Sucrose availability on the aerial part of the plant promotes morphogenesis and flowering of Arabidopsis in the dark. *Plant J* **20**: 581–590
- Rolland F, Baena-Gonzalez E, Sheen J** (2006) Sugar sensing and signaling in plants: conserved and novel mechanisms. *Annu Rev Plant Biol* **57**: 675–709
- Rosso MG, Li Y, Strizhov N, Reiss B, Dekker K, Weisshaar B** (2003) An *Arabidopsis thaliana* T-DNA mutagenized population (GABI-Kat) for flanking sequence tag-based reverse genetics. *Plant Mol Biol* **53**: 247–259
- Schaffer R, Landgraf J, Accerbi M, Simon V, Larson M, Wisman E** (2001) Microarray analysis of diurnal and circadian-regulated genes in *Arabidopsis*. *Plant Cell* **13**: 113–123
- Schlüter U, Köpke D, Altmann T, Müssig C** (2002) Analysis of carbohydrate metabolism of *CPD*-antisense plants and the brassinosteroid-deficient *cbf1* mutant. *Plant Cell Environ* **25**: 783–791
- Schröder F, Lisso J, Lange P, Müssig C** (2009) The extracellular EXO protein mediates cell expansion in Arabidopsis leaves. *BMC Plant Biol* **9**: 20
- Shen W, Reyes MI, Hanley-Bowdoin L** (2009) Arabidopsis protein kinases GRIK1 and GRIK2 specifically activate SnRK1 by phosphorylating its activation loop. *Plant Physiol* **150**: 996–1005
- Smeekens S, Ma J, Hanson J, Rolland F** (2010) Sugar signals and molecular networks controlling plant growth. *Curr Opin Plant Biol* **13**: 274–279
- Smith AM, Stitt M** (2007) Coordination of carbon supply and plant growth. *Plant Cell Environ* **30**: 1126–1149
- Smith SM, Fulton DC, Chia T, Thorncroft D, Chapple A, Dunstan H, Hylton C, Zeeman SC, Smith AM** (2004) Diurnal changes in the transcriptome encoding enzymes of starch metabolism provide evidence for both transcriptional and posttranscriptional regulation of starch metabolism in Arabidopsis leaves. *Plant Physiol* **136**: 2687–2699
- Stitt M, Gibon Y, Lunn JE, Piques M** (2007) Multilevel genomics analysis of carbon signalling during low carbon availability: coordinating the supply and utilisation of carbon in a fluctuating environment. *Funct Plant Biol* **34**: 526–549
- Sugden C, Donaghy PG, Halford NG, Hardie DG** (1999) Two SNF1-related protein kinases from spinach leaf phosphorylate and inactivate 3-hydroxy-3-methylglutaryl-coenzyme A reductase, nitrate reductase, and sucrose phosphate synthase in vitro. *Plant Physiol* **120**: 257–274
- Thimm O, Bläsing O, Gibon Y, Nagel A, Meyer S, Krüger P, Selbig J, Müller LA, Rhee SY, Stitt M** (2004) MAPMAN: a user-driven tool to display genomics data sets onto diagrams of metabolic pathways and other biological processes. *Plant J* **37**: 914–939
- Usadel B, Bläsing OE, Gibon Y, Retzlaff K, Höhne M, Günther M, Stitt M** (2008) Global transcript levels respond to small changes of the carbon status during progressive exhaustion of carbohydrates in Arabidopsis rosettes. *Plant Physiol* **146**: 1834–1861
- Wang Z-Y, Seto H, Fujioka S, Yoshida S, Chory J** (2001) BRI1 is a critical component of a plasma-membrane receptor for plant steroids. *Nature* **410**: 380–383
- Winter D, Vinegar B, Nahal H, Ammar R, Wilson GV, Provart NJ** (2007) An “Electronic Fluorescent Pictograph” browser for exploring and analyzing large-scale biological data sets. *PLoS ONE* **2**: e718
- Wu CY, Trieu A, Radhakrishnan P, Kwok SF, Harris S, Zhang K, Wang J, Wan J, Zhai H, Takatsuto S, et al** (2008) Brassinosteroids regulate grain filling in rice. *Plant Cell* **20**: 2130–2145
- Yamamoto YY, Obokata J** (2008) PPDB: a plant promoter database. *Nucleic Acids Res (Database issue)* **36**: D977–D981
- Yanagisawa S, Yoo SD, Sheen J** (2003) Differential regulation of EIN3 stability by glucose and ethylene signalling in plants. *Nature* **425**: 521–525
- Zhang J, Vemuri G, Nielsen J** (2010) Systems biology of energy homeostasis in yeast. *Curr Opin Microbiol* **13**: 382–388
- Zhang Y, Primavesi LF, Jhurrea D, Andralojc PJ, Mitchell RA, Powers SJ, Schlupepmann H, Delatte T, Winkler A, Paul MJ** (2009) Inhibition of SNF1-related protein kinase1 activity and regulation of metabolic pathways by trehalose-6-phosphate. *Plant Physiol* **149**: 1860–1871
- Zhou L, Jang J-C, Jones TL, Sheen J** (1998) Glucose and ethylene signal transduction crosstalk revealed by an Arabidopsis glucose-insensitive mutant. *Proc Natl Acad Sci USA* **95**: 10294–10299