

A Comparative Study of Ethylene Growth Response Kinetics in Eudicots and Monocots Reveals a Role for Gibberellin in Growth Inhibition and Recovery¹[W][OA]

Joonyup Kim, Rebecca L. Wilson, J. Brett Case², and Brad M. Binder*

Department of Biochemistry, Cellular, and Molecular Biology, University of Tennessee, Knoxville, Tennessee 37996

Time-lapse imaging of dark-grown *Arabidopsis thaliana* hypocotyls has revealed new aspects about ethylene signaling. This study expands upon these results by examining ethylene growth response kinetics of seedlings of several plant species. Although the response kinetics varied between the eudicots studied, all had prolonged growth inhibition for as long as ethylene was present. In contrast, with continued application of ethylene, white millet (*Panicum miliaceum*) seedlings had a rapid and transient growth inhibition response, rice (*Oryza sativa* 'Nipponbare') seedlings had a slow onset of growth stimulation, and barley (*Hordeum vulgare*) had a transient growth inhibition response followed, after a delay, by a prolonged inhibition response. Growth stimulation in rice correlated with a decrease in the levels of rice *ETHYLENE INSENSITIVE3-LIKE2* (*OsEIL2*) and an increase in rice *F-BOX DOMAIN AND LRR CONTAINING PROTEIN7* transcripts. The gibberellin (GA) biosynthesis inhibitor paclobutrazol caused millet seedlings to have a prolonged growth inhibition response when ethylene was applied. A transient ethylene growth inhibition response has previously been reported for *Arabidopsis ethylene insensitive3-1* (*ein3-1*) *eil1-1* double mutants. Paclobutrazol caused these mutants to have a prolonged response to ethylene, whereas constitutive GA signaling in this background eliminated ethylene responses. Sensitivity to paclobutrazol inversely correlated with the levels of EIN3 in *Arabidopsis*. Wild-type *Arabidopsis* seedlings treated with paclobutrazol and mutants deficient in GA levels or signaling had a delayed growth recovery after ethylene removal. It is interesting to note that ethylene caused alterations in gene expression that are predicted to increase GA levels in the *ein3-1* *eil1-1* seedlings. These results indicate that ethylene affects GA levels leading to modulation of ethylene growth inhibition kinetics.

Ethylene is a simple, unsaturated hydrocarbon that affects diverse processes throughout the lifetime of a plant, including seed germination, growth, formation of the apical hook, organ senescence, fruit ripening, abscission, gravitropism, and responses to various stresses (Mattoo and Suttle, 1991; Abeles et al., 1992). Of these many processes much attention has focused on the effects of ethylene on dark-grown seedlings. The Russian scientist Dimitry Neljubow discovered that ethylene is biologically active when he showed that ethylene was the active compound in illuminating gas that altered growth of etiolated pea (*Pisum sativum*) seedlings (Neljubow, 1901). The growth inhibition response from prolonged exposure to ethylene is a sensitive and easily quantified bioassay that has been used to screen for

mutants to acquire information and generate models about the ethylene signaling pathway (Bleecker et al., 1988; Guzmán and Ecker, 1990).

The model plant *Arabidopsis thaliana* contains five receptor isoforms localized to the membranes of the endoplasmic reticulum and Golgi apparatus that mediate responses to ethylene (Chang et al., 1993; Hua and Meyerowitz, 1998; Hua et al., 1998; Sakai et al., 1998; Chen et al., 2002, 2010; Dong et al., 2008; Grefen et al., 2008). The ethylene receptors form a complex with CONSTITUTIVE TRIPLE RESPONSE1 (CTR1; Clark et al., 1998; Cancel and Larsen, 2002; Gao et al., 2003; Huang et al., 2003; Mayerhofer et al., 2012). CTR1 negatively regulates the ethylene response pathway by either directly or indirectly inhibiting activity of ETHYLENE INSENSITIVE2 (EIN2; Kieber et al., 1993; Alonso et al., 1999; Huang et al., 2003). The EIN3, EIN3-LIKE1 (EIL1), and perhaps EIL2 transcription factors act downstream of EIN2 (Chao et al., 1997; Solano et al., 1998; Alonso et al., 2003b; Binder et al., 2004a, 2007). These, in turn, regulate other transcription factors such as ETHYLENE RESPONSE FACTOR1 (ERF1) and ETHYLENE RESPONSE DNA BINDING FACTOR1 to EDF4 (Chao et al., 1997; Solano et al., 1998; Alonso et al., 2003a; Stepanova et al., 2007). EIN3 and EIL1 are required for prolonged responses to ethylene but transient growth responses still occur in double *ein3* *eil1* loss-of-function mutants (Alonso et al., 2003b; Binder et al., 2004a). At least part of the

¹ This work was supported by the National Science Foundation (grant no. MCB-0918430 to B.M.B.).

² Present address: Department of Medicine, Vanderbilt University Medical Center, Nashville, TN 37232.

* Corresponding author; e-mail bbinder@utk.edu.

The author responsible for distribution of materials integral to the findings presented in this article in accordance with the policy described in the Instructions for Authors (www.plantphysiol.org) is: Brad M. Binder (bbinder@utk.edu).

[W] The online version of this article contains Web-only data.

[OA] Open Access articles can be viewed online without a subscription.

www.plantphysiol.org/cgi/doi/10.1104/pp.112.205799

regulation of EIN3 and EIL1 involves selective proteolysis via the ubiquitin/26S proteasome pathway using S-PHASE KINASE-ASSOCIATED1-CULLIN-F-BOX (SCF) E3 ubiquitin ligase complexes containing EIN3-BINDING F-BOX1 and EBF2 F-box proteins (Guo and Ecker, 2003; Potuschak et al., 2003; Yanagisawa et al., 2003; Gagne et al., 2004; Binder et al., 2007; An et al., 2010). This model posits that ethylene binding to the receptors reduces the activity of the receptors, leading to reduced activity of CTR1 and an increase in EIN2 protein levels along with subsequent signaling associated with it (Qiao et al., 2009). EIN2 acts in part to directly or indirectly reduce ubiquitination of EIN3 and EIL1 by SCF^{EBF1/2} leading to an increase in EIN3 and EIL1 levels and most ethylene responses. Similar genes have been described in other plant species such as rice (*Oryza sativa*), tomato (*Solanum lycopersicum*), and strawberry (*Fragaria vesca*; Rzewuski and Sauter, 2008; Ma et al., 2010; Klee and Giovannoni, 2011; Shulaev et al., 2011) suggesting similar signaling pathways are present across land plant species. Events downstream of transcriptional changes are more complicated and less understood. However, in cases where ethylene stimulates growth, ethylene treatment is modeled to cause an increase in GA levels and sensitivity leading to an increase in growth (Rijnders et al., 1997; Kende et al., 1998; Voesenek et al., 2003; Cox et al., 2004; Pierik et al., 2004, 2006). The role of GA in the ethylene growth inhibition response in dark-grown seedlings is less clear.

Although end-point analysis of seedling growth continues to be an informative bioassay, it has the limitation of only examining the long-term, end-point effects of ethylene on seedlings, so that transient events are overlooked. Time-lapse imaging has led to a more refined understanding about the ethylene receptors, feedback control mechanisms, signaling cross talk, and the roles that various proteins have in ethylene signaling in Arabidopsis (Binder et al., 2004a, 2004b, 2006, 2007; Potuschak et al., 2006; Gao et al., 2008; van Zanten et al., 2010; Vandenbussche et al., 2010; Žádníková et al., 2010; Kim et al., 2011; McDaniel and Binder, 2012). Even though many new details about ethylene signaling and responses have come out of studies using this approach, it is important to determine whether this information is specific to Arabidopsis or applicable to other plant species. With this in mind, we undertook a comparative study examining the growth response kinetics to the application and removal of ethylene in the shoots of several eudicot and monocot plant species.

Coupled with analyses of Arabidopsis mutants, these experiments revealed EIN3/EIL1-dependent and -independent regulation of GA levels, leading to modulation of both ethylene-induced growth inhibition kinetics and growth recovery kinetics after removal of ethylene.

RESULTS

Ethylene Growth Responses in Eudicots

In this study, Arabidopsis Columbia (Col) seedling hypocotyls grew in air with a rate of 0.32 ± 0.01 mm

h^{-1} that diminished to a steady-state rate of 0.04 ± 0.01 mm h^{-1} upon application of $10 \mu\text{L L}^{-1}$ ethylene (Supplemental Fig. S1). This is consistent with our prior results (Binder et al., 2004a, 2004b). We examined the rate of growth in air and ethylene for a variety of other eudicot plant species including kale (*Brassica oleracea* 'Red Russian'), canola (*Brassica napus*), two cultivars of tomato (*Solanum lycopersicum* 'Floradade' and 'German Queen'), poppy (*Papaver orientale* 'Oriental Scarlet'), and beetberry (*Chenopodium capitatum*). The growth rates in air for these species ranged from 0.15 ± 0.01 to 0.76 ± 0.06 mm h^{-1} . The growth rate of each of these eudicots was inhibited by the application of ethylene and ranged from 0.04 ± 0.01 to 0.20 ± 0.01 mm h^{-1} (Supplemental Fig. S1).

Our earlier studies on Arabidopsis seedlings uncovered two phases of hypocotyl growth inhibition when $10 \mu\text{L L}^{-1}$ ethylene was applied (Binder et al., 2004a, 2004b). The first phase of inhibition starts about 10 min after ethylene is added and lasts approximately 15 min when it reaches a new steady state. This new steady state lasts approximately 20 min when the second phase of growth inhibition occurs that further inhibits growth. A new steady-state growth rate is reached after approximately 20 min. This second, prolonged phase of growth inhibition requires the EIN3 and EIL1 transcription factors and lasts for as long as saturating concentrations of ethylene are present (Binder et al., 2004a, 2006, 2007). Similar onset kinetics are seen in both the Col and Wassilewskija ecotypes (Binder et al., 2004b). When ethylene is removed after a 2-h treatment, hypocotyl growth recovers to pretreatment rates in approximately 85 min in Col and 120 min in Wassilewskija seedlings. A dampened oscillation in growth rate often occurs during this recovery phase that involves a growth rate overshoot where the growth rate increases over the rate initially observed prior to ethylene treatment (Binder et al., 2004b, 2007). The growth response kinetics of Col hypocotyls to a 2-h application of $10 \mu\text{L L}^{-1}$ ethylene (Fig. 1) were indistinguishable from our previous results (Binder et al., 2004a, 2006, 2007).

We examined kale and canola as two other members of the Brassicaceae family. The onset of the second phase of growth inhibition was less distinct in these species than Arabidopsis (Fig. 1A). Additionally, after the removal of ethylene, kale and canola seedlings had a growth rate overshoot that continued to increase rather than the dampened oscillation in growth rate observed with Arabidopsis. By 5 h after the removal of ethylene, the growth rates of both species reached a plateau but had not yet started to return to pretreatment growth rates (data not shown). It is possible that this is part of a dampened oscillation in growth rate that had a longer period beyond the time frame of these experiments. The shoots of both cultivars of tomato showed a measurable decrease in growth rate approximately 10 min after application of ethylene (Fig. 1B). cv German Queen reached steady-state growth inhibition within approximately 60 min whereas cv Floradade reached maximum growth inhibition approximately




Figure 1. Ethylene growth response kinetics in dark-grown shoots of various eudicots. Seedlings were grown in air 1 h before $10 \mu\text{L L}^{-1}$ ethylene was applied (downward arrow). Ethylene was removed 2 h later (upward arrow), and seedlings allowed to grow another 3 h. The growth rate was normalized to the growth rate in the 1-h air pretreatment. A, Arabidopsis, canola, and kale. B, Two cultivars of tomato. C, Poppy and beetberry. In B and C, responses of Arabidopsis seedlings are shown in gray for comparison. Data represent the average \pm SEM of at least five seedlings.

90 min after ethylene was added. Both took approximately 1 h longer than Col to recover to pretreatment growth rates after the removal of ethylene. To gain a better understanding about the variability of ethylene growth response kinetics in eudicots, we examined ethylene growth responses in poppy and beetberry to represent two other plant families (Fig. 1C). Poppy seedling shoots showed growth inhibition kinetics very similar to cv German Queen seedlings (Fig. 1, B and C). However, after the removal of ethylene, poppy had growth recovery kinetics that were more similar to Col seedlings with one difference being the lack of a growth rate overshoot. In contrast, beetberry seedling shoots

showed ethylene growth inhibition kinetics very similar to Col seedlings. Of the eudicots observed, beetberry seedlings had the shortest delay in the onset of growth recovery and the fastest rate of growth recovery after ethylene removal (Fig. 1C; Supplemental Table S1). These seedlings also had a growth rate overshoot that rose to approximately 2.5-fold over the growth rate initially observed in air before treatment with ethylene (Fig. 1C).

These results show that eudicots vary in their ethylene growth response and recovery kinetics even within the same species. The underlying mechanisms for these differences are unknown. From the above data, five growth parameters were measured including growth rate in air, steady-state growth rate in ethylene, time to maximum growth inhibition after addition of ethylene, delay in onset of recovery after ethylene removal, and recovery time after ethylene removal (Supplemental Table S1). To determine if any of these parameters might be mechanistically linked, we compared them pairwise and calculated the correlation coefficient (r^2). Most have an r^2 value well below 0.5 indicating a weaker correlation between the traits (Supplemental Table S2). The steady-state growth rate in air correlated with the steady-state growth rate in ethylene with an $r^2 = 0.51$ whereas the delay in the onset of growth recovery correlated with the recovery time with an $r^2 = 0.49$. These low correlations suggest that these traits are not tightly linked mechanistically.

Ethylene Growth Responses in Monocots

To gain more insights into the action of ethylene, we examined the ethylene rapid growth response kinetics of the coleoptiles of seedlings from rice 'Nipponbare', white millet (*Panicum miliaceum*), and barley (*Hordeum vulgare*). In air, under these conditions, rice coleoptiles grew at $0.16 \pm 0.02 \text{ mm h}^{-1}$, millet coleoptiles at $0.44 \pm 0.05 \text{ mm h}^{-1}$, and barley coleoptiles at $0.71 \pm 0.03 \text{ mm h}^{-1}$. Application of $10 \mu\text{L L}^{-1}$ ethylene inhibited coleoptile growth in millet, although the growth rate began to recover prior to ethylene removal 2 h later (data not shown). This prompted us to examine the growth response kinetics of millet with prolonged exposure to ethylene. For comparison, we also measured the growth inhibition kinetics of Arabidopsis seedlings with prolonged ethylene treatment. Consistent with our prior research (Binder et al., 2004a, 2006), Col seedling hypocotyls had prolonged growth inhibition in the continued presence of $10 \mu\text{L L}^{-1}$ ethylene (Fig. 2). Ethylene treatment of millet seedlings resulted in a transient growth inhibition response that corresponded in timing to the first phase of growth inhibition observed in Arabidopsis hypocotyls (Fig. 2). Millet showed maximum growth inhibition approximately 20 min after ethylene was added. This growth rate stabilized for approximately 30 min at $0.19 \pm 0.04 \text{ mm h}^{-1}$ after which the growth rate increased, returning to pretreatment growth rates approximately 90 min after ethylene was added.




Figure 2. Ethylene growth response kinetics in dark-grown shoots of several monocots. Seedlings were grown in air 1 h before $10 \mu\text{L L}^{-1}$ ethylene was applied (arrow). The growth rate was normalized to the growth rate in the 1-h air pretreatment. The growth response kinetics of Arabidopsis are shown for comparison. Data represent the average \pm SEM of at least five seedlings.

We obtained very different growth response kinetics with rice seedlings. Application of ethylene caused no measurable growth inhibition of rice coleoptiles (Fig. 2). Approximately 70 min after the addition of ethylene, the growth rate started to increase, reaching a maximum about 3.5 h after ethylene was applied. The maximum growth rate was approximately double the air pretreatment growth rate. This growth stimulation response to ethylene is consistent with prior studies on rice (Ku et al., 1970; Suge, 1972, 1985; Suge et al., 1971; Raskin and Kende, 1984; Rose-John and Kende, 1985; Satler and Kende, 1985; Keith et al., 1986; Hoffmann-Benning and Kende, 1992; Watanabe et al., 2007). It is interesting to note that whereas adult cv Nipponbare rice is considered submergence intolerant, at the seedling stage this cultivar is flooding tolerant and has increased coleoptile growth upon flooding (Lee et al., 2009).

In contrast to millet and rice, application of ethylene to barley resulted in a small, transient growth inhibition response that was followed, after a delay, with a prolonged growth inhibition response (Fig. 2). These results show that the ethylene growth response kinetics in these monocots are more variable and distinct from what are seen in the eudicots studied.

Ethylene Alters Transcript Levels of *OsEIL2* and *OsFBL7* in Rice Coleoptiles

The EIN3 and EIL1 transcription factors are required for long-term growth inhibition responses to ethylene in Arabidopsis (Chao et al., 1997; Solano et al., 1998; Alonso et al., 2003b; Binder et al., 2004a, 2007). Because of this, we were interested to examine the levels of the homologous genes, *OsEIL1* and *OsEIL2*, from rice. These genes have been linked to ethylene responses in rice (Mao et al., 2006; Hiraga et al., 2009). In Arabidopsis, growth inhibition caused by application of ethylene correlates with increases in EIN3 and EIL1 levels (Gao et al.,

2003; Yanagisawa et al., 2003; Gagne et al., 2004). In rice, coleoptile growth is stimulated by ethylene. We therefore predicted that the transcript levels for *OsEIL1* or *OsEIL2* or both would decrease in coleoptiles treated with ethylene. The levels of these transcripts relative to *OsUBIQUITIN5* (*OsUBQ5*) were analyzed using quantitative real-time reverse transcriptase (qRT)-PCR. This revealed that in coleoptiles of dark-grown rice seedlings maintained in air, the transcript levels of *OsEIL2* were approximately 2.5-fold higher than transcript levels of *OsEIL1* (Fig. 3). Upon treatment with $10 \mu\text{L L}^{-1}$ ethylene for 3 h, the *OsEIL2* transcript levels decreased 50% but no significant change ($P = 0.4$) was observed in the transcript levels of *OsEIL1* (Fig. 4). Thus, ethylene treatment may lead to lower levels of *OsEIL2* because of decreased transcript abundance.

In Arabidopsis, EIN3 and EIL1 levels are controlled by SCF E3 complexes containing the EBF1 and EBF2 F-box proteins (Guo and Ecker, 2003; Potuschak et al., 2003; Yanagisawa et al., 2003; Gagne et al., 2004; Binder et al., 2007; An et al., 2010). The rice genome contains two related genes, rice *F-BOX DOMAIN AND LRR CONTAINING PROTEIN7* (*OsFBL7*) and *OsFBL30* (Wang et al., 2009). We therefore examined the transcript levels of these two genes in coleoptiles of dark-grown rice seedlings maintained in air versus $10 \mu\text{L L}^{-1}$ ethylene for 3 h. We found that in air, *OsFBL7* transcript levels were approximately 24-fold higher than *OsFBL30* (Fig. 3). Application of ethylene caused *OsFBL7* transcript levels to increase but no significant alterations ($P = 0.13$) in *OsFBL30* transcript levels occurred. These observations suggest that application of ethylene results in lower levels of *OsEIL1* and *OsEIL2* due to higher levels of the *OsFBL7* F-box protein, resulting in higher rates of ubiquitination and proteolysis.




Figure 3. Transcript levels for select ethylene signaling genes in rice coleoptiles. The levels of transcript for *OsEIL1*, *OsEIL2*, *OsFBL7*, and *OsFBL30* were normalized relative to *OsUBQ5*. Seedlings were grown in the dark in air for 3 d, RNA was extracted from coleoptiles, and transcript levels were analyzed with qRT-PCR. Transcript levels in untreated seedlings were compared with seedlings treated with $10 \mu\text{L L}^{-1}$ ethylene for 3 h before RNA extraction. Data show the average \pm SEM for three biological replicates with two to three technical replicates each. *Statistical difference compared with untreated seedlings ($P \leq 0.05$).




Figure 4. Effects of paclobutrazol on growth in air and ethylene growth response kinetics. Seedlings of millet, Col, *ein3-1 eil1-1* double mutants, and *ebf2-3* single mutants were treated with paclobutrazol. A, Paclobutrazol dose dependency on growth in air. The responses of millet shoots at several doses of paclobutrazol are shown for comparison. Data represent the average \pm SEM of at least four seedlings. B, Effects of paclobutrazol on ethylene growth response kinetics of millet. C, Effects of paclobutrazol and paclobutrazol plus GA on *ein3-1 eil1-1* mutants. In B and C, seedlings were grown 1 h in air prior to application of $10 \mu\text{L L}^{-1}$ ethylene (arrow). Data represent the average \pm SEM of at least five seedlings.

Paclobutrazol Prolongs the Ethylene Growth Inhibition Response of Millet Seedlings

A model has developed that posits that when rice is submerged, ethylene levels increase that in turn lead to an increase in GA levels, causing increased growth of aerial tissues (Kende et al., 1998; Bailey-Serres and Voeselek, 2008). Similarly, GA is modeled to be involved

in ethylene stimulated growth of Arabidopsis hypocotyls that occurs under specific growth conditions (Vandenbussche et al., 2007). This raises the possibility that the transient growth response in millet seedlings is caused by an increase in GA levels in ethylene-treated seedlings that counteracts the growth inhibition initially caused by application of ethylene. Inhibition of GA biosynthesis has previously been shown to reduce flood-induced and ethylene-stimulated growth of rice (Raskin and Kende, 1984; Setter and Laureles, 1996; Das et al., 2005; Hattori et al., 2009). Therefore, to test this hypothesis, we germinated millet seedlings on paclobutrazol-containing medium to block GA biosynthesis. Both 30 and 120 μM paclobutrazol reduced the growth rate of millet coleoptiles in air approximately 30% (Fig. 4A) from $0.39 \pm 0.02 \text{ mm h}^{-1}$ to $0.27 \pm 0.01 \text{ mm h}^{-1}$. Dimethyl sulfoxide (DMSO), solvent-control seedlings had a transient growth inhibition response (Fig. 4B) similar to what we observed above in the absence of DMSO (Fig. 2). Millet seedlings treated with 30 μM paclobutrazol still had a transient growth inhibition response, but took longer than control seedlings to recover to pretreatment growth rates whereas 120 μM paclobutrazol blocked the growth rate reversal so that the growth rate remained inhibited while ethylene was applied (Fig. 4B). Thus, GA appears to be important for the reversal in growth inhibition when millet seedlings are treated with ethylene.

Paclobutrazol Prolongs and Constitutive GA Signaling Attenuates the Ethylene Growth Inhibition Response of *ein3 eil1* Seedlings

Unlike wild-type Col seedlings, *ein3-1 eil1-1* double mutant seedlings only have a transient growth inhibition response to the application of ethylene (Binder et al., 2004a, 2007) resembling the growth kinetics observed with millet (this study). Therefore, we treated *ein3-1 eil1-1* double mutant seedlings with paclobutrazol to determine whether GA biosynthesis is involved in the growth response kinetics of this mutant. The *ein3-1 eil1-1* double mutants were more responsive to paclobutrazol than millet (Fig. 4A). Also, consistent with prior studies on Arabidopsis (Jacobsen and Olszewski, 1993; Peng et al., 1999; Wen and Chang, 2002), *ein3-1 eil1-1* seeds failed to germinate well on paclobutrazol (data not shown). Therefore, *ein3-1 eil1-1* seeds were germinated on media without paclobutrazol and then gently transferred onto plates containing paclobutrazol or DMSO and allowed to grow in air for 3 h prior to time-lapse imaging. Solvent control *ein3-1 eil1-1* double mutants had a transient growth inhibition response with complete growth recovery taking approximately 2 h after the addition of ethylene (Fig. 4C) that was similar to our prior results with this double mutant (Binder et al., 2004a, 2007). Application of 30 μM paclobutrazol reduced the growth rate of these seedlings from $0.50 \pm 0.01 \text{ mm h}^{-1}$ to $0.10 \pm 0.004 \text{ mm h}^{-1}$ and completely blocked the growth reversal of these double mutants, so that ethylene caused prolonged growth inhibition (Fig. 4C). To

determine whether the effect of paclobutrazol on the *ein3-1 eil1-1* mutants was caused by GA, we treated the double mutants with both 30 μM paclobutrazol and 10 μM GA. The presence of GA slightly increased the growth rate in air to $0.13 \pm 0.005 \text{ mm h}^{-1}$ and restored the transient response to ethylene that was eliminated by paclobutrazol (Fig. 4C).

To confirm that GA is involved in reversing the first phase of growth inhibition, we compared the ethylene growth inhibition kinetics of *ein3-1 eil1-1* mutants with *ein3-1 eil1-1 della* mutants where all five *DELLA* genes have also been mutated (Feng et al., 2008; An et al., 2012). The quintuple *della* mutant seedlings have a constitutive GA response phenotype and hypocotyl growth in dark-grown *della* seedlings is insensitive to application of either GA or paclobutrazol (Feng et al., 2008). In air, the growth rate of the double mutants was indistinguishable from the septuple mutants (data not shown). The *ein3-1 eil1-1 della* mutants had a very attenuated or no growth inhibition response when ethylene was added compared with *ein3-1 eil1-1* mutants (Fig. 5).

Together, these data suggest that GA is involved in regulating the long-term growth inhibition response to ethylene. One possibility is that GA and ethylene are regulating long-term growth via parallel pathways. Alternatively, EIN3 and EIL1 could be negatively regulating GA levels, so that when ethylene levels are high, EIN3 and EIL1 levels increase and GA biosynthesis is reduced or catabolism is increased leading to reduced GA levels and long-term growth inhibition. The second hypothesis predicts that wild-type Arabidopsis seedlings should be more sensitive to paclobutrazol than the *ein3-1 eil1-1* mutants because EIN3 and EIL1 in wild-type seedlings would also be negatively regulating GA levels. Consistent with the second hypothesis, Col seedlings grew slower than *ein3-1 eil1-1* mutants in air (0.32 ± 0.01 versus $0.50 \pm 0.01 \text{ mm h}^{-1}$ respectively) and were also slightly more sensitive to paclobutrazol than the double mutants (Fig. 4A). We also examined the effects of paclobutrazol on an EIN3 overexpression line (EIN3-GFP) that has previously been shown to have a constitutive ethylene response phenotype (Guo and Ecker, 2003). Whereas paclobutrazol inhibited growth of these seedlings, we could not obtain clear dose-response measurements because these seedlings grew very slowly ($0.09 \pm 0.01 \text{ mm h}^{-1}$) in the absence of paclobutrazol (data not shown). Therefore, we examined *ebf2-3* mutants that have previously been shown to accumulate EIN3 protein to higher levels than wild-type seedlings (Gagne et al., 2004). Whereas *ebf2-3* mutants grew at a similar rate in air as Col seedlings, the *ebf2-3* seedlings were more sensitive to paclobutrazol than Col seedlings (Fig. 4A) providing more support for the idea that EIN3 and EIL1 negatively regulate GA levels.

Altered GA Levels or Signaling Affect Ethylene Growth Response Kinetics in Arabidopsis

These results prompted us to investigate whether GA is involved in regulating the ethylene growth kinetic




Figure 5. Constitutive GA signaling attenuates the ethylene growth inhibition response of *ein3 eil1* seedlings. Seedlings were grown 1 h in air prior to application of 10 $\mu\text{L L}^{-1}$ ethylene (arrow). The growth response of *ein3-1 eil1-1* versus *ein3-1 eil1-1 della* mutants are compared. Data represent the average \pm SEM of at least 11 seedlings.

responses of wild-type Arabidopsis seedlings. Addition of 1 μM GA had no obvious effect on the growth rate in air or ethylene growth response kinetics (data not shown). This is consistent with prior results showing that addition of GA had little effect on hypocotyl elongation in the dark suggesting that GA levels are saturated (Cowling and Harber, 1999; Feng et al., 2008). Treating Col seedlings with 5 μM paclobutrazol had no measurable effect on the inhibition kinetics upon addition of 10 $\mu\text{L L}^{-1}$ ethylene (Fig. 6A), although the growth rate in air was reduced from $0.29 \pm 0.01 \text{ mm h}^{-1}$ to $0.15 \pm 0.01 \text{ mm h}^{-1}$ when paclobutrazol was added. In contrast, the paclobutrazol-treated seedlings had altered growth recovery kinetics after the removal of ethylene. It is most notable that the paclobutrazol-treated seedlings failed to completely recover to pre-treatment growth rates in the time scale of these measurements. These results indicate that GA levels may also modulate normal growth recovery after the removal of ethylene. To further address the role of GA in modulating these responses when EIN3 and EIL1 are present, we examined the ethylene growth response kinetics of *ga3-oxidase1-3* (*ga3ox1-3*) that is mutated in the *GA3ox1* gene and produces less GA (Mitchum et al., 2006). In air, the *ga3ox1-3* mutants grew at $0.28 \text{ mm} \pm 0.01 \text{ mm h}^{-1}$, which is slightly slower than the Col wild-type controls ($P < 0.05$). The *ga3ox1-3* seedlings had similar growth inhibition kinetics compared with the wild type, but appeared to reach a slower growth rate during the plateau in growth prior to the onset of the second phase of growth inhibition response (Fig. 6B). As expected, based upon the results obtained with paclobutrazol-treated seedlings, these mutants also recovered more slowly after the removal of ethylene. We examined another mutant of *GA3ox1* (*ga4-1*) that produces less GA (Talon et al., 1990; Chiang et al., 1995; Cowling et al., 1998) and obtained similar results (data not shown). Because reduced GA levels altered the ethylene




Figure 6. Altered GA levels or signaling modulate ethylene growth response kinetics. Seedlings were then grown for 1 h in air, followed by the application of $10 \mu\text{L L}^{-1}$ ethylene (downward arrow). Ethylene was removed 2 h later (upward arrow). A, Col seedlings were treated with $5 \mu\text{M}$ paclobutrazol or DMSO as a solvent control for 3 h prior to time-lapse imaging. B, *ga3ox1-3* compared with Col controls. Data for Col were taken from Figure 2. C, *gai-1* and *della* mutants were compared with *Ler* controls. Data represent the average \pm SEM of at least eight seedlings.

growth response, we also examined the ethylene growth response kinetics of the *ga insensitive-1* (*gai-1*) mutant that has reduced GA responses (Koorneef et al., 1985; Peng and Harberd, 1993, 1997; Wilson and Somerville, 1995; Peng et al., 1997). In air, hypocotyls of *gai-1* seedlings grew at approximately 50% the rate of *Landsberg erecta* (*Ler*) control seedlings (0.14 ± 0.01 versus $0.27 \pm 0.01 \text{ mm h}^{-1}$ respectively) consistent with prior studies

(Cowling and Harberd, 1999). Hypocotyls of *Ler* seedlings had growth inhibition kinetics similar to Col seedlings but took longer to recover after ethylene was removed (Fig. 6C; Supplemental Table S1). Like the *ga3ox1-3* mutants, the *gai-1* mutants appeared to reach a slower growth rate during the plateau in growth inhibition prior to the onset of the second phase of growth inhibition response (Fig. 6C). In addition, the mutants had slower growth recovery after removal of ethylene and never fully recovered to pretreatment growth rates in the timescale of these experiments (Fig. 6C). It is interesting to note that in ethylene the *gai-1* mutants reached a steady-state growth rate of $0.02 \pm 0.002 \text{ mm h}^{-1}$, which is significantly slower ($P < 0.001$) than *Ler* seedlings that grew at $0.04 \pm 0.004 \text{ mm h}^{-1}$.

We also examined quintuple *della* mutant seedlings where all five *DELLA* genes are mutated, and seedlings have a constitutive GA response phenotype (Feng et al., 2008). The hypocotyl growth rate of *della* seedlings in air was similar to wild-type seedlings consistent with prior research (Feng et al., 2008). The ethylene growth inhibition kinetics of the *della* seedlings were similar to the wild type except that they seemed to reach a slightly faster growth rate during the plateau in growth prior to the onset of the second phase of growth inhibition response (Fig. 6C). The *della* mutant seedlings had similar recovery kinetics to the wild type after the removal of ethylene, except that the mutants had a growth rate overshoot that was missing in the *Ler* controls (Fig. 6C). Both of these alterations are opposite to what we observed in the *gai3ox1-3*, *ga4-1*, and *gai-1* mutants that have reduced GA levels and signaling (Fig. 6, B and C).

Transcript Levels for GA Metabolism Genes Are Altered in *ein3 eil1* Seedlings

From the above, we were interested to know whether altered levels of EIN3 correlated with altered GA biosynthesis or degradation. To address this, RNA was extracted from 2-d-old Col, *ebf2-3*, and *ein3-1 eil1-1* seedlings that were grown in the dark in hydrocarbon-free air. Some seedlings were treated with $1 \mu\text{L L}^{-1}$ ethylene for 2 h prior to RNA extraction. We used qRT-PCR to compare the transcript levels of select genes related to GA metabolism. We focused on several GA biosynthesis genes (*ent*-kaurene synthase [*KS*], *GA20ox1*–*GA20ox3*, and *GA3ox1*) and one gene coding for a GA degradation enzyme (*GA2ox2*). The levels of these transcripts relative to *ACTIN2* (*ACT2*) were normalized to levels in untreated, Col seedlings (Fig. 7).

In air, we found that the transcript levels for *KS* increased in the *ebf2-3* background relative to Col seedlings, whereas the transcript levels for the other genes did not change. A different pattern was observed in transcripts from *ein3-1 eil1-1* seedlings where the level of *KS* transcript was unchanged but the levels of *GA2ox2*, *GA3ox1*, *GA20ox1*, *GA20ox2*, and *GA20ox3* transcripts increased (Fig. 7). Therefore, the levels of

these transcripts in dark-grown seedlings maintained in air are affected by mutations that alter EIN3 levels.

When Col seedlings were treated with $1 \mu\text{L L}^{-1}$ ethylene for 2 h and examined for changes in gene expression, we observed increases in the transcript levels of *KS* and *GA3ox1* and decreases in the levels of *GA2ox2*, *GA2ox1*, and *GA2ox3* transcripts. The levels of *KS* transcript were similar between ethylene-treated Col and untreated *ebf2-3* mutants. The pattern of ethylene-induced changes was different in the *ebf2-3* mutants where application of ethylene led to a slight increase in the level of *GA2ox3* transcripts. Other apparent differences in the *ebf2-3* mutants caused by ethylene were considered statistically insignificant (using a $P \leq 0.05$). Surprisingly, application of ethylene to the *ein3-1 eil1-1* double mutants caused a decrease in *GA2ox2* and an increase in *GA3ox1* transcript levels. Thus, ethylene may cause an increase in GA levels in the *ein3-1 eil1-1* double mutants that is predicted to increase the growth rate of these seedlings.

It is also interesting to note that after treatment with ethylene, the *ein3-1 eil1-1* double mutants had lower levels of *KS* transcript but higher levels of *GA3ox1*, *GA2ox1*, *GA2ox2*, and *GA2ox3* compared with ethylene-treated Col that would likely result in higher GA levels in the double mutants compared with Col. By contrast, ethylene treatment of the *ebf2-3* mutants resulted in an increase in the level of *GA2ox2* transcript and a smaller increase in *GA2ox3* transcripts compared with ethylene-treated Col. These differences would potentially lead to lower levels of GA in the *ebf2-3* mutants and perhaps underlie the delay in growth recovery after removal of ethylene that has been observed in these mutants (Binder et al., 2007). These results show that ethylene affects GA metabolism genes both via EIN3/EIL1 and independently of these transcription factors.

DISCUSSION

It is believed that all land plants bind and respond to ethylene. However, the range and number of responses in dark-grown seedlings differs between species (Goeschl, 1975; Abeles et al., 1992). We used high-resolution, time-lapse imaging of dark-grown seedlings to compare the effects of ethylene on the growth of shoots from several eudicot and monocot species. All eudicots studied here had a growth inhibition response with a delay of between 10 and 15 min before the onset of growth inhibition upon application of $10 \mu\text{L L}^{-1}$ ethylene. Similar latent times for growth inhibition with ethylene have generally been observed in other eudicot species using other techniques (Warner and Leopold, 1971; Burg, 1973; Goeschl, 1975; Rauser and Horton, 1975; Jackson, 1983). Whereas the delay for growth inhibition was fairly constant between the eudicots studied, the time for full growth inhibition after ethylene was added and the time for full growth recovery after ethylene was removed were both more variable and differed by over 1 h between species.




Figure 7. Transcript levels for select GA metabolism genes in Arabidopsis seedlings. The levels of transcript for *KS*, *GA2ox2*, *GA3ox1*, *GA2ox1*, *GA2ox2*, and *GA2ox3* relative to *ACT2* in *ebf2-3* and *ein3-1 eil1-1* mutants were normalized to levels in untreated Col seedlings. Seedlings were grown in the dark in air for 2 d, RNA was extracted, and the transcript was analyzed with qRT-PCR. Transcript levels in untreated seedlings (closed bars) were compared with seedlings treated with $1 \mu\text{L L}^{-1}$ ethylene for 2 h before RNA extraction (open bars). Data shows the average \pm SEM for three biological replicates with two to three technical replicates each. ^aStatistical difference compared with untreated Col ($P \leq 0.05$); ^bstatistical difference caused by treatment with ethylene compared with untreated seedlings of that background ($P \leq 0.05$); ^cstatistical difference caused by ethylene compared with ethylene-treated Col seedlings ($P \leq 0.05$).

Similarly, the delay in the onset of growth recover after removal of ethylene varied up to approximately 50 min between the different eudicots. It remains to be seen whether the ethylene growth kinetics of shoots from other eudicots are similar to those observed in this study. The three monocots studied, millet, rice, and barley were more diverged in their growth kinetic responses to ethylene. Barley seedlings had growth inhibition kinetics that were somewhat similar to the eudicots, except that the initial growth inhibition response was attenuated, and there was a delay prior to prolonged growth inhibition. In contrast, millet seedlings had a rapid growth inhibition response when ethylene was added. However, whereas eudicots had a second phase of growth inhibition that was slower in onset and more prolonged, millet had a transient growth response to the continued application of

10 $\mu\text{L L}^{-1}$ ethylene. The underlying mechanism for this first, transient growth inhibition response is unknown but in *Arabidopsis* occurs independently of EIN3 and EIL1 (Binder et al., 2004a). The transient response in millet is similar to our prior results with wild-type *Arabidopsis* seedlings treated with low, subsaturating dosages ($\leq 0.01 \mu\text{L L}^{-1}$) of ethylene (Binder et al., 2004a). In this study, we used 1,000-fold higher ethylene concentrations suggesting that the transient growth response in millet is not caused by subsaturating dosages of ethylene. Rice seedlings lacked a growth inhibition response. Rather, rice seedlings had a slow onset of growth stimulation after ethylene was added. It is interesting to note that phase II growth inhibition in *Arabidopsis* is predicted to have a similar delay in onset and time for maximum response (Binder, 2007) as the growth stimulation response in rice (compared in Supplemental Fig. S2), indicating that the two responses may have common molecular components even though ethylene is having opposite effects.

Prior studies have shown that when ethylene treatment leads to stimulation of growth, GA levels increase to mediate this growth stimulation (Kende et al., 1998; Voesenek et al., 2003; Vandenbussche et al., 2007). We therefore explored the role of GA in ethylene growth response kinetics. Treatment of millet seedlings with paclobutrazol to block GA biosynthesis attenuated or prevented the growth reversal seen during continuous ethylene treatment. In other words, growth inhibition caused by application of ethylene was prolonged by paclobutrazol treatment. This suggests that ethylene treatment of millet leads to an increase in GA biosynthesis that in turn leads to an increase in the growth rate. In the case of millet, this overcomes the initial growth inhibition effect of ethylene whereas in rice there is no initial growth inhibition response, so ethylene only causes a stimulation of growth. Some links between ethylene and GA metabolism have been uncovered in light-grown seedlings (Achard et al., 2007; Vandenbussche et al., 2007). However, the role for GA metabolism in the ethylene growth inhibition response in the shoots of dark-grown eudicot seedlings remains largely unexplored. Whereas GA has been linked to growth stimulation caused by ethylene, here we show that regulation of GA levels also has a role in modulating growth inhibition caused by application of ethylene and recovery after ethylene removal. GA stimulates hypocotyl elongation in eudicots (Sun, 2008), which suggests a model where ethylene causes a decrease in GA levels in dark-grown eudicot seedlings to affect the second phase of growth inhibition. Support for this model comes from the observation that ethylene treatment causes a decrease in GA content in sunflowers (*Helianthus annuus*; Pearce et al., 1991), and *ctr1* mutants in *Arabidopsis* with constitutive ethylene signaling have decreased GA content (Achard et al., 2007).

In *Arabidopsis*, the prolonged growth inhibition response to ethylene is controlled by EIN3, EIL1, and probably EIL2 (Chao et al., 1997; Binder et al., 2004a, 2007). EIN3 levels rise when *Arabidopsis* is treated with ethylene (Guo and Ecker, 2003; Yanagisawa et al., 2003;

Gagne et al., 2004). Also, EIN3 overexpression in *Arabidopsis* leads to slower growth in air and slower recovery after ethylene is removed (Guo and Ecker, 2003; Binder et al., 2007). These observations are consistent with lower GA levels. It is interesting to note that mutants deficient in GA signaling or levels also recover slower after removal of ethylene. In contrast, *ein3-1 eil1-1* loss-of-function mutants grow faster in air than wild-type plants and lack a prolonged growth inhibition response when ethylene is added (Binder et al., 2004a, 2007; this study) consistent with higher GA levels. This supports a model where ethylene-induced activation of the EIN3 and EIL transcription factors in eudicots leads to reduced GA levels to modulate growth inhibition (Fig. 8). Consistent with this, we found that sensitivity to paclobutrazol was inversely correlated to the levels of EIN3 indicating that paclobutrazol and EIN3 inhibit GA accumulation via independent mechanisms. Whereas ethylene altered the expression levels of genes for GA metabolic enzymes in Col seedlings, ethylene probably also modulates GA levels independently of EIN3 and EIL1 because ethylene treatment of *ein3-1 eil1-1* seedlings resulted in decreased levels of *GA2ox2* transcript and increased levels of *GA3ox1* transcript. Both changes are predicted to lead to an increase in GA content. It is interesting to note that paclobutrazol prolonged the growth response of *ein3-1 eil1-1* double mutants whereas coapplication of GA with paclobutrazol did not, suggesting that increased GA content in this double mutant leads to the transient ethylene growth inhibition response. It is also interesting to note that the *ein3-1 eil1-1 della* mutants had a highly attenuated growth inhibition response compared with *ein3-1 eil1-1* mutants. Because *della* mutants have a constitutive GA response phenotype, this further supports a role for GA signaling in modulating the growth inhibition response to ethylene.

We have previously proposed two models invoking feedback mechanisms to explain the two phases of growth inhibition and the transient nature of the first phase of growth inhibition in *Arabidopsis* (Binder et al., 2004a). In both of these models, we proposed dual functions for EIN2. The results from this study suggest a third possible model involving GA (Fig. 8). In this model, EIN2 levels are low, so there are no ethylene responses and growth is fast; growth rate is modulated by feedback control of GA levels where higher growth rate leads to lower GA levels. Ethylene treatment of *Arabidopsis* seedlings leads to an initial growth inhibition response via EIN2 that is independent of EIN3 and EIL1. This is modeled to be independent of GA. After a delay, the reduced growth leads to an increase in GA levels or signaling that overcomes the initial growth inhibition response caused by EIN2. Also after a delay, EBF1 and EBF2 activity is inhibited resulting in higher levels of EIN3 and EIL1. In this model, we propose that this leads to decreased levels of GA resulting in the second, prolonged phase of growth inhibition. There also is a GA-independent inhibition of growth by EIN3 and EIL1. This model predicts that when EIN3 and EIL1




Figure 8. Models for modulation of ethylene growth responses by GA in Arabidopsis, rice, and millet. In Arabidopsis, this model proposes that in the absence of ethylene, EIN2 levels are low so there are no ethylene responses and growth is fast. Growth rate is modulated by feedback control of GA levels where a higher growth rate leads to lower levels. Ethylene leads to an accumulation of EIN2. One output of EIN2 reduces growth via an EIN3/EIL1-independent mechanism. On a slower time course, this reduced growth leads to an increase in GA levels resulting in an increase in growth. This pathway is proposed to underlie the transient first phase of growth inhibition response. On a slower time course, EIN2 also inhibits degradation of EIN3/EIL1 mediated by the EBF1 and EBF2 F-box proteins. Accumulation of EIN3 and EIL1 causes a lower growth rate leading to the second phase of prolonged growth inhibition response. Our results indicate that this is occurring via a GA-dependent pathway where EIN3 and EIL1 inhibit GA accumulation. There also is likely to be a GA-independent pathway. Thus, when EIN3 and EIL1 are not present, ethylene initially causes growth inhibition, but this inhibition is reversed by the delayed feedback involving GA. The model predicts that in rice, OsEIN2 activation by ethylene has no growth inhibition effect. On a slower time scale, activation of OsEIN2 leads to enhanced degradation of OsEIL1 and OsEIL2 mediated by the OsFBL7 and OsFBL30 F-box proteins. This leads to lower levels of OsEIL1 and OsEIL2, resulting in higher levels of GA and enhanced growth. In millet, the model proposes that PmEILs are either absent or at very low levels. Thus, PmEIN2 activity initially leads to growth inhibition, but feedback resulting in higher GA levels reverses this inhibition leading to a transient growth response.

are removed, ethylene activation of EIN2 initially causes growth inhibition, but after a delay the feedback loop causes an increase in GA levels that overcomes this initial growth inhibition response leading to a transient growth inhibition response. Addition of paclobutrazol blocks or reduces this increase in GA, leading to prolonged growth inhibition. Similarly, mutants with reduced GA levels or signaling have a stronger growth inhibition response when ethylene is added and take longer to recover after removal of ethylene. This model also provides a framework to understand the variety of growth inhibition responses in the eudicots studied, as well as barley, where the timing and relative strengths of the proposed outputs from EIN2 and feedback varies between species.

This basic model needs to be modified to explain the results obtained with rice and millet. Six *EIL* genes have been identified in the rice genome (Mao et al., 2006). *OsEIL1* and *OsEIL2* are the most closely related to *EIN3*, *EIL1*, and *EIL2* from Arabidopsis and have been linked to ethylene signaling in rice (Mao et al., 2006; Hiraga et al., 2009). Overexpression of *OsEIL1* in rice leads to decreased growth (Mao et al., 2006) much like overexpression of *EIN3* in Arabidopsis decreases growth (Guo and Ecker, 2003). Thus, these transcription factors have similar roles in Arabidopsis and rice, yet ethylene has the opposite effect on the growth of shoots of these species under the conditions used in this study. If these differences in the growth effects of ethylene are mediated by the EILs, it is likely that the levels of the EILs are controlled differently in these two species when ethylene is applied. Consistent with this, we found that ethylene caused a decrease in the mRNA levels of *OsEIL2* in coleoptiles correlating with the growth stimulation caused by application of ethylene. Also, application of ethylene enhanced transcript levels of *OsFBL7*. Increased levels of this F-box protein would lead to decreased levels of *OsEIL1* and *OsEIL2*. The Arabidopsis homologs of *OsFBL7* and *OsFBL30* are *EBF2* and *EBF1*, respectively (Rzewuski and Sauter, 2008; Ma et al., 2010). *EBF2* appears to have a larger role in long-term responses to ethylene than *EBF1* (Binder et al., 2007), paralleling the results in this study suggesting that *FBL7* may have a larger role than *FBL30* in the slow onset of growth stimulation caused by the application of ethylene. Several ethylene response factors have been identified that act downstream of the *OsEILs* to mediate ethylene responses in rice (Xu et al., 2000, 2006; Yang et al., 2002; Fukao and Bailey-Serres, 2008; Hu et al., 2008; Fukao et al., 2009, 2011; Hattori et al., 2009; Jung et al., 2010; Qi et al., 2011). At least some of these ethylene response factors act to modulate GA levels. The results from these prior studies coupled with results in this study indicate that the output from EIN2 in Arabidopsis that causes a rapid growth inhibition response is lacking from *OsEIN2* in dark-grown cv Nipponbare rice coleoptiles (Fig. 8). Rather, increased activity from *OsEIN2* leads to increases in *OsFBL7*, causing increased proteolysis of *OsEIL1* and *OsEIL2*. The lower levels of these transcription factors are predicted to lead to increased GA biosynthesis and increased growth (Fig. 8). We cannot rule out that there is also an *OsEIL1/OsEIL2*-independent mechanism that leads to increased GA levels. Whereas ethylene stimulates coleoptile growth of dark-grown rice seedlings, it inhibits growth of roots (Ma et al., 2010). The differences in mechanisms by which ethylene causes growth stimulation versus inhibition has yet to be resolved. Arabidopsis *EIN3* has been shown to be affected by factors other than ethylene (Yanagisawa et al., 2003; Lee et al., 2006; Zhu et al., 2011; An et al., 2012), suggesting that *EIN3* and the *EILs* are part of a complex regulatory network that could underlie these opposite responses to ethylene. Although we have no data on the *PmEILs* from millet, we predict that *PmEIL* levels are low and ethylene causes little or no change in their levels (Fig.

8). Therefore, ethylene is predicted to initially cause growth inhibition via output from PmEIN2. On a slower timescale, this inhibition is counteracted by a feedback mechanism similar to what is predicted to occur in Arabidopsis.

Based upon the limited number of species studied, the shoots of eudicot species appear to have diverged only slightly in the kinetics of growth inhibition upon addition of ethylene. In contrast, shoots of monocots have evolved more divergent of ethylene growth responses. Even though the number of ethylene receptor isoforms can vary between species and receptor function within a species is not entirely overlapping (Binder et al., 2012), it is likely that much of this divergence in ethylene growth responses arises from differences in downstream signaling components. Our data indicate that the control of the EIN3 and EIL transcription factors by ethylene is one locus that differs between plant species and under different environmental and developmental conditions. This leads to differences in GA levels that affect growth responses to ethylene.

MATERIALS AND METHODS

A variety of eudicot plant species were used in this study including kale (*Brassica oleracea* 'Red Russian'), canola (*Brassica napus*), two cultivars of tomato (*Solanum lycopersicum* 'Floradade' and 'German Queen'), poppy (*Papaver orientale* 'Oriental Scarlet'), and beetberry (*Chenopodium capitatum*). Three monocots species, rice (*Oryza sativa* 'Nipponbare'), white millet (*Panicum miliaceum*), and barley (*Hordeum vulgare*) were also examined. Seeds for these different species were obtained from various sources (Supplemental Table S3). These species were chosen both to represent a variety of plant families as well as because of their small seed size, which made them compatible with the time-lapse imaging methods used with Arabidopsis (*Arabidopsis thaliana*) seedlings. The *ein3-1 eil1-1* double mutant was originally obtained from Joseph Ecker and has previously been characterized (Alonso et al., 2003b; Binder et al., 2004a). The *ga4-1*, *ga3ox1-3*, and *gai-1* mutants were from the Arabidopsis Biological Resource Center and have previously been characterized (Talon et al., 1990; Jacobsen and Olszewski, 1993; Chiang et al., 1995; Jacobsen et al., 1996; Mitchum et al., 2006), the quintuple *della* mutant seeds where all five *DELLA* genes are mutated (*rga-t2 gai-t6 rgl1-1 rgl2-1 rgl3-1*) were obtained from Xiuhua Gao and Xiangdong Fu (Feng et al., 2008), the EIN3-GFP over-expressing seeds and *ein3 eil1 della* septuple mutants were from Hongwei Guo (Guo and Ecker, 2003; An et al., 2012). The *ebf2-3* loss-of-function mutant has been previously described and characterized (Gagne et al., 2004; Binder et al., 2007). The *ein3-1 eil1-1*, *ebf2-3*, and *ga3ox1-3* mutants are in the Col background whereas the *ga4-1*, *gai-1*, and *della* mutants are in the *Ler* background. Gibberellic acid was from Fisher Scientific, and paclobutrazol was a gift from Elena Shpak (University of Tennessee-Knoxville).

Seed Preparation

Most seeds were surface-sterilized by treatment with 70% alcohol for 30 s. Rice and barley seeds were surface-sterilized with a solution containing 10% (v/v) bleach with 0.01% (v/v) Triton X-100 for 5 min prior to treatment with 70% (v/v) alcohol. Unless otherwise specified, surface sterilized seeds were placed on sterile filter paper to dry and then placed on agar plates containing 0.8% (w/v) agar, one-half-strength Murashige and Skoog basal salt mixture, (Murashige and Skoog, 1962), pH 5.7, fortified with vitamins and with no added sugar. Arabidopsis, kale, and canola seeds were cold treated at 4°C for 2 to 4 d prior to being light treated for 2 to 8 h at 21°C. Tomato seeds were plated and light treated without cold treatment. To ensure germination in the dark, the *ga3ox1-3*, *ga4-1*, *gai-1*, and *della* mutants were soaked in sterile, distilled water with 1 μ M GA at 4°C in the dark for 1 d prior to being placed on agar plates and cold treated at 4°C for 2 d prior to being light treated. The *ga3ox1-3* and *ga4-1* seeds were light treated for 4 h, and *gai-1* and *della* seeds for 12 to 24 h at 21°C. Light-treated seeds were then allowed to germinate in the dark at 21°C. Millet, poppy, and beetberry seeds were plated and allowed to

germinate in the dark at 21°C without light treatment, and rice seeds were germinated in the dark at 28°C and then transferred to 21°C for 2 h prior to time-lapse imaging.

Paclobutrazol and GA Treatment of Seedlings

In some cases, seedlings were treated with paclobutrazol to block GA biosynthesis (Dalziel and Lawrence, 1984; Hedden and Graebe, 1985). Paclobutrazol was made as a 1,000 \times stock in DMSO, filter sterilized, and added to the agar after heat sterilization to the indicated final concentration. Millet seedlings were germinated on paclobutrazol-containing plates whereas solvent control millet seedlings were germinated and grown on agar plates containing DMSO. However, paclobutrazol inhibited Arabidopsis seed germination as has been previously reported (Jacobsen and Olszewski, 1993; Peng et al., 1999; Wen and Chang, 2002). Therefore, we germinated Arabidopsis seeds on media without paclobutrazol, then gently transferred the seedlings onto plates containing the indicated concentration of paclobutrazol (or DMSO as a control) and allowed them to grow in air 3 h prior to time-lapse imaging. We have previously shown that this transfer protocol has no measurable effect on the ethylene growth inhibition response (Binder et al., 2006). These manipulations were done under dim green light. In some experiments, GA was also included in the agar at the indicated concentration.

Time-Lapse Imaging and Analyses

Seedlings were allowed to germinate in the dark until approximately 4 to 8 mm long as previously described (Binder et al., 2004a, 2004b, 2006). Images of growing shoots were captured in the dark with Marlin CCD cameras (Allied Vision Technology) using infrared radiation and at 21°C. All manipulations prior to imaging were done under dim green light.

Growth rate experiments used two different protocols. In one, seedlings were allowed to grow 1 h in air followed by 2 h with 10 μ L L⁻¹ ethylene. Ethylene was then removed and seedlings allowed to grow an additional 3 h in air to allow for growth recovery. Alternatively, seedlings were grown in air for 1 h followed by 10 μ L L⁻¹ ethylene treatment for 5 h. In both cases, images were captured every 5 min. To determine the growth rate of shoots, the height in pixels of each seedling in each frame was analyzed using custom software written by Edgar Spalding (Parks and Spalding, 1999; Folta and Spalding, 2001) as previously described (Binder et al., 2004a, 2004b). Growth rates in air were determined from the 1 h of air pretreatment prior to applying ethylene. Growth rates were normalized to the growth rate during the air pretreatment for each seedling to make response kinetic comparisons easier, and trend lines were drawn by hand based upon moving averages of two consecutive data points. To calculate the growth rate in ethylene, the average growth rate was determined after the growth rate had stabilized at a new rate after ethylene application.

RNA Isolation and qRT-PCR

The transcript levels of several Arabidopsis GA biosynthesis-related genes including: *K5*, *GA3ox1*, *GA2ox2*, *GA20ox1*, *GA20ox2*, and *GA20ox3* were analyzed using qRT-PCR. Similar analysis was used to determine the transcript levels of *OsEIL1*, *OsEIL2*, *OsFBL7*, and *OsFBL30* from rice. For qRT-PCR analysis of Arabidopsis genes, total RNA was extracted from 80 or more 2-d-old, dark-grown Arabidopsis seedlings using a TRIzol Reagent (Invitrogen). Some seedlings were treated with 1 μ L L⁻¹ ethylene for 2 h prior to RNA extraction. For qRT-PCR analysis of rice genes, total RNA was extracted from five coleoptiles of 3-d-old, dark-grown seedlings. Some seedlings were treated with 10 μ L L⁻¹ ethylene for 3 h prior to extractions. The quality and quantity of the extracted RNAs were examined on a 1.2% agarose gel and by using a Nanodrop (ND-1000) spectrophotometer (Thermo Fisher Scientific). One microgram of total RNA was used for complementary DNA (cDNA) synthesis using the ImProm-II Reverse Transcription System (Promega) according to the manufacturer's instructions. The RNA was treated with one unit of DNase I (Invitrogen) prior to cDNA synthesis, and the cDNAs were diluted 10-fold. The qRT-PCR was performed with a Bio-Rad iQ5 Real-Time PCR Detection System (Bio-Rad) using SsoFast EvaGreen Supermix (Bio-Rad). Gene-specific primers (see below) were used in a total volume of 10 μ L with 3 μ L of cDNA, 0.5 μ M of gene-specific primers and 4 μ L of SsoFast EvaGreen Supermix (Bio-Rad). The qRT-PCR conditions were: 45 cycles consisting of 95°C for 15 s, 58°C for 30 s, 72°C for 10 s followed by the initial denaturation step at 95°C for 1 min. All qRT-PCR values are from three biological replicates with two to three technical replicates each. Data represents the average \pm SEM.

Arabidopsis transcript data were normalized to *ACT2* as an internal control. The cycle threshold values for the *ACT2* reference gene varied by no more than 1 between treatments and plant backgrounds. The primers used for qRT-PCR were: *KS* 5'-AATCAGCTCTACAAGCTCGTGAG-3' (forward), *KS* 5'-ATTCCTTGGGAACCACTTCCTT-3' (reverse), *GA20x2* 5'-CTGTGTG-AAAGATGGAAGTTGGGT-3' (forward), *GA20x2* 5'-ATCTTGCTCAGGGA-CAAGGCAT-3' (reverse); *GA30x1* 5'-TCGCCTCTCAACGATTCCGT-3' (forward), *GA30x1* 5'-ATTTAGCTGGAGAGCAGCTTGG-3' (reverse); *GA200x1* 5'-CATTCTGCCAACCCCAA-3' (forward), *GA200x1* 5'-TGATGCTGTCCAAAAGCTCTCT-3' (reverse); *GA200x2* 5'-TCGTTGCAATATTGG-TGACACTTC-3' (forward), *GA200x2* 5'-TCITCGGACACAAGAAAACCGC-3' (reverse); *GA200x3* 5'-TCGTGGTGAACATAGGCGACA-3' (forward), *GA200x3* 5'-CGTTACTAGTCTCTCGTGGCTTC-3' (reverse). *ACT2* primers were from Cartieaux et al. (2008). Data were normalized to transcript levels in untreated Col seedlings. Rice transcript data were normalized *OsUBQ5*. The cycle threshold values for the *OsUBQ5* reference gene varied by no more than 1 between treatments. The primers used were: *OsFBL30* 5'-ACTTGAACGGCTGCGAGAA-3' (forward), 5'-ATCCTGCTGCAACCTTCGAGA-3' (reverse); *OsFBL7* 5'-AGA-TAACCGATGCTAGCCTCTTT-3' (forward), 5'-CAGGGTAGCAACACCATTG-TCA-3' (reverse). Primers for *OsEIL1* and *OsEIL2* were from Hiraga et al. (2009) and for *OsUBQ5* was from Jain et al. (2006).

Ethylene Concentration Measurements

Ethylene concentrations were determined using a Hewlett-Packard 6890 gas chromatograph with an HP Plot/Q column (Agilent Technologies) or an ETD-300 ethylene detector (Sensor Sense).

Statistical Analysis

Five growth parameters (growth rate in air, growth rate in ethylene, time to maximum growth inhibition after addition of ethylene, delay in the onset of growth recovery after ethylene removal, recovery time after ethylene removal) were measured in this study (Supplemental Table S1) and compared pairwise. The square of r^2 was determined using Microsoft Excel. These values are shown in Supplemental Table S2. qRT-PCR data were analyzed with Student's *t* tests and considered statistically different with a *P* value of ≤ 0.05 .

Arabidopsis Genome Initiative accession numbers for Arabidopsis genes described in this article are: *EBF1*, At2g25490; *EBF2*, At5g25350; *EIN3*, At3g20770; *EIL1*, At2g27050; *EIL2*, At5g21120; *CTR1*, At5g03730; *EIN2*, At5g03280; *KS*, At1g79460; *GA30x1* (*GA4*), At1g15550; *GA20x2*, At1g30040; *GA200x1*, At4g25420; *GA200x2*, At5g51810; *GA200x3*, At5g07200; *GAL*, At1g14920. The Gene Loci for rice genes described in this article are: *OsEIL1*, Os03g20780; *OsEIL2*, Os07g48630; *OsFBL7*, Os02g10700; *OsFBL30*, Os06g40360.

Supplemental Data

The following materials are available in the online version of this article.

Supplemental Figure S1. Growth rates of dark-grown eudicot seedling shoots in air and ethylene.

Supplemental Figure S2. Comparison of ethylene growth response kinetics of rice shoots and predicted phase II growth inhibition kinetics of Arabidopsis.

Supplemental Table S1. Growth parameter measurements for eudicots studied.

Supplemental Table S2. Correlations between growth parameters in eudicots.

Supplemental Table S3. List of species used and source for seeds.

ACKNOWLEDGMENTS

The authors thank Joe Williams, Simon Gilroy and members of the Binder lab for helpful discussions, and Amanda Wehner for technical assistance. We thank Joseph Ecker, Xiangdong Fu, Hongwei Guo, Harry Klee, Sara Patterson, Amber Smith, and Albrecht von Arnim who kindly supplied us with seeds.

Received August 19, 2012; accepted September 12, 2012; published September 13, 2012.

LITERATURE CITED

- Abeles F, Morgan P, Saltveit MJ (1992) Ethylene in Plant Biology, Ed 2. Academic Press, San Diego, CA
- Achard P, Baghour M, Chapple A, Hedden P, van Der Straeten D, Genschik P, Moritz T, Harberd NP (2007) The plant stress hormone ethylene controls floral transition via DELLA-dependent regulation of floral meristem-identity genes. *Proc Natl Acad Sci USA* **104**: 6484–6489
- Alonso JM, Hirayama T, Roman G, Nourizadeh S, Ecker JR (1999) EIN2, a bifunctional transducer of ethylene and stress responses in Arabidopsis. *Science* **284**: 2148–2152
- Alonso JM, Stepanova AN, Leisse TJ, Kim CJ, Chen HM, Shinn P, Stevenson DK, Zimmerman J, Barajas P, Cheuk R, et al (2003a) Genome-wide insertional mutagenesis of Arabidopsis thaliana. *Science* **301**: 653–657
- Alonso JM, Stepanova AN, Solano R, Wisman E, Ferrari S, Ausubel FM, Ecker JR (2003b) Five components of the ethylene-response pathway identified in a screen for weak ethylene-insensitive mutants in Arabidopsis. *Proc Natl Acad Sci USA* **100**: 2992–2997
- An F, Zhang X, Zhu Z, Ji Y, He W, Jiang Z, Li M, Guo H (2012) Coordinated regulation of apical hook development by gibberellins and ethylene in etiolated Arabidopsis seedlings. *Cell Res* **22**: 915–927
- An F, Zhao Q, Ji Y, Li W, Jiang Z, Yu X, Zhang C, Han Y, He W, Liu Y, et al (2010) Ethylene-induced stabilization of ETHYLENE INSENSITIVE3 and EIN3-LIKE1 is mediated by proteasomal degradation of EIN3 binding F-box 1 and 2 that requires EIN2 in Arabidopsis. *Plant Cell* **22**: 2384–2401
- Bailey-Serres J, Voeselek LACJ (2008) Flooding stress: acclimations and genetic diversity. *Annu Rev Plant Biol* **59**: 313–339
- Binder BM (2007) Rapid kinetic analysis of ethylene growth responses in seedlings: new insights into ethylene signal transduction. *J Plant Growth Regul* **26**: 131–142
- Binder BM, Chang C, Schaller GE (2012) Perception of ethylene by plants: ethylene receptors. In *McManus MT*, ed, Annual Plant Reviews, The Plant Hormone Ethylene, Vol 44. Wiley-Blackwell, Hoboken, NJ, pp 117–145
- Binder BM, Mortimore LA, Stepanova AN, Ecker JR, Bleeker AB (2004a) Short-term growth responses to ethylene in Arabidopsis seedlings are EIN3/EIL1 independent. *Plant Physiol* **136**: 2921–2927
- Binder BM, O'Malley RC, Wang W, Moore JM, Parks BM, Spalding EP, Bleeker AB (2004b) Arabidopsis seedling growth response and recovery to ethylene. A kinetic analysis. *Plant Physiol* **136**: 2913–2920
- Binder BM, O'Malley RC, Wang W, Zutz TC, Bleeker AB (2006) Ethylene stimulates mutations that are dependent on the ETR1 receptor. *Plant Physiol* **142**: 1690–1700
- Binder BM, Walker JM, Gagne JM, Emborg TJ, Hemmann G, Bleeker AB, Vierstra RD (2007) The Arabidopsis EIN3 binding F-Box proteins EBF1 and EBF2 have distinct but overlapping roles in ethylene signaling. *Plant Cell* **19**: 509–523
- Bleeker AB, Estelle MA, Somerville C, Kende H (1988) Insensitivity to ethylene conferred by a dominant mutation in Arabidopsis thaliana. *Science* **241**: 1086–1089
- Burg SP (1973) Ethylene in plant growth. *Proc Natl Acad Sci USA* **70**: 591–597
- Cancel JD, Larsen PB (2002) Loss-of-function mutations in the ethylene receptor ETR1 cause enhanced sensitivity and exaggerated response to ethylene in Arabidopsis. *Plant Physiol* **129**: 1557–1567
- Cartieaux F, Contesto C, Gallou A, Desbrosses G, Kopka J, Taconnat L, Renou J-P, Touraine B (2008) Simultaneous interaction of Arabidopsis thaliana with Bradyrhizobium Sp. strain ORS278 and Pseudomonas syringae pv. tomato DC3000 leads to complex transcriptome changes. *Mol Plant Microbe Interact* **21**: 244–259
- Chang C, Kwok SF, Bleeker AB, Meyerowitz EM (1993) Arabidopsis ethylene-response gene ETR1: similarity of product to two-component regulators. *Science* **262**: 539–544
- Chao QM, Rothenberg M, Solano R, Roman G, Terzaghi W, Ecker JR (1997) Activation of the ethylene gas response pathway in Arabidopsis by the nuclear protein ETHYLENE-INSENSITIVE3 and related proteins. *Cell* **89**: 1133–1144
- Chen Y-F, Gao Z, Kerris RJ III, Wang W, Binder BM, Schaller GE (2010) Ethylene receptors function as components of high-molecular-mass protein complexes in Arabidopsis. *PLoS ONE* **5**: e8640.
- Chen Y-F, Randlett MD, Findell JL, Schaller GE (2002) Localization of the ethylene receptor ETR1 to the endoplasmic reticulum of Arabidopsis. *J Biol Chem* **277**: 19861–19866

- Chiang HH, Hwang I, Goodman HM (1995) Isolation of the *Arabidopsis* GA4 locus. *Plant Cell* 7: 195–201
- Clark KL, Larsen PB, Wang XX, Chang C (1998) Association of the *Arabidopsis* CTR1 Raf-like kinase with the ETR1 and ERS ethylene receptors. *Proc Natl Acad Sci USA* 95: 5401–5406
- Cowling RJ, Harberd NP (1999) Gibberellins control *Arabidopsis* hypocotyl growth via regulation of cellular elongation. *J Exp Bot* 50: 1351–1357
- Cowling RJ, Kamiya Y, Seto H, Harberd NP (1998) Gibberellin dose-response regulation of GA4 gene transcript levels in *Arabidopsis*. *Plant Physiol* 117: 1195–1203
- Cox MCH, Benschop JJ, Vreeburg RAM, Wagemaker CAM, Moritz T, Peeters AJM, Voeselek LACJ (2004) The roles of ethylene, auxin, abscisic acid, and gibberellin in the hyponastic growth of submerged *Rumex palustris* petioles. *Plant Physiol* 136: 2948–2960
- Dalziel J, Lawrence DK (1984) Biochemical and biological effects of kaurene oxidase inhibitors, such as paclobutrazol. In Menhenett R, Lawrence DK, eds, *Biochemical Aspects of Synthetic and Naturally Occurring Plant Growth Regulators*, Monograph No. 11. British Plant Growth Regulator Group, Wantage, UK, pp 43–57
- Das KK, Sarkar RK, Ismail AM (2005) Elongation ability and non-structural carbohydrate levels in relation to submergence tolerance in rice. *Plant Sci* 168: 131–136
- Dong C-H, Rivarola M, Resnick JS, Maggin BD, Chang C (2008) Subcellular co-localization of *Arabidopsis* RTE1 and ETR1 supports a regulatory role for RTE1 in ETR1 ethylene signaling. *Plant J* 53: 275–286
- Feng S, Martinez C, Gusmaroli G, Wang Y, Zhou J, Wang F, Chen L, Yu L, Iglesias-Pedraz JM, Kircher S, et al (2008) Coordinated regulation of *Arabidopsis thaliana* development by light and gibberellins. *Nature* 451: 475–479
- Folta KM, Spalding EP (2001) Unexpected roles for cryptochrome 2 and phototropin revealed by high-resolution analysis of blue light-mediated hypocotyl growth inhibition. *Plant J* 26: 471–478
- Fukao T, Bailey-Serres J (2008) Submergence tolerance conferred by Sub1A is mediated by SLR1 and SLRL1 restriction of gibberellin responses in rice. *Proc Natl Acad Sci USA* 105: 16814–16819
- Fukao T, Harris T, Bailey-Serres J (2009) Evolutionary analysis of the Sub1 gene cluster that confers submergence tolerance to domesticated rice. *Ann Bot (Lond)* 103: 143–150
- Fukao T, Yeung E, Bailey-Serres J (2011) The submergence tolerance regulator SUB1A mediates crosstalk between submergence and drought tolerance in rice. *Plant Cell* 23: 412–427
- Gagne JM, Smalle J, Gingerich DJ, Walker JM, Yoo SD, Yanagisawa S, Vierstra RD (2004) *Arabidopsis* EIN3-binding F-box 1 and 2 form ubiquitin-protein ligases that repress ethylene action and promote growth by directing EIN3 degradation. *Proc Natl Acad Sci USA* 101: 6803–6808
- Gao Z, Wen C-K, Binder BM, Chen Y-F, Chang J, Chiang Y-H, Kerris RJ III, Chang C, Schaller GE (2008) Heteromeric interactions among ethylene receptors mediate signaling in *Arabidopsis*. *J Biol Chem* 283: 23801–23810
- Gao ZY, Chen YF, Randlett MD, Zhao XC, Findell JL, Kieber JJ, Schaller GE (2003) Localization of the Raf-like kinase CTR1 to the endoplasmic reticulum of *Arabidopsis* through participation in ethylene receptor signaling complexes. *J Biol Chem* 278: 34725–34732
- Goeschl JD (1975) Concentration dependencies of some effects of ethylene on etiolated pea, peanut, bean, and cotton seedlings. *Plant Physiol* 55: 670–677
- Grefen C, Städele K, Ruzicka K, Obrdlík P, Harter K, Horák J (2008) Subcellular localization and in vivo interactions of the *Arabidopsis thaliana* ethylene receptor family members. *Mol Plant* 1: 308–320
- Guo HW, Ecker JR (2003) Plant responses to ethylene gas are mediated by SCF(EBF1/EBF2)-dependent proteolysis of EIN3 transcription factor. *Cell* 115: 667–677
- Guzmán P, Ecker JR (1990) Exploiting the triple response of *Arabidopsis* to identify ethylene-related mutants. *Plant Cell* 2: 513–523
- Hattori Y, Nagai K, Furukawa S, Song X-J, Kawano R, Sakakibara H, Wu J, Matsumoto T, Yoshimura A, Kitano H, et al (2009) The ethylene response factors SNORKEL1 and SNORKEL2 allow rice to adapt to deep water. *Nature* 460: 1026–1030
- Hedden P, Graebe JE (1985) Inhibition of gibberellin biosynthesis by paclobutrazol in cell-free homogenates of *Cucurbita maxima* endosperm and *Malus pumila* embryos. *J Plant Growth Regul* 4: 111–122
- Hiraga S, Sasaki K, Hibi T, Yoshida H, Uchida E, Kosugi S, Kato T, Mie T, Ito H, Katou S, et al (2009) Involvement of two rice *ETHYLENE* INSENSITIVE3-LIKE genes in wound signaling. *Mol Genet Genomics* 282: 517–529
- Hoffmann-Benning S, Kende H (1992) On the role of abscisic acid and gibberellin in the regulation of growth in rice. *Plant Physiol* 99: 1156–1161
- Hu Y, Zhao L, Chong K, Wang T (2008) Overexpression of OsERF1, a novel rice ERF gene, up-regulates ethylene-responsive genes expression besides affects growth and development in *Arabidopsis*. *J Plant Physiol* 165: 1717–1725
- Hua J, Meyerowitz EM (1998) Ethylene responses are negatively regulated by a receptor gene family in *Arabidopsis thaliana*. *Cell* 94: 261–271
- Hua J, Sakai H, Nourizadeh S, Chen QG, Bleecker AB, Ecker JR, Meyerowitz EM (1998) EIN4 and ERS2 are members of the putative ethylene receptor gene family in *Arabidopsis*. *Plant Cell* 10: 1321–1332
- Huang YF, Li H, Hutchison CE, Laskey J, Kieber JJ (2003) Biochemical and functional analysis of CTR1, a protein kinase that negatively regulates ethylene signaling in *Arabidopsis*. *Plant J* 33: 221–233
- Jackson MB (1983) Regulation of root growth and morphology by ethylene and other externally applied growth substances. In M Jackson, A Stead, eds, *Growth Regulators in Root Development*, Monograph No. 10. British Plant Growth Regulator Group, London, pp 103–116
- Jacobsen SE, Binkowski KA, Olszewski NE (1996) SPINDLY, a tetratricopeptide repeat protein involved in gibberellin signal transduction in *Arabidopsis*. *Proc Natl Acad Sci USA* 93: 9292–9296
- Jacobsen SE, Olszewski NE (1993) Mutations at the SPINDLY locus of *Arabidopsis* alter gibberellin signal transduction. *Plant Cell* 5: 887–896
- Jain M, Nijhawan A, Tyagi AK, Khurana JP (2006) Validation of house-keeping genes as internal control for studying gene expression in rice by quantitative real-time PCR. *Biochem Biophys Res Commun* 345: 646–651
- Jung K-H, Seo Y-S, Walia H, Cao P, Fukao T, Canlas PE, Amonpant F, Bailey-Serres J, Ronald PC (2010) The submergence tolerance regulator Sub1A mediates stress-responsive expression of AP2/ERF transcription factors. *Plant Physiol* 152: 1674–1692
- Keith KA, Raskin I, Kende H (1986) A comparison of the submergence response of deepwater and non-deepwater rice. *Plant Physiol* 80: 479–482
- Kende H, van der Knaap E, Cho HT (1998) Deepwater rice: a model plant to study stem elongation. *Plant Physiol* 118: 1105–1110
- Kieber JJ, Rothenberg M, Roman G, Feldmann KA, Ecker JR (1993) CTR1, a negative regulator of the ethylene response pathway in *Arabidopsis*, encodes a member of the raf family of protein kinases. *Cell* 72: 427–441
- Kim H, Helmbrecht EE, Stalans MB, Schmitt C, Patel N, Wen C-K, Wang W, Binder BM (2011) Ethylene receptor ETHYLENE RECEPTOR1 domain requirements for ethylene responses in *Arabidopsis* seedlings. *Plant Physiol* 156: 417–429
- Klee HJ, Giovannoni JJ (2011) Genetics and control of tomato fruit ripening and quality attributes. *Annu Rev Genet* 45: 41–59
- Koorneef M, Elgersma A, Hanhart CJ, van Loenen-Martinet EP, van Rijn L, Zeevaert JAD (1985) A gibberellin insensitive mutant of *Arabidopsis thaliana*. *Physiol Plant* 65: 33–39
- Ku HS, Suge H, Rappaport L, Pratt HK (1970) Stimulation of rice coleoptile growth by ethylene. *Planta* 90: 333–339
- Lee J-H, Deng XW, Kim WT (2006) Possible role of light in the maintenance of EIN3/EIL1 stability in *Arabidopsis* seedlings. *Biochem Biophys Res Commun* 350: 484–491
- Lee K-W, Chen P-W, Lu C-A, Chen S, Ho T-HD, Yu S-M (2009) Coordinated responses to oxygen and sugar deficiency allow rice seedlings to tolerate flooding. *Sci Signal* 2: ra61
- Ma B, Chen SY, Zhang JS (2010) Ethylene signaling in rice. *Chin Sci Bull* 55: 2204–2210
- Mao C, Wang S, Jia Q, Wu P (2006) OsEIL1, a rice homolog of the *Arabidopsis* EIN3 regulates the ethylene response as a positive component. *Plant Mol Biol* 61: 141–152
- Mattoo AK, Suttle JC (1991) *The Plant Hormone Ethylene*. CRC Press Inc., Boca Raton, FL
- Mayerhofer H, Panneerselvam S, Mueller-Dieckmann J (2012) Protein kinase domain of CTR1 from *Arabidopsis thaliana* promotes ethylene receptor cross talk. *J Mol Biol* 415: 768–779
- McDaniel BK, Binder BM (2012) Ethylene Receptor 1 (ETR1) is sufficient and has the predominant role in mediating inhibition of ethylene responses by silver in *Arabidopsis thaliana*. *J Biol Chem* 287: 26094–26103
- Mitchum MG, Yamaguchi S, Hanada A, Kuwahara A, Yoshioka Y, Kato T, Tabata S, Kamiya Y, Sun TP (2006) Distinct and overlapping roles

- of two gibberellin 3-oxidases in Arabidopsis development. *Plant J* **45**: 804–818
- Murashige T, Skoog F** (1962) A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiol Plant* **15**: 473–497
- Neljubow DN** (1901) Über die horizontale Nutation der Stengel von *Pisum sativum* und einiger anderer Pflanzen. *Beitr Bot Centralbl* **10**: 128–138
- Parks BM, Spalding EP** (1999) Sequential and coordinated action of phytochromes A and B during Arabidopsis stem growth revealed by kinetic analysis. *Proc Natl Acad Sci USA* **96**: 14142–14146
- Pearce DW, Reid DM, Pharis RP** (1991) Ethylene-mediated regulation of gibberellin content and growth in *Helianthus annuus* L. *Plant Physiol* **95**: 1197–1202
- Peng J, Carol P, Richards DE, King KE, Cowling RJ, Murphy GP, Harberd NP** (1997) The Arabidopsis *GAI* gene defines a signaling pathway that negatively regulates gibberellin responses. *Genes Dev* **11**: 3194–3205
- Peng J, Harberd NP** (1993) Derivative alleles of the Arabidopsis gibberellin-insensitive (*gai*) mutation confer a wild-type phenotype. *Plant Cell* **5**: 351–360
- Peng J, Harberd NP** (1997) Gibberellin deficiency and response mutations suppress the stem elongation phenotype of phytochrome-deficient mutants of Arabidopsis. *Plant Physiol* **113**: 1051–1058
- Peng J, Richards DE, Moritz T, Caño-Delgado A, Harberd NP** (1999) Extragenic suppressors of the Arabidopsis *gai* mutation alter the dose-response relationship of diverse gibberellin responses. *Plant Physiol* **119**: 1199–1208
- Pierik R, Cuppens MLC, Voeselek LACJ, Visser EJW** (2004) Interactions between ethylene and gibberellins in phytochrome-mediated shade avoidance responses in tobacco. *Plant Physiol* **136**: 2928–2936
- Pierik R, Tholen D, Poorter H, Visser EJW, Voeselek LACJ** (2006) The Janus face of ethylene: growth inhibition and stimulation. *Trends Plant Sci* **11**: 176–183
- Potuschak T, Lechner E, Parmentier Y, Yanagisawa S, Grava S, Konz C, Genschik P** (2003) EIN3-dependent regulation of plant ethylene hormone signaling by two arabidopsis F box proteins: EBF1 and EBF2. *Cell* **115**: 679–689
- Potuschak T, Vansiri A, Binder BM, Lechner E, Vierstra RD, Genschik P** (2006) The exoribonuclease XRN4 is a component of the ethylene response pathway in Arabidopsis. *Plant Cell* **18**: 3047–3057
- Qi W, Sun F, Wang Q, Chen M, Huang Y, Feng Y-Q, Luo X, Yang J** (2011) Rice ethylene-response AP2/ERF factor OsEATB restricts internode elongation by down-regulating a gibberellin biosynthetic gene. *Plant Physiol* **157**: 216–228
- Qiao H, Chang KN, Yazaki J, Ecker JR** (2009) Interplay between ethylene, ETP1/ETP2 F-box proteins, and degradation of EIN2 triggers ethylene responses in Arabidopsis. *Genes Dev* **23**: 512–521
- Raskin I, Kende H** (1984) Role of gibberellin in the growth response of submerged deep water rice. *Plant Physiol* **76**: 947–950
- Rausser WE, Horton RF** (1975) Rapid effects of indoleacetic acid and ethylene on the growth of intact pea roots. *Plant Physiol* **55**: 443–447
- Rijnders JGHM, Yang Y-Y, Kamiya Y, Takahashi N, Barendse GWM, Blom CWPM, Voeselek LACJ** (1997) Ethylene enhances gibberellin levels and petiole sensitivity in flooding-tolerant *Rumex palustris* but not in flooding-intolerant *R. acetosa*. *Planta* **203**: 20–25
- Rose-John S, Kende H** (1985) Short-term growth response of deep-water rice to submergence and ethylene. *Plant Sci* **38**: 129–134
- Rzewuski G, Sauter M** (2008) Ethylene biosynthesis and signaling in rice. *Plant Sci* **175**: 32–42
- Sakai H, Hua J, Chen QG, Chang C, Medrano LJ, Bleecker AB, Meyerowitz EM** (1998) ETR2 is an ETR1-like gene involved in ethylene signaling in Arabidopsis. *Proc Natl Acad Sci USA* **95**: 5812–5817
- Satler SO, Kende H** (1985) Ethylene and the growth of rice seedlings. *Plant Physiol* **79**: 194–198
- Setter TL, Laureles EV** (1996) The beneficial effect of reduced elongation growth on submergence tolerance of rice. *J Exp Bot* **47**: 1551–1559
- Shulaev V, Sargent DJ, Crowhurst RN, Mockler TC, Folkerts O, Delcher AL, Jaiswal P, Mockaitis K, Liston A, Mane SP, et al** (2011) The genome of woodland strawberry (*Fragaria vesca*). *Nat Genet* **43**: 109–116
- Solano R, Stepanova A, Chao QM, Ecker JR** (1998) Nuclear events in ethylene signaling: a transcriptional cascade mediated by ETHYLENE-INSENSITIVE3 and ETHYLENE-RESPONSE-FACTOR1. *Genes Dev* **12**: 3703–3714
- Stepanova AN, Yun J, Likhacheva AV, Alonso JM** (2007) Multilevel interactions between ethylene and auxin in Arabidopsis roots. *Plant Cell* **19**: 2169–2185
- Suge H** (1972) Mesocotyl elongation of japonica rice: effect of high temperature pre-treatment and ethylene. *Plant Cell Physiol* **13**: 401–405
- Suge H** (1985) Ethylene and gibberellin: regulation of internodal elongation and nodal root development in floating rice. *Plant Cell Physiol* **26**: 607–614
- Suge H, Katsura N, Inada K** (1971) Ethylene-light relationship in the growth of the rice coleoptile. *Planta* **101**: 365–368
- Sun T-P** (2008) Gibberellin metabolism, perception and signaling pathways in Arabidopsis. *The Arabidopsis Book* **6**: e0103, doi/10.199/tab.0103
- Talon M, Koornneef M, Zeevaart JAD** (1990) Accumulation of C₁₉-gibberellins in the gibberellin-insensitive dwarf mutant *gai* of Arabidopsis thaliana (L.) Heynh. *Planta* **182**: 501–505
- van Zanten M, Basten Snoek L, van Eck-Stouten E, Proveniers MCG, Torii KU, Voeselek LACJ, Peeters AJM, Millenaar FF** (2010) Ethylene-induced hyponastic growth in Arabidopsis thaliana is controlled by ERECTA. *Plant J* **61**: 83–95
- Vandenbussche F, Petrásek J, Zádňíková P, Hoyerová K, Pešek B, Raz V, Swarup R, Bennett M, Zázimalová E, Benková E, et al** (2010) The auxin influx carriers AUX1 and LAX3 are involved in auxin-ethylene interactions during apical hook development in Arabidopsis thaliana seedlings. *Development* **137**: 597–606
- Vandenbussche F, Vancompernelle B, Rieu I, Ahmad M, Phillips A, Moritz T, Hedden P, Van Der Straeten D** (2007) Ethylene-induced Arabidopsis hypocotyl elongation is dependent on but not mediated by gibberellins. *J Exp Bot* **58**: 4269–4281
- Voeselek LACJ, Benschop JJ, Bou J, Cox MCH, Groeneveld HW, Millenaar FF, Vreeburg RAM, Peeters AJM** (2003) Interactions between plant hormones regulate submergence-induced shoot elongation in the flooding-tolerant dicot *Rumex palustris*. *Ann Bot (Lond)* **91**(Spec No): 205–211
- Wang X, Kong H, Ma H** (2009) F-box proteins regulate ethylene signaling and more. *Genes Dev* **23**: 391–396
- Warner HL, Leopold AC** (1971) Timing of growth regulator responses in peas. *Biochem Biophys Res Commun* **44**: 989–994
- Watanabe H, Hase S, Saigusa M** (2007) Effects of the combined application of ethephon and gibberellin on growth of rice (*Oryza sativa* L.) seedlings. *Plant Prod Sci* **10**: 468–472
- Wen C-K, Chang C** (2002) Arabidopsis RGL1 encodes a negative regulator of gibberellin responses. *Plant Cell* **14**: 87–100
- Wilson RN, Somerville CR** (1995) Phenotypic suppression of the gibberellin-insensitive mutant (*gai*) of Arabidopsis. *Plant Physiol* **108**: 495–502
- Xu K, Xu X, Fukao T, Canlas P, Maghirang-Rodriguez R, Heuer S, Ismail AM, Bailey-Serres J, Ronald PC, Mackill DJ** (2006) Sub1A is an ethylene-response-factor-like gene that confers submergence tolerance to rice. *Nature* **442**: 705–708
- Xu K, Xu X, Ronald PC, Mackill DJ** (2000) A high-resolution linkage map of the vicinity of the rice submergence tolerance locus *Sub1*. *Mol Gen Genet* **263**: 681–689
- Yanagisawa S, Yoo SD, Sheen J** (2003) Differential regulation of EIN3 stability by glucose and ethylene signalling in plants. *Nature* **425**: 521–525
- Yang H-J, Shen H, Chen L, Xing Y-Y, Wang Z-Y, Zhang J-L, Hong M-M** (2002) The *OsEBP-89* gene of rice encodes a putative EREBP transcription factor and is temporally expressed in developing endosperm and intercalary meristem. *Plant Mol Biol* **50**: 379–391
- Zádňíková P, Petrásek J, Marhavý P, Raz V, Vandenbussche F, Ding Z, Schwarzerová K, Morita MT, Tasaka M, Hejático J, et al** (2010) Role of PIN-mediated auxin efflux in apical hook development of Arabidopsis thaliana. *Development* **137**: 607–617
- Zhu Z, An F, Feng Y, Li P, Xue L, A M, Jiang Z, Kim JM, To TK, Li W, et al** (2011) Derepression of ethylene-stabilized transcription factors (EIN3/EIL1) mediates jasmonate and ethylene signaling synergy in Arabidopsis. *Proc Natl Acad Sci USA* **108**: 12539–12544