

Regulation of Carotenoid Biosynthesis by Shade Relies on Specific Subsets of Antagonistic Transcription Factors and Cofactors^{1[OPEN]}

Jordi Bou-Torrent², Gabriela Toledo-Ortiz², Miriam Ortiz-Alcaide², Nicolas Cifuentes-Esquivel³, Karen J. Halliday, Jaime F. Martinez-García*, and Manuel Rodriguez-Concepcion*

Centre for Research in Agricultural Genomics, CSIC-IRTA-UAB-UB, Campus UAB Bellaterra, 08193 Barcelona, Spain (J.B.-T., G.T.-O., M.O.-A., N.C.-E., J.F.M.-G., M.R.-C.); School of Biological Sciences, Edinburgh University, Edinburgh EH9 3JF, United Kingdom (G.T.-O., K.J.H.); and Institutió Catalana de Recerca i Estudis Avançats, 08010 Barcelona, Spain (J.F.M.-G.)

ORCID IDs: 0000-0003-1516-0341 (J.F.M.-G.); 0000-0002-1280-2305 (M.R.-C.).

Carotenoids are photosynthetic pigments essential for the protection against excess light. During deetiolation, their production is regulated by a dynamic repression-activation module formed by PHYTOCHROME-INTERACTING FACTOR1 (PIF1) and LONG HYPOCOTYL5 (HY5). These transcription factors directly and oppositely control the expression of the gene encoding PHYTOENE SYNTHASE (PSY), the first and main rate-determining enzyme of the carotenoid pathway. Antagonistic modules also regulate the responses of deetiolated plants to vegetation proximity and shade (i.e. to the perception of far-red light-enriched light filtered through or reflected from neighboring plants). These responses, aimed to adapt to eventual shading from plant competitors, include a reduced accumulation of carotenoids. Here, we show that PIF1 and related photolabile PIFs (but not photostable PIF7) promote the shade-triggered decrease in carotenoid accumulation. While HY5 does not appear to be required for this process, other known PIF antagonists were found to modulate the expression of the Arabidopsis (*Arabidopsis thaliana*) *PSY* gene and the biosynthesis of carotenoids early after exposure to shade. In particular, PHYTOCHROME-RAPIDLY REGULATED1, a transcriptional cofactor that prevents the binding of true transcription factors to their target promoters, was found to interact with PIF1 and hence directly induce *PSY* expression. By contrast, a change in the levels of the transcriptional cofactor LONG HYPOCOTYL IN FAR RED1, which also binds to PIF1 and other PIFs to regulate shade-related elongation responses, did not impact *PSY* expression or carotenoid accumulation. Our data suggest that the fine-regulation of carotenoid biosynthesis in response to shade relies on specific modules of antagonistic transcriptional factors and cofactors.

During their lifetime, plants are exposed to large variations in the quantity and quality of the incoming light that eventually determine their growth and development. Strong sunlight sometimes overwhelms the photosynthetic capacity of plants and, hence, leads to the production of highly reactive oxygen species that can potentially damage photosynthetic and cell structures. To cope with this danger, plant chloroplasts accumulate carotenoids, photoprotective compounds that channel energy away from chlorophylls and protect against reactive oxygen species and free radicals (Niyogi, 1999; Domonkos et al., 2013). On the other hand, photosynthesis and growth can be heavily compromised by the shading of nearby plants that compete for light. Phytochromes, a family of plant photoreceptors, are able to perceive changes in light quality associated with crowded (i.e. high-density) plant environments and to rapidly transduce them into changes in gene expression aimed to anticipate and avoid shading by overgrowing neighboring plants (i.e. promoting elongation growth), readjusting photosynthetic metabolism (i.e. decreasing the production of chlorophylls and carotenoids), or launching reproductive development (Franklin, 2008; Martinez-Garcia et al., 2010; Casal, 2013; Gommers et al., 2013).

Although regulating carotenoid biosynthesis and accumulation is central for plants to successfully adapt to changes in light quantity and quality, our understanding of how light cues regulate the accumulation of these essential metabolites is still limited (Ruiz-Sola and Rodríguez-Concepción, 2012). A first level of regulation of carotenoid accumulation in plants is the control of the transcription of biosynthetic genes. Recent reports have shown that the expression of the Arabidopsis (*Arabidopsis thaliana*) gene encoding PHYTOENE SYNTHASE (PSY), the first and main rate-determining enzyme of the carotenoid pathway (Ruiz-Sola and Rodríguez-Concepción, 2012), is under the direct control of two transcription factors involved in the transduction of light signals: PHYTOCHROME-INTERACTING FACTOR1 (PIF1) and LONG HYPOCOTYL5 (HY5; Toledo-Ortiz et al., 2010, 2014). PIF1 is a basic helix-loop-helix (bHLH) protein that, like other members of the so-called PIF quartet (formed by PIF1, PIF3, PIF4, and PIF5, collectively referred to as PIFq), accumulates in the dark and is degraded in the light (Leivar et al., 2008a; Leivar and Quail, 2011). By contrast, HY5 belongs to the basic Leu zipper (bZIP) family, accumulates in the light, and is degraded in the dark (Lau and Deng, 2010). PIFq and HY5 act antagonistically for a broad set of responses (Kami et al.,

2010; Lau and Deng, 2010; Leivar and Quail, 2011; Chen et al., 2013), including the control of *PSY* expression and carotenoid biosynthesis (Toledo-Ortiz et al., 2010, 2014). PIF1 (repressor) and HY5 (activator) were demonstrated to bind to the same G-box motif in the promoter of *PSY*, forming a dynamic repression-activation transcriptional module that provides robustness in response to light but also to temperature cues (Toledo-Ortiz et al., 2014).

The combination of positive and negative regulators also appears to be instrumental in the responses of plants to shade (i.e. to light signals generated by the presence of nearby vegetation; Franklin, 2008; Martinez-Garcia et al., 2010; Casal, 2013). When sunlight is filtered through leaves or reflected from the surface of neighboring plants, the red light wavelengths of the spectrum are preferentially absorbed and the resulting light becomes enriched in far-red light. The reduction in the red light-far-red light ratio (R:FR) caused by plant proximity or canopy shade displaces the photoequilibrium toward the inactive form of phytochromes, hence inducing the so-called shade-avoidance syndrome (SAS) in shade-intolerant species like *Arabidopsis* and most crops. The four PIFq members and the photostable PIF7 protein have been demonstrated to directly contribute to the SAS (Lorrain et al., 2008; Hornitschek et al., 2012; Leivar et al., 2012; Li et al., 2012; Sellaro et al., 2012). Exposure to low R:FR promotes PIFq protein accumulation (Lorrain et al., 2008; Leivar et al., 2012) and binding of PIF7 to its target promoters (Li et al., 2012). Based on their role in promoting elongation, all PIFs are considered as positive

regulators of the SAS (Lorrain et al., 2008; Leivar et al., 2012; Li et al., 2012). A decrease in the R:FR also triggers the production of negative regulators of hypocotyl elongation, including PHYTOCHROME-RAPIDLY REGULATED1 (PAR1), its paralog PAR2, and LONG HYPOCOTYL IN FAR RED1 (HFR1). These are helix-loop-helix proteins that lack a proper DNA-binding domain and act as transcriptional cofactors (Hornitschek et al., 2009; Galstyan et al., 2011, 2012): that is, they regulate gene expression by heterodimerizing with true bHLH transcription factors. For example, both PAR1 and HFR1 interact with PIFs and prevent their binding to their target promoters (Hornitschek et al., 2009; Hao et al., 2012; Shi et al., 2013). This dynamic balance of positive (PIFs) and negative (PAR1 and HFR1) regulators of the SAS was proposed to function as a gas-and-brake mechanism that prevents an excessive elongation in response to the initial low R:FR signal (Sessa et al., 2005; Roig-Villanova et al., 2007). Although a role for HY5 in the SAS has been proposed in plants grown under prolonged low R:FR (Sellaro et al., 2011; Ciolfi et al., 2013), the relevance of the antagonistic PIFq-HY5 module in the response to plant proximity remains to be determined.

Whereas the main output response of the SAS studied to date is elongation growth in seedlings, low R:FR also triggers a reduction in the levels of photosynthetic pigments (including carotenoids) in seedlings and adult plants (Roig-Villanova et al., 2007; Patel et al., 2013). Here, we investigated to what extent PIFs and their known antagonists in the control of light responses regulate carotenoid biosynthesis, and particularly *PSY* expression, in response to plant proximity signals.

RESULTS AND DISCUSSION

PIFq Proteins Repress *PSY* Gene Expression and Eventually Carotenoid Biosynthesis under Simulated Shade

Treatment of *Arabidopsis* plants with low R:FR results in a fast increase in the level of PIFq proteins (Lorrain et al., 2008; Leivar et al., 2012) and a decreased accumulation of chlorophyll and carotenoid pigments (Roig-Villanova et al., 2007; Patel et al., 2013; Supplemental Fig. S1). To investigate whether PIFs participated in the control of carotenogenesis in response to shade, we used *Arabidopsis* wild-type (ecotype Columbia) plants, a knockout allele (*pif7-2*) impaired in PIF7 function (Leivar et al., 2008a), and a quadruple mutant line (*pifq*) defective in all four PIFq proteins (Leivar et al., 2008b). After germination and growth for 2 d under continuous white light (W), seedlings were either left under W or exposed to far-red light-supplemented white light (W+FR; a treatment referred to as simulated shade) for another 5 d. At day 7, W and W+FR samples were collected to quantify the accumulation of total carotenoids (Fig. 1). Treatment with simulated shade led to decreased carotenoid accumulation in wild-type plants but had no effect on the *pifq*

¹ This work was supported by the Spanish Dirección General de Investigación (grant nos. BIO2011–23680 [including an FPI fellowship for M.O.-A.] and BIO2011–23489), the Consejo Superior de Investigaciones Científicas (JAE-DOC contract for G.T.-O.), the Gobierno de Chile (predoctoral fellowship to N.C.-E.), the Generalitat de Catalunya (grant nos. 2014SGR1434 and 2014SGR447), the Programa Iberoamericano de Ciencia y Tecnología para el Desarrollo (IBERCAROT), the U.K. Biotechnology and Biological Sciences Research Council (ROBuST, grant no. BB/F005237/1 to K.J.H.), and the European Union FP7 (TiMet, contract no. 245143) and Marie Curie (grant no. PCIG11–GA–2012–321649) programs.

² These authors contributed equally to the article.

³ Present address: Centro de Biotecnología Vegetal, Universidad Andrés Bello, 8370146 Santiago, Chile.

* Address correspondence to jaume.martinez@cragenomica.es and manuel.rodriguez@cragenomica.es.

The author responsible for distribution of materials integral to the findings presented in this article in accordance with the policy described in the Instructions for Authors (www.plantphysiol.org) is: Manuel Rodríguez-Concepción (manuel.rodriguez@cragenomica.es).

J.B.-T., G.T.-O., M.O.-A., J.F.M.-G., and M.R.-C. conceived the project and designed the experiments; J.B.-T., G.T.-O., and M.O.-A. performed most of the experiments; J.F.M.-G. and N.C.-E. carried out the analysis of the Y2H screen results; G.T.-O. and K.J.H. analyzed the ChIP assay data; J.B.-T., G.T.-O., M.O.-A., J.F.M.-G., and M.R.-C. analyzed the rest of the results; J.F.M.-G. and M.R.-C. wrote the article; J.B.-T., G.T.-O., M.O.-A., N.C.-E., J.F.M.-G., K.J.H., M.R.-C. discussed the data and revised the article.

[OPEN] Articles can be viewed without a subscription.

www.plantphysiol.org/cgi/doi/10.1104/pp.15.00552




Figure 1. Carotenoid and *PSY* transcript levels in wild-type and PIF-defective plants exposed to simulated shade. A, Carotenoids quantified

mutant (Fig. 1A). By contrast, the *pif7-2* mutant showed a virtually wild-type phenotype in terms of carotenoid accumulation under both W and W+FR (Fig. 1A). The higher carotenoid content in W-grown *pifq* plants is consistent with the proposed role for PIFq proteins as repressors of carotenoid biosynthesis (Toledo-Ortiz et al., 2010), whereas the lack of response to simulated shade indicates that PIFq activity is required to repress carotenoid accumulation in response to low R:FR (W+FR). These results together suggest that PIFq proteins, but not PIF7, are negative regulators of carotenoid biosynthesis both before and after the shade signal is perceived.

Our previous work indicated that one of the mechanisms by which PIFq proteins control carotenogenesis is the direct repression of the transcriptional activity of the *PSY* gene (Toledo-Ortiz et al., 2010). To investigate the relevance of PIFq-mediated changes in *PSY* expression for the regulation of carotenoid biosynthesis in response to plant proximity light signals, the levels of both *PSY* transcripts and total carotenoids were quantified in W-grown wild-type and *pifq* seedlings at different time points after treatment with W+FR at day 7. Consistent with Figure 1A, a decline in carotenoid levels was observed in wild-type plants after exposure to simulated shade, whereas no significant ($P < 0.05$) changes occurred in the *pifq* mutant even after 24 h of W+FR illumination (Fig. 1B). *PSY* transcripts also declined in wild-type plants after transferring them to simulated shade (Fig. 1C). A virtually identical profile of *PSY* gene expression was observed in the *pif7-2* mutant (Fig. 1C). By contrast, *pifq* mutant plants showed higher levels of *PSY* transcripts under W, as anticipated (Toledo-Ortiz et al., 2010), and a reproducible increase in *PSY* expression early (1 h) after W+FR exposure. At later time points, *PSY* transcript accumulation decreased to levels similar to those of wild-type plants (Fig. 1C), an unexpected finding based on the absence of changes in the levels of carotenoids observed in *pifq* seedlings (Fig. 1B). A possible explanation is that *PSY* protein stability is increased or its degradation rate is decreased in a PIFq-defective background. Another possibility is that high PIFq activity in wild-type plants grown under low R:FR eventually results in a reduced accumulation of carotenoids in a

in wild-type (WT), *pifq*, and *pif7-2* seedlings grown for 2 d under W and then either left under W (white bars) or transferred to W+FR (gray bars) for an additional 5 d. B, Carotenoids measured in wild-type and *pifq* seedlings grown for 7 d under W and then treated with W+FR for the indicated times. C, Quantitative reverse transcription-PCR (qPCR) analysis of *PSY* transcript levels in wild-type and *pifq* seedlings grown as described in B. The *UBIQUITIN-CONJUGATING ENZYME* (*UBC*) gene (At5g25760) was used for normalization. Values are shown relative to those in W-grown wild-type samples and correspond to means and SD of biological triplicates ($n = 3$). The mean values for total carotenoid levels in W-grown wild-type samples were $111.03 \mu\text{g g}^{-1}$ fresh weight (A) and $119.14 \mu\text{g g}^{-1}$ fresh weight (B). Asterisks in A and B mark statistically significant differences ($P < 0.05$) relative to W-grown plants. In C, W+FR exposure led to statistically significant differences ($P < 0.05$) in transcript levels relative to untreated (0-h) samples for all the genes and genotypes tested.

PSY-independent manner. Consistently, PIFq proteins are known to repress the development of chloroplasts and photosynthetic structures (Leivar et al., 2008b; Leivar and Quail, 2011; Zhang et al., 2013), hence decreasing carotenoid storage capacity. We conclude that PIFq (but not PIF7) proteins repress *PSY* early after treatment with simulated shade, whereas at later stages, they repress carotenoid accumulation by mechanisms that do not necessarily rely on controlling *PSY* expression.

HY5 Is Not Required for the Shade-Triggered Down-Regulation of Carotenoid Biosynthesis

Because the combination of negative and positive regulators has been found to be instrumental in the SAS responses (Franklin, 2008; Martinez-Garcia et al., 2010; Casal, 2013), we next aimed to identify positive regulators of carotenoid biosynthesis that could counterbalance the negative role of PIFq proteins. The first selected candidate was HY5, because this transcription factor has been shown to act antagonistically with PIFq proteins in a number of physiological processes, including light-regulated elongation growth and carotenoid biosynthesis (Kami et al., 2010; Lau and Deng, 2010; Leivar and Quail, 2011; Chen et al., 2013; Toledo-Ortiz

et al., 2014). In agreement with the described role of HY5 in repressing hypocotyl elongation in the light, wild-type plants grown under W showed shorter hypocotyls than the HY5-defective *SALK_056405* line (Oh and Montgomery, 2013), here referred to as *hy5-2* (Fig. 2). Also consistent with the proposed role of HY5 as an inducer of *PSY* expression and carotenoid biosynthesis, mutant *hy5-2* plants showed significantly ($P < 0.05$) reduced levels of *PSY* transcripts (Fig. 2B) and carotenoids (Fig. 2C) under both W and W+FR compared with the wild type. But since wild-type and *hy5-2* plants had very similar responses to simulated shade in terms of hypocotyl elongation, *PSY* expression, and carotenoid accumulation (Fig. 2), we conclude that HY5 is not required for the developmental (i.e. hypocotyl elongation) or the metabolic (i.e. carotenoid biosynthesis) responses to our W+FR treatment. In particular, the described repression-activation module formed by PIFq and HY5 proteins might not be relevant to regulate *PSY* expression in response to W+FR.

PAR1 Is a Positive Regulator of *PSY* Gene Expression and Carotenoid Accumulation during the SAS

Besides HY5, other negative (i.e. antagonistic) regulators of PIF activity and SAS elongation responses are





Figure 2. Contribution of HY5 to shade-triggered responses. A, Hypocotyl elongation response in wild-type (WT) and *hy5-2* seedlings grown for 2 d under W and then either kept in W (white bars) or transferred to W+FR (gray bars) for another 5 d. Hypocotyl length was measured using the ImageJ software (<http://rsb.info.nih.gov/>) on digital images. Columns represent means and SD of $n = 2$ independent experiments, each containing more than 30 seedlings of every genotype per treatment. B, qPCR analysis of *PSY* transcript levels in wild-type and *hy5-2* seedlings grown for 7 d under W and then treated with W+FR for the indicated times. C, Carotenoid levels in seedlings grown as described in A. The mean amount in W-grown wild-type samples was $105.74 \mu\text{g g}^{-1}$ fresh weight. D, *HY5* transcript levels in the samples described in B. Values in B to D are shown relative to those in W-grown wild-type samples and correspond to means and SD of biological triplicates ($n = 3$). Asterisks mark statistically significant differences ($P < 0.05$) relative to W-grown plants.



PAR1 and HFR1 (Sessa et al., 2005; Roig-Villanova et al., 2006, 2007; Hornitschek et al., 2009). But unlike *HY5* expression, which only slightly increases at late time points (8 h) following W+FR treatment (Ciolfi et al., 2013; Fig. 2D), the expression of *PAR1* and *HFR1* is induced soon (1 h) after exposure to low R:FR (Fig. 3). Under our experimental conditions, *PAR1* transcript levels peak early (1 h) after the W+FR treatment (Fig. 3A), similar to that observed for *PSY* transcripts in the *pifq* mutant (Fig. 1C), whereas *HFR1* expression increases steadily when plants are transferred to simulated shade (Fig. 3B). Interestingly, an increase in PAR1 activity in transgenic plants producing the PAR1-GFP fusion protein (Roig-Villanova et al., 2007) results in increased *PSY* transcript levels compared with untransformed plants (Fig. 3C). By contrast, no changes in *PSY* expression were observed when HFR1 function was up-regulated in lines producing a GFP-tagged version of a truncated HFR1 protein (here referred to as GFP- Δ N_HFR1) previously shown to be more stable than the full-length protein and highly active in vivo (Galstyan et al., 2011). Consistent with these results, analysis of carotenoid levels showed increased contents in plants with enhanced PAR1 levels, but not in those with higher HFR1 activity, relative to untransformed controls (Fig. 3C). Therefore, we conclude that *PSY* gene expression and carotenoid biosynthesis are positively regulated by PAR1 but not by HFR1. When exposed to simulated shade, lines overproducing PAR1-GFP showed no changes in carotenoid levels and a strongly attenuated down-regulation of *PSY* transcript levels (Fig. 4). These results confirm a positive role for PAR1 in the regulation of *PSY* gene expression and carotenoid accumulation during the SAS. Moreover, the similar carotenoid-related phenotypes observed in lines with down-regulated PIFq levels (*pifq* mutant) or up-regulated PAR1 levels (PAR1-GFP-overexpressing lines) compared with the wild-type controls (compare Figs. 1 and 4) further supports the conclusion that PAR1 and PIFq proteins antagonistically contribute to the control of carotenoid accumulation both under normal light conditions and in response to shade. This antagonistic effect, however, was not observed in the dark (Supplemental Fig. S1). Comparison of carotenoid

Figure 3. Contribution of PAR1 and HFR1 to the control of *PSY* expression and carotenoid biosynthesis. A, Levels of *PAR1* transcripts quantified by qPCR analysis of RNA samples from wild-type seedlings grown for 7 d under W and then treated for the indicated times with W+FR. B, Levels of *HFR1* transcripts in the samples described in A. Values in A and B are represented relative to those before the simulated shade treatment (0 h). Average and SD values of $n = 3$ independent samples are shown. C, Carotenoid levels (white bars) and *PSY* transcript abundance (gray bars) in plants producing the fusion proteins GFP- Δ N_HFR1 and PAR1-GFP and in wild-type controls (WT) grown under W for 7 d. Values are shown relative to those in wild-type samples (the mean carotenoid amount was $112.84 \mu\text{g g}^{-1}$ fresh weight) and correspond to means and SD of $n = 3$ independent samples. Asterisks mark statistically significant differences ($P < 0.05$) relative to wild-type plants.




Figure 4. Carotenoid accumulation and *PSY* expression in plants with increased PAR1 levels. A, Carotenoids measured in wild-type (WT) plants and transgenic lines overexpressing a GFP-tagged PAR1 protein (PAR1-GFP). Plants were grown for 7 d under W and then treated with W+FR for the indicated times. The mean carotenoid level in W-grown wild-type plants was $105.17 \mu\text{g g}^{-1}$ fresh weight. B, *PSY* transcript levels quantified by qPCR analysis of RNA samples from the plants described in A. Values are shown relative to those in W-grown wild-type samples and correspond to means and SD of $n = 3$ experiments. Asterisks mark statistically significant differences ($P < 0.05$) relative to W-grown plants.

levels in etiolated seedlings of wild-type, *pifq*, and PAR1-GFP-overexpressing lines showed wild-type levels in the transgenic seedlings, whereas the absence of PIFq proteins in the mutant resulted in increased carotenoid levels, as reported previously (Toledo-Ortiz et al., 2010). A possible explanation is that, opposite to PIFq proteins, PAR1-GFP is degraded in darkness (Zhou et al., 2014).

PAR1 Can Interact with PIF1 and Prevent Its Binding to the *PSY* Promoter to Directly Induce Gene Expression

Both PAR1 and HFR1 directly regulate gene expression by heterodimerizing with true bHLH transcription factors such as PIFs to prevent their binding to the promoters of target genes (Hornitschek et al., 2009; Carretero-Paulet et al., 2010; Galstyan et al., 2011; Hao et al., 2012; Cifuentes-Esquivel et al., 2013; Shi et al., 2013). In particular, HFR1 has been shown to interact with PIF1 and PIF3 in vitro and in the yeast two-hybrid (Y2H) assay (Fairchild et al., 2000; Bu et al., 2011), whereas coimmunoprecipitation and bimolecular fluorescence complementation (BiFC) assays demonstrated its interaction with PIF1, PIF4, and PIF5 in vivo (Hornitschek et al., 2009; Shi et al., 2013). PAR1 was recently shown to bind to PIF4 in vivo to prevent its binding to DNA for transcriptional activity (Hao et al., 2012). To test whether PAR1 could also interact with PIF1, the main repressor of *PSY* (Toledo-Ortiz et al., 2010), we carried out two complementary assays, Y2H and BiFC, using HFR1 as a positive control (Fig. 5; Supplemental Fig. S2). The Y2H assay showed a clear interaction between PAR1 and PIF1 (Fig. 5A), whereas BiFC experiments performed by microbombarding leek (*Allium ampeloprasum*) epidermal cells confirmed that PAR1 binds PIF1 in the nucleus (Fig. 5B). Together, these results show that PAR1 can physically interact with PIF1 in vivo.

To next determine whether the interaction of PAR1 and PIF1 could modulate *PSY* expression in response to shade, chromatin immunoprecipitation assays were carried out using double transgenic Arabidopsis lines constitutively producing the fusion proteins PAR1-GR and TAP-PIF1 (Fig. 6). PAR1-GR, a fusion of PAR1 to the glucocorticoid receptor (GR), was previously shown to be biologically active when targeted to the nucleus upon treatment of transgenic plants with dexamethasone (DEX), a synthetic glucocorticoid (Roig-Villanova et al., 2007; Fig. 6A). The TAP-PIF1 protein contains a tandem affinity purification (TAP) tag with nine copies of the MYC repeat that allows immunoprecipitation of PIF1-bound chromatin regions using an anti-MYC serum (Toledo-Ortiz et al., 2010, 2014). After germination and growth for 2 d under W, double transgenic seedlings were either left under W or exposed to W+FR for another 5 d in the presence or absence of DEX. At day 7, samples were collected and protein-DNA complexes formed by TAP-PIF1 were immunoprecipitated. Enriched DNA sequences were then quantified by qPCR using primers that amplified the PIF1 target sequence on the *PSY* promoter (Toledo-Ortiz et al., 2010). As shown in Figure 6B, W+FR treatment resulted in a higher ($P < 0.05$) proportion of immunoprecipitated PIF1-bound *PSY* promoter sequences. Therefore, it is likely that an enhanced accumulation of the PIF1 protein under simulated shade conditions results in enhanced binding to the *PSY* promoter. Moreover, treatment with both W+FR and DEX caused a reduction in the level of immunoprecipitated *PSY* promoter sequences compared




Figure 5. Interaction of PAR1 with PIF1 in vivo. A, Y2H assay of the indicated proteins fused to the activation domain (AD) or the binding domain (BD) of GAL4. Mating cells (positive transformants) were selected on synthetically defined medium (SD) lacking Leu and Trp (SD-LT), whereas protein-protein interactions were assessed on medium that also lacked His (SD-HLT). Proteins p53 and SV40 are known to interact

with samples treated only with W+FR (Fig. 6B). These results show that increased PAR1 levels in the nucleus interfere with PIF1 binding to the *PSY* promoter (Fig. 6A). Because PIF1 is a repressor of *PSY* expression (Toledo-Ortiz et al., 2010), PAR1 would then act as a positive regulator of *PSY* expression, consistent with the increased levels of *PSY* transcripts observed in light-grown PAR1-GFP-overexpressing lines (Figs. 3C and 4B).

To further confirm that PAR1 directly activates *PSY* expression by preventing the binding of transcriptional repressors like PIF1 and likely other PIFq proteins to the promoter, we used transgenic *Arabidopsis* lines producing the PAR1-GR protein in a genetic background in which PIFq proteins were either present (wild type) or absent (*pifq*). As shown in Figure 6C, treatment of W-grown PAR1-GR seedlings with DEX for 4 h resulted in a significant ($P < 0.05$) increase in *PSY* transcript levels compared with untreated plants, whereas no changes were detected in *pifq* PAR1-GR seedlings. This result confirms that *PSY* expression can be up-regulated when PAR1 levels increase in the nucleus and that this response requires the presence of PIFq proteins. Moreover, treatment of PAR1-GR plants with the protein synthesis inhibitor cycloheximide had no effect on the DEX-mediated increase of *PSY* transcript levels (Fig. 6C). Together, our data support the conclusion that PAR1 directly activates *PSY* gene expression by forming heterodimers with PIF1 and likely other PIFq proteins that repress *PSY* expression, hence preventing the binding of these transcription factors to their target sequences in the *PSY* promoter (Fig. 6A).

Specific Interactions of PAR1 and HFR1 with Distinct Subsets of Transcription Factors Might Control Different SAS Responses

Because both PAR1 and HFR1 can bind to PIF1 (Fig. 5; Supplemental Fig. S2) and other PIFq proteins but only PAR1 regulates *PSY* gene expression and carotenoid biosynthesis in plants (Fig. 3C), it can be speculated that the PAR1-PIF and HFR1-PIF interaction modules might somehow unleash different signaling pathways that eventually regulate common sets of responses (such as those causing hypocotyl elongation) but also specific ones (like carotenoid biosynthesis). It is also possible that

and were used as a positive control. B, BiFC assay of the indicated proteins fused to the N-terminal domain (YN) or the C-terminal domain (YC) of yellow fluorescent protein (YFP). Leek cells were cobombarded with plasmids to express the indicated fusion proteins and an additional vector producing the DsRed protein as a marker of cells expressing the microbombarded constructs. The top row shows the DsRed fluorescence, the middle row shows the fluorescence of reconstituted YFP (indicative of a positive interaction of YN and YC fusions), and the bottom row shows the bright-field images of the same area. Proteins MEDEA (MEA) and FERTILIZATION-INDEPENDENT ENDOSPERM (FIE), known to interact in both the nucleus and the cytoplasm of plant cells (Bracha-Drori et al., 2004), were used as a positive control. A negative empty plasmid control is also shown in the right column. Additional controls are shown in Supplemental Figure S2. Bars = 20 μ m.




Figure 6. PAR1 is a direct positive regulator of *PSY* expression. *A*, Proposed model for the direct regulation of *PSY* gene expression by PIF1 and PAR1. PIF1 is unstable under high R:FR (W) but accumulates under low R:FR achieved by simulated shade (W+FR) treatment, eventually binding to the *PSY* promoter to repress it. However, when PAR1 levels are high (e.g. when nuclear translocation of the chimeric PAR1-GR protein is promoted by DEX treatment), interaction with PIF1 prevents its binding to the promoter and *PSY* expression increases. *B*, Chromatin immunoprecipitation assay of the influence of PAR1 on PIF1 binding to the *PSY* promoter. Double transgenic lines producing the fusion proteins TAP-PIF1 and PAR1-GR were grown for 2 d under W and then either left under W (white column) or exposed to W+FR for another 5 d in the presence (black column) or absence (gray column) of DEX. Chromatin immunoprecipitation was carried out using an anti-MYC antibody recognizing the tandem affinity purification tag. A nonantibody control sample was processed in parallel in each case. Immunoprecipitated DNA was analyzed by qPCR using specific primers covering the PIF1-binding motif in the *PSY* promoter. Results are represented relative to W-grown samples after normalization with the nonantibody control. Columns represent means and *sd* of biological triplicates. *C*, Single transgenic plants expressing only the PAR1-GR fusion in a wild-type or a PIF-defective (*pifq*) background were grown for 7 d under W and then treated (+) with DEX, cycloheximide (CHX), and/or mock solutions (–). Samples were collected 4 h after treatment to quantify *PSY* transcript levels. Values are shown relative to those in mock-treated PAR1-GR (wild-type) samples and correspond to means and *sd* of biological triplicates ($n = 3$). Asterisks mark statistically significant differences ($P < 0.05$) relative to mock samples.

PAR1 and HFR1 might differentially interact with other bHLH transcription factors that could further contribute to provide the observed specificity in terms of SAS responses. As a first step to understand the molecular mechanism behind the differential role of PAR1 and HFR1 on the regulation of *PSY* expression and carotenoid biosynthesis, we aimed to compare the profiles of

transcription factors that could potentially interact with these transcriptional cofactors. First, we searched for bHLH family proteins among those identified in a previous Y2H screening for PAR1-interacting partners (Cifuentes-Esquivel et al., 2013). As shown in Supplemental Table S1, 28 partners of PAR1 were found in eight of the 12 subfamilies described for bHLH proteins (Heim et al., 2003; Fig. 7). The interaction of PAR1 with individual members of subfamilies VII (PIF4 and HFR1), VIII (PAR1 and PAR2), and XII (BRASSINOSTEROID ENHANCED EXPRESSION1 [BEE1], BEE2, and BEE3) has been experimentally confirmed (Carretero-Paulet et al., 2010; Hao et al., 2012; Cifuentes-Esquivel et al., 2013). Additionally, previous work showed that PAR1 can also interact with other bHLH proteins not identified in this screen, including the atypical non-DNA-binding protein PACLOBUTRAZOL RESISTANCE1 (PRE1; Hao et al., 2012) and the subfamily V member BES1-INTERACTING MYC-LIKE1 (BIM1; Cifuentes-Esquivel et al., 2013), suggesting that the variety of bHLH targets of PAR1




Figure 7. Differential roles for PAR1 and HFR1 during the SAS. *A*, Number of members of each bHLH subfamily found to interact with PAR1, HFR1, both transcription cofactors, and none of them (N.I.) in Y2H experiments. *B*, Model for antagonistic transcriptional modules regulating SAS responses. Low R:FR up-regulates the activity of transcription factors like PIFq and PIF7 and the expression of transcription cofactors like PAR1 and HFR1 (in boldface). Repression-activation modules formed by the indicated proteins thus modulate the induction of genes required for elongation growth and the repression of genes involved in carotenoid accumulation (including *PSY*).

action might be even wider than that unveiled by the Y2H screen.

A similar Y2H screening for HFR1-interacting partners identified only six bHLH proteins, all of them belonging to subfamily VII (Supplemental Table S2). They include all four PIFq proteins, previously shown to be able to heterodimerize with HFR1 (Fairchild et al., 2000; Hornitschek et al., 2009; Bu et al., 2011; Shi et al., 2013). Y2H screenings identified only one bHLH protein (PIF5) as a common target for PAR1 and HFR1 (Fig. 7A). However, other PIFq proteins (PIF1 and PIF4) have been found to also be able to interact with both transcriptional cofactors (Hornitschek et al., 2009; Hao et al., 2012; Shi et al., 2013; Fig. 5; Supplemental Fig. S2). Nevertheless, the results suggest that HFR1 might be much more target specific than PAR1. The observation that HFR1 appears to preferentially bind PIFs (i.e. members of its own subfamily) whereas PAR1 interacts with PIFs but also with a wide variety of other bHLH proteins (Fig. 7A) suggests that the impact on *PSY* expression and carotenoid accumulation caused by increasing the levels of PAR1 (but not HFR1) might be further mediated by non-PIF bHLH transcription factors that interact with PAR1 but not with HFR1. Consistent with this conclusion, the peak in *PAR1* gene expression observed early (1 h) after W+FR treatment (Fig. 3A) correlates with a similar up-regulation of *PSY* transcript levels in the absence of PIFq proteins (i.e. in the *pifq* mutant; Fig. 1C). Identifying specific non-PIFq *PSY* repressors antagonized by PAR1 upon heterodimerization, however, will require further work.

CONCLUSION

Our data show that PIFq proteins (but not PIF7) are repressors of *PSY* expression and carotenoid biosynthesis both under W and in response to shade (Fig. 1). Besides stimulating PIF activity, low R:FR perception results in a strong up-regulation of genes coding for SAS antagonists, including the transcription cofactors PAR1 and, later, HFR1 (Sessa et al., 2005; Roig-Villanova et al., 2006, 2007; Hornitschek et al., 2009). HFR1 does not appear to regulate *PSY* expression or carotenoid biosynthesis (Fig. 3). By contrast, PAR1 promotes carotenoid biosynthesis and directly induces *PSY* expression (Fig. 4), most likely by forming nonfunctional heterodimers with PIFq proteins such as PIF1 (Figs. 5 and 6) and, perhaps, with other bHLH transcription factors that might act as *PSY* repressors (Fig. 7A). The fast and transient up-regulation of *PAR1* expression that takes place after simulated shade exposure (Fig. 3A), therefore, might act as an early break to prevent an excessive reduction in *PSY* expression, allowing the plant to rapidly resume carotenoid biosynthesis if the low R:FR signal disappears (e.g. if a commitment to the shade-avoidance lifestyle is unnecessary). If the low R:FR signal persists, carotenoid accumulation decreases independently of *PSY* expression (Fig. 1). Our results with the *hy5-2* mutant suggest that the antagonistic PIFq-HY5 module

previously described to control *PSY* expression and carotenoid biosynthesis in response to environmental (light and temperature) signals (Toledo-Ortiz et al., 2014) might not be relevant during the SAS. Consistently, PAR1 and HY5 have been shown to act in separate pathways to inhibit hypocotyl elongation under different light conditions (Zhou et al., 2014).

Our data further suggest that specific modules of bHLH family members might control different sets of SAS-associated responses (Fig. 7B). Thus, PIF7 and HFR1 are key regulators of shade-induced elongation responses (Sessa et al., 2005; Li et al., 2012) but have no apparent role in regulating carotenoid biosynthesis, whereas PIFq and PAR1 proteins regulate both SAS responses. We propose that activation-suppression transcriptional modules formed by PAR1-PIFq proteins likely exert a dynamic control of early events in the regulation of both hypocotyl growth and carotenoid biosynthesis in response to low R:FR, whereas HFR1-PIFq and potentially HFR1-PIF7 modules might specifically control elongation responses (Fig. 7B). The lack of correlation between shade-induced hypocotyl elongation and flowering initiation in over 100 Arabidopsis ecotypes (Botto and Smith, 2002) previously supported the conclusion that different signaling pathways control distinct SAS responses (Martinez-Garcia et al., 2010). The results presented here, however, go a step beyond by highlighting the coexistence of different shade signaling pathways controlling several SAS responses (hypocotyl elongation and carotenoid accumulation) at the very same stage of development (photosynthetic seedlings).

In summary, we conclude that different modules formed by activators and repressors are recruited to regulate *PSY* expression and eventually carotenoid biosynthesis in response to low R:FR. Some of them regulate other responses to the same signal (e.g. the PAR1-PIFq module also controls hypocotyl elongation in response to shade). Most strikingly, individual module components do not always regulate carotenoid biosynthesis in other developmental stages (e.g. PAR1 does not appear to control carotenoid production in etiolated seedlings; Supplemental Fig. S1) or in response to different stimuli (e.g. the burst in *PSY* transcript accumulation that takes place in Arabidopsis roots exposed to a salt stress does not require the activity of PIFq proteins; Ruiz-Sola et al., 2014). Future work should help to unveil the complexity underlying the control of carotenoid biosynthesis in order to better understand and manipulate the carotenoid profiles of particular tissues in response to specific treatments.

MATERIALS AND METHODS

Plant Material and Growth Conditions

All the Arabidopsis (*Arabidopsis thaliana*) lines used in this work are in the Columbia background. Transgenic lines producing PAR1-GFP, PAR1-GR, and GFP-ΔN_HFR1 have been described previously (Roig-Villanova et al., 2007; Galstyan et al., 2011). Double transgenic lines producing PAR1-GR and TAP-PIF1

proteins were generated by crossing the corresponding single transgenic lines (Roig-Villanova et al., 2007; Toledo-Ortiz et al., 2010). Single *pif7-2* (Leivar et al., 2008a) and quadruple *pifq* (Leivar et al., 2008b) mutants were also available in the laboratory. The *pifq* PAR1-GFP lines were constructed by transforming *pifq* plants with the PAR1-GR construct (Roig-Villanova et al., 2007). A DEX-responsive homozygous *pifq* PAR1-GFP line displaying PAR1-GR transcript levels similar to those in the original PAR1-GR line (in a wild-type background) was selected for the experiments. The SALK_056405 line (here referred to as *hy5-2*), previously shown to be a knockout mutant (Oh and Montgomery, 2013), was obtained from the Nottingham Arabidopsis Stock Centre.

Seeds were surface sterilized and sown on petri dishes with solid growth medium without Suc (Roig-Villanova et al., 2007). When comparing different lines (e.g. wild type versus mutant), they were grown together on the same plate instead of growing each line on a different plate. After stratification for at least 3 d at 4°C in the dark, the plates were incubated in growth chambers at 22°C under continuous W (25 $\mu\text{mol m}^{-2} \text{s}^{-1}$ photosynthetically active radiation, R:FR of 2.1–3.5). Simulated shade (W+FR; 25 $\mu\text{mol m}^{-2} \text{s}^{-1}$ photosynthetically active radiation, R:FR of 0.05) was generated by enriching W with supplementary far-red light provided by QB1310CS-670-735 light-emitting diode hybrid lamps (Quantum Devices). Fluence rates were measured using an EPP2000 spectrometer (StellarNet) as described (Sorin et al., 2009). For experiments involving pharmacological treatments, seeds were germinated and grown under W on sterile filter paper placed on top of solid growth medium in petri dishes. At the indicated time, the filter paper with the seedlings was transferred to fresh plates containing cycloheximide and/or DEX as described (Roig-Villanova et al., 2007). For seed production, plants were grown in the greenhouse under long-day conditions.

Quantification of Transcript and Carotenoid Levels

Whole seedlings were ground in liquid nitrogen, and part of the resulting powder was used for the quantification of carotenoid levels by HPLC or spectrometric methods as described (Rodríguez-Villalón et al., 2009). HPLC separation and quantification of carotenoids and chlorophylls were performed in an initial set of experiments. Once confirmed that all individual carotenoids showed similar responses to simulated shade (Supplemental Fig. S1), total carotenoid levels were estimated spectrophotometrically (Lichtenthaler, 1987). Concentration was calculated relative to fresh weight. The rest of the powder was used for RNA extraction using the RNeasy Plant Mini Kit (Qiagen). The synthesis of complementary DNA (cDNA) from DNase-treated RNA was carried out using the Transcriptor First-Strand cDNA Synthesis Kit (Roche). The qPCR experiments were performed as described (Rodríguez-Villalón et al., 2009) using Fast Start Universal SYBR Green Master Mix (Roche) on a Light Cycler 480 apparatus (Roche). The *UBC* (At5g25760) gene was used for normalization (Czechowski et al., 2005). Primer sequences for qPCR are listed in Supplemental Table S3. Two-tailed Student's *t* tests were performed to analyze statistical differences.

Protein-DNA and Protein-Protein Interaction Analyses

Chromatin immunoprecipitation assays were carried out as described (Toledo-Ortiz et al., 2014). The Y2H screen using HFR1 as a bait was performed by Hybrigenics (www.hybrigenics-services.com) as described previously for PAR1 (Cifuentes-Esquivel et al., 2013). Briefly, the coding sequence for HFR1 was cloned into pB27 as a C-terminal fusion to LexA, and the resulting construct (pB27-LexA-HFR1) was used as a bait to screen a random-primed Arabidopsis cDNA library from 7-d-old wild-type seedlings grown at 24°C under a long-day photoperiod. Screening of 76.6 million clones was carried out using a mating approach, and positive clones were used to sequence the corresponding prey inserts by PCR. Supplemental Tables S1 and S2 summarize the results for bHLH proteins found to interact with PAR1 and HFR1, respectively, including a confidence score (predicted biological score) attributed to each interaction.

Individual Y2H assays were performed by cell mating with the Matchmaker system (Clontech). Yeast (*Saccharomyces cerevisiae*) cells of two different strains, the *MATa* Leu auxotroph YM4271a (Liu et al., 1993) and the *MAT α* Trp auxotroph PJ694a (Uetz et al., 2000), were transformed with constructs to express fusion proteins to either the GAL4 activation domain (AD) or binding domain (BD), respectively, and selected in appropriate medium. PIF1 was fused to the activation domain by subcloning the full-length *PIF1* cDNA into the pGADT7 plasmid, yielding pJB62 [AD-PIF1]. A BD-PAR1 fusion in plasmid pCL1 was available (Galstyan et al., 2011), whereas a similar construct was generated for HFR1 by subcloning the full-length *HFR1* cDNA into the pGBKT7 plasmid to

yield pJB37 [BD-HFR1]. Independent transformants were selected on SD lacking either Leu (pJB62) or Trp (pCL1 and pJB37), grown in liquid medium, and then allowed to mate by mixing equal volumes of the two types of transformed yeast cells. Mated cells were selected on SD-Leu-Trp and then transferred to SD-His-Leu-Trp to test for protein-protein interaction. Mating was repeated at least twice, with identical results.

For BiFC experiments, cDNA sequences encoding PIF1, PAR1, and HFR1 proteins were fused to the N-terminal or C-terminal regions of YFP using appropriate vectors (Bracha-Drori et al., 2004). The full-length *PIF1* cDNA was cloned into pSY736 to yield pJB72 [YN-PIF1]. Plasmid pSY735 was used to construct vectors pMS2 [YC-PAR1] and pCN6 [YC-HFR1] as described (Cifuentes-Esquivel et al., 2013). Leek (*Allium ampeloprasum*) epidermal peels were microbombarded with DNA-coated gold microcarriers using a Biolistic PDS-1000/helium system (Bio-Rad) and incubated at 22°C in the dark for 24 h prior to observation with an Olympus FV1000-ASW confocal laser scanning microscope.

Supplemental Data

The following supplemental materials are available.

Supplemental Figure S1. HPLC profiles and quantification of carotenoid and chlorophyll pigments.

Supplemental Figure S2. BiFC assay of the interaction of HFR1 and PIF1.

Supplemental Table S1. PAR1-interacting bHLH proteins identified in the Y2H screen.

Supplemental Table S2. HFR1-interacting bHLH proteins identified in the Y2H screen.

Supplemental Table S3. Primers used in this work for qPCR analyses.

ACKNOWLEDGMENTS

We thank the Nottingham Arabidopsis Stock Centre for seeds from mutant lines, Dr. Luis Oñate for Y2H yeast strains, and M. Rosa Rodríguez-Goberna and members of the Centre for Research in Agricultural Genomics Services for technical support.

Received April 13, 2015; accepted June 16, 2015; published June 16, 2015.

LITERATURE CITED

- Botto JH, Smith H** (2002) Differential genetic variation in adaptive strategies to a common environmental signal in Arabidopsis accessions: phytochrome-mediated shade avoidance. *Plant Cell Environ* 25: 53–63
- Bracha-Drori K, Shichrur K, Katz A, Oliva M, Angelovici R, Yalovsky S, Ohad N** (2004) Detection of protein-protein interactions in plants using bimolecular fluorescence complementation. *Plant J* 40: 419–427
- Bu Q, Castillon A, Chen F, Zhu L, Huq E** (2011) Dimerization and blue light regulation of PIF1 interacting bHLH proteins in Arabidopsis. *Plant Mol Biol* 77: 501–511
- Carretero-Paulet L, Galstyan A, Roig-Villanova I, Martínez-García JF, Bilbao-Castro JR, Robertson DL** (2010) Genome-wide classification and evolutionary analysis of the bHLH family of transcription factors in Arabidopsis, poplar, rice, moss, and algae. *Plant Physiol* 153: 1398–1412
- Casal JJ** (2013) Photoreceptor signaling networks in plant responses to shade. *Annu Rev Plant Biol* 64: 403–427
- Chen D, Xu G, Tang W, Jing Y, Ji Q, Fei Z, Lin R** (2013) Antagonistic basic helix-loop-helix/bZIP transcription factors form transcriptional modules that integrate light and reactive oxygen species signaling in *Arabidopsis*. *Plant Cell* 25: 1657–1673
- Cifuentes-Esquivel N, Bou-Torrent J, Galstyan A, Gallemí M, Sessa G, Salla Martret M, Roig-Villanova I, Ruberti I, Martínez-García JF** (2013) The bHLH proteins BEE and BIM positively modulate the shade avoidance syndrome in Arabidopsis seedlings. *Plant J* 75: 989–1002
- Ciolfi A, Sessa G, Sassi M, Possenti M, Salvucci S, Carabelli M, Morelli G, Ruberti I** (2013) Dynamics of the shade-avoidance response in Arabidopsis. *Plant Physiol* 163: 331–353
- Czechowski T, Stitt M, Altmann T, Udvardi MK, Scheible WR** (2005) Genome-wide identification and testing of superior reference genes for transcript normalization in Arabidopsis. *Plant Physiol* 139: 5–17

- Domonkos I, Kis M, Gombos Z, Ughy B** (2013) Carotenoids, versatile components of oxygenic photosynthesis. *Prog Lipid Res* **52**: 539–561
- Fairchild CD, Schumaker MA, Quail PH** (2000) HFR1 encodes an atypical bHLH protein that acts in phytochrome A signal transduction. *Genes Dev* **14**: 2377–2391
- Franklin KA** (2008) Shade avoidance. *New Phytol* **179**: 930–944
- Galstyan A, Bou-Torrent J, Roig-Villanova I, Martínez-García JF** (2012) A dual mechanism controls nuclear localization in the atypical basic-helix-loop-helix protein PAR1 of *Arabidopsis thaliana*. *Mol Plant* **5**: 669–677
- Galstyan A, Cifuentes-Esquivel N, Bou-Torrent J, Martínez-García JF** (2011) The shade avoidance syndrome in *Arabidopsis*: a fundamental role for atypical basic helix-loop-helix proteins as transcriptional cofactors. *Plant J* **66**: 258–267
- Gommers CM, Visser EJ, St Onge KR, Voesenek LA, Pierik R** (2013) Shade tolerance: when growing tall is not an option. *Trends Plant Sci* **18**: 65–71
- Hao Y, Oh E, Choi G, Liang Z, Wang ZY** (2012) Interactions between HLH and bHLH factors modulate light-regulated plant development. *Mol Plant* **5**: 688–697
- Heim MA, Jakoby M, Werber M, Martin C, Weisshaar B, Bailey PC** (2003) The basic helix-loop-helix transcription factor family in plants: a genome-wide study of protein structure and functional diversity. *Mol Biol Evol* **20**: 735–747
- Hornitschek P, Kohnen MV, Lorrain S, Rougemont J, Ljung K, López-Vidriero I, Franco-Zorrilla JM, Solano R, Trevisan M, Pradervand S, et al** (2012) Phytochrome interacting factors 4 and 5 control seedling growth in changing light conditions by directly controlling auxin signaling. *Plant J* **71**: 699–711
- Hornitschek P, Lorrain S, Zoete V, Michielin O, Fankhauser C** (2009) Inhibition of the shade avoidance response by formation of non-DNA binding bHLH heterodimers. *EMBO J* **28**: 3893–3902
- Kami C, Lorrain S, Hornitschek P, Fankhauser C** (2010) Light-regulated plant growth and development. *Curr Top Dev Biol* **91**: 29–66
- Lau OS, Deng XW** (2010) Plant hormone signaling lightens up: integrators of light and hormones. *Curr Opin Plant Biol* **13**: 571–577
- Leivar P, Monte E, Al-Sady B, Carle C, Storer A, Alonso JM, Ecker JR, Quail PH** (2008a) The *Arabidopsis* phytochrome-interacting factor PIF7, together with PIF3 and PIF4, regulates responses to prolonged red light by modulating phyB levels. *Plant Cell* **20**: 337–352
- Leivar P, Monte E, Oka Y, Liu T, Carle C, Castillon A, Huq E, Quail PH** (2008b) Multiple phytochrome-interacting bHLH transcription factors repress premature seedling photomorphogenesis in darkness. *Curr Biol* **18**: 1815–1823
- Leivar P, Quail PH** (2011) PIFs: pivotal components in a cellular signaling hub. *Trends Plant Sci* **16**: 19–28
- Leivar P, Tepperman JM, Cohn MM, Monte E, Al-Sady B, Erickson E, Quail PH** (2012) Dynamic antagonism between phytochromes and PIF family basic helix-loop-helix factors induces selective reciprocal responses to light and shade in a rapidly responsive transcriptional network in *Arabidopsis*. *Plant Cell* **24**: 1398–1419
- Li L, Ljung K, Breton G, Schmitz RJ, Pruneda-Paz J, Cowing-Zitron C, Cole BJ, Ivans LJ, Pedmale UV, Jung HS, et al** (2012) Linking photoreceptor excitation to changes in plant architecture. *Genes Dev* **26**: 785–790
- Lichtenthaler HK** (1987) Chlorophylls and carotenoids: pigments of photosynthetic biomembranes. *Methods Enzymol* **148**: 351–382
- Liu H, Styles CA, Fink GR** (1993) Elements of the yeast pheromone response pathway required for filamentous growth of diploids. *Science* **262**: 1741–1744
- Lorrain S, Allen T, Duek PD, Whitelam GC, Fankhauser C** (2008) Phytochrome-mediated inhibition of shade avoidance involves degradation of growth-promoting bHLH transcription factors. *Plant J* **53**: 312–323
- Martínez-García JF, Galstyan A, Salla-Martret M, Cifuentes-Esquivel N, Gallemi M, Bou-Torrent J** (2010) Regulatory components of shade avoidance syndrome. *Adv Bot Res* **53**: 65–116
- Niyogi KK** (1999) Photoprotection revisited: genetic and molecular approaches. *Annu Rev Plant Physiol Mol Biol* **50**: 333–359
- Oh S, Montgomery BL** (2013) Phytochrome-induced SIG2 expression contributes to photoregulation of phytochrome signalling and photomorphogenesis in *Arabidopsis thaliana*. *J Exp Bot* **64**: 5457–5472
- Patel D, Basu M, Hayes S, Majláth I, Hetherington FM, Tschaplinski TJ, Franklin KA** (2013) Temperature-dependent shade avoidance involves the receptor-like kinase ERECTA. *Plant J* **73**: 980–992
- Rodríguez-Villalón A, Gas E, Rodríguez-Concepción M** (2009) Phytoene synthase activity controls the biosynthesis of carotenoids and the supply of their metabolic precursors in dark-grown *Arabidopsis* seedlings. *Plant J* **60**: 424–435
- Roig-Villanova I, Bou J, Sorin C, Devlin PF, Martínez-García JF** (2006) Identification of primary target genes of phytochrome signaling: early transcriptional control during shade avoidance responses in *Arabidopsis*. *Plant Physiol* **141**: 85–96
- Roig-Villanova I, Bou-Torrent J, Galstyan A, Carretero-Paulet L, Portolés S, Rodríguez-Concepción M, Martínez-García JF** (2007) Interaction of shade avoidance and auxin responses: a role for two novel atypical bHLH proteins. *EMBO J* **26**: 4756–4767
- Ruiz-Sola MA, Rodríguez-Concepción M** (2012) Carotenoid biosynthesis in *Arabidopsis*: a colorful pathway. *The Arabidopsis Book* **10**: e0158, doi/10.1199/tab.0158
- Ruiz-Sola MA, Rodríguez-Villalón A, Rodríguez-Concepción M** (2014) Light-sensitive Phytochrome-Interacting Factors (PIFs) are not required to regulate phytoene synthase gene expression in the root. *Plant Signal Behav* **9**: e29248
- Sellaro R, Pacin M, Casal JJ** (2012) Diurnal dependence of growth responses to shade in *Arabidopsis*: role of hormone, clock, and light signaling. *Mol Plant* **5**: 619–628
- Sellaro R, Yanovsky MJ, Casal JJ** (2011) Repression of shade-avoidance reactions by sunfleck induction of HY5 expression in *Arabidopsis*. *Plant J* **68**: 919–928
- Sessa G, Carabelli M, Sassi M, Ciolfi A, Possenti M, Mittempergher F, Becker J, Morelli G, Ruberti I** (2005) A dynamic balance between gene activation and repression regulates the shade avoidance response in *Arabidopsis*. *Genes Dev* **19**: 2811–2815
- Shi H, Zhong S, Mo X, Liu N, Nezames CD, Deng XW** (2013) HFR1 sequesters PIF1 to govern the transcriptional network underlying light-initiated seed germination in *Arabidopsis*. *Plant Cell* **25**: 3770–3784
- Sorin C, Salla-Martret M, Bou-Torrent J, Roig-Villanova I, Martínez-García JF** (2009) ATHB4, a regulator of shade avoidance, modulates hormone response in *Arabidopsis* seedlings. *Plant J* **59**: 266–277
- Toledo-Ortiz G, Huq E, Rodríguez-Concepción M** (2010) Direct regulation of phytoene synthase gene expression and carotenoid biosynthesis by phytochrome-interacting factors. *Proc Natl Acad Sci USA* **107**: 11626–11631
- Toledo-Ortiz G, Johansson H, Lee KP, Bou-Torrent J, Stewart K, Steel G, Rodríguez-Concepción M, Halliday KJ** (2014) The HY5-PIF regulatory module coordinates light and temperature control of photosynthetic gene transcription. *PLoS Genet* **10**: e1004416
- Uetz P, Giot L, Cagney G, Mansfield TA, Judson RS, Knight JR, Lockshon D, Narayan V, Srinivasan M, Pochart P, et al** (2000) A comprehensive analysis of protein-protein interactions in *Saccharomyces cerevisiae*. *Nature* **403**: 623–627
- Zhang Y, Mayba O, Pfeiffer A, Shi H, Tepperman JM, Speed TP, Quail PH** (2013) A quartet of PIF bHLH factors provides a transcriptionally centered signaling hub that regulates seedling morphogenesis through differential expression-patterning of shared target genes in *Arabidopsis*. *PLoS Genet* **9**: e1003244
- Zhou P, Song M, Yang Q, Su L, Hou P, Guo L, Zheng X, Xi Y, Meng F, Xiao Y, et al** (2014) Both PHYTOCHROME RAPIDLY REGULATED1 (PAR1) and PAR2 promote seedling photomorphogenesis in multiple light signaling pathways. *Plant Physiol* **164**: 841–852