

On the Cover: Photosynthetic activity of *pp7l* mutants during seedling development is most strongly affected in young seedlings and recovers gradually over 10 days during subsequent seedling development. These seedling and the PAM images shown represent the cotyledon phenotypes and various photosynthetic parameters of 4-to-10-day-old Col-0, *pp7l-1* and *sig2* and *sig6* mutants. Credit: Duorong Xu and Tatjana Kleine.

ON THE INSIDE

Peter V. Minorsky 1

NEWS AND VIEWS

Salicylic Acid, Senescence, and Heterosis. *Lisa M. Smith* 3

Diverse Strategies Coping with Winter in Barley and its Relatives. *Yunqing Yu* 5

Epigenetic Regulation of mRNA Polyadenylation Site Selection. *Lisa M. Smith* 7

A Bridge between Kingdoms: Introduction of a Golden Gate-Based Tool Kit for Cyanobacteria. *Raimund Nagel* 10

Plastid Sulfate Transporters Open Doors to Abiotic Stress Resistance. *Charlotte M.M. Gommers* 12

Cutting the Mustard: Evolving Endoplasmic Reticulum Structures into Endoplasmic Reticulum Bodies for Plant Defense. *Kim L. Johnson* 14

Two Is Better than One: Dual SEC11 Binding Sites Regulate SYP121-Mediated Vesicle Trafficking. *Lynn G.L. Richardson* 16

BRacing for Water Stress: Brassinosteroid Signaling Promotes Drought Survival in Wheat. *Scott Hayes* 18

Balance under Stress: A Case of Mixed Signals. *Bethany Huot* 20

LETTERS

^[OPEN]Nitrate Modulates the Differentiation of Root Distal Stem Cells. *Yalu Wang, Zhizhong Gong, Jiri Friml, and Jing Zhang*

Nitrate regulation of root stem cell activity is auxin-dependent. 22

UPDATE

^[OPEN]Genetic Engineering for Disease Resistance in Plants: Recent Progress and Future Perspectives. *Oliver Xiaou Dong and Pamela C. Ronald*

A review of the recent progress in plant genetic engineering for disease resistance highlights future challenges and opportunities in the field. 26

BREAKTHROUGH TECHNOLOGIES

^[OPEN]CyanoGate: A Modular Cloning Suite for Engineering Cyanobacteria Based on the Plant MoClo Syntax. *Ravendran Vasudevan, Grant A.R. Gale, Alejandra A. Schiavon, Anton Puzorjov, John Malin, Michael D. Gillespie, Konstantinos Vavitsas, Valentin Zulkower, Baojun Wang, Christopher J. Howe, David J. Lea-Smith, and Alistair J. McCormick*

A Golden Gate-based assembly standard developed for cloning and transformation in cyanobacteria is compatible with, and builds on, the broadly established plant modular cloning syntax. 39

An Efficient System for Ds Transposon Tagging in *Brachypodium distachyon*. *Hongyu Wu, Xiaodong Xue, Caihua Qin, Yi Xu, Yuyu Guo, Xiang Li, Wei Lv, Qinxia Li, Chuangxue Mao, Luzhao Li, Suzhen Zhao, Xiaoquan Qi, and Hailong An*

Ds transposon tagging is an efficient system in Brachypodium distachyon for large-scale insertional mutagenesis. 56

Continued on next page

RESEARCH REPORT

[OPEN] Synergistic Pectin Degradation and Guard Cell Pressurization Underlie Stomatal Pore Formation. Yue Rui, Yintong Chen, Hojae Yi, Taylor Purzycki, Virendra M. Puri, and Charles T. Anderson

Degradation of pectic homogalacturonan, but not other cell wall components, promotes stomatal pore initiation, and homogalacturonan degradation and turgor increase both contribute to pore enlargement. 66

[OPEN] Regeneration of *Solanum tuberosum* Plants from Protoplasts Induces Widespread Genome Instability. Michelle Fossi, Kirk Amundson, Sundaram Kuppu, Anne Britt, and Luca Comai

Genome sequencing in potato plants regenerated from protoplasts reveals widespread changes in chromosome number and structure. 78

RESEARCH ARTICLES

BIOCHEMISTRY AND METABOLISM

[OPEN] A MYB Triad Controls Primary and Phenylpropanoid Metabolites for Pollen Coat Patterning. Maor Battat, Asa Eitan, Ilana Rogachev, Kati Hanhineva, Alisdair Fernie, Takayuki Tohge, Jules Beekwilder, and Asaph Aharoni

A triad of Arabidopsis MYB transcription factors controls the production of flavonols and additional phenylpropanoids that are embedded as pollen coat material. 87

[OPEN] Extensive Variations in Diurnal Growth Patterns and Metabolism Among *Ulva* spp. Strains. Antoine Fort, Morgane Lebrault, Margot Allaire, Alberto A. Esteves-Ferreira, Marcus McHale, Francesca Lopez, Jose M. Fariñas-Franco, Saleh Alseekh, Alisdair R. Fernie, and Ronan Sulpice

*Sea lettuce (*Ulva* spp.) strains show extensive variation in growth and metabolism, demonstrating a large potential for strain selection in boosting biomass and metabolite yields in aquaculture.* 109

[OPEN] Isoprene Acts as a Signaling Molecule in Gene Networks Important for Stress Responses and Plant Growth. Zhaojiang Zuo, Sarathi M. Weraduwage, Alexandra T. Lantz, Lydia M. Sanchez, Sean E. Weise, Jie Wang, Kevin L. Childs, and Thomas D. Sharkey

Arabidopsis, which does not normally emit isoprene, was engineered to emit isoprene, and growth and development as well as gene expression were analyzed to determine how isoprene affects plants. 124

Deficiency in the Phosphorylated Pathway of Serine Biosynthesis Perturbs Sulfur Assimilation. Armand D. Anoman, María Flores-Tornero, Ruben M. Benstein, Samira Blau, Sara Rosa-Téllez, Andrea Bräutigam, Alisdair R. Fernie, Jesús Muñoz-Bertomeu, Sören Schilasky, Andreas J. Meyer, Stanislav Kopriva, Juan Segura, Stephan Krueger, and Roc Ros

The Phosphorylated Pathway of L-Ser Biosynthesis in Arabidopsis thaliana provides Ser for thiol assimilation in heterotrophic tissues such as roots. 153

CYP79D73 Participates in Biosynthesis of Floral Scent Compound 2-Phenylethanol in *Plumeria rubra*. Savitha Dhandapani, Jingjing Jin, Vishweshwaran Sridhar, Nam-Hai Chua, and In-Cheol Jang

A cytochrome P450 enzyme contributes to the production of 2-phenylethanol and nitrogen-containing volatiles in Plumeria. 171

[OPEN] Appropriate Thiamin Pyrophosphate Levels Are Required for Acclimation to Changes in Photoperiod. Laise Rosado-Souza, Sebastian Proost, Michael Moulin, Susan Bergmann, Samuel E. Bocobza, Asaph Aharoni, Teresa B. Fitzpatrick, Marek Mutwil, Alisdair R. Fernie, and Toshihiro Obata

Plants over-accumulating vitamin B₁ lose metabolic network flexibility and therefore cannot acclimate to an altered photoperiod. 185

CELL BIOLOGY

[OPEN] Mitochondrial Pyruvate Carriers Prevent Cadmium Toxicity by Sustaining the TCA Cycle and Glutathione Synthesis. Lilong He, Ying Jing, Jianlin Shen, Xining Li, Huiping Liu, Zilong Geng, Mei Wang, Yongqing Li, Donghua Chen, Jianwei Gao, and Wei Zhang

Response to heavy metals requires glutamate and mitochondrial pyruvate carriers. 198

Continued on next page

[^{OPEN}] A Family of NAI2-Interacting Proteins in the Biogenesis of the ER Body and Related Structures. *Zhe Wang, Xifeng Li, Nana Liu, Qi Peng, Yuexia Wang, Baofang Fan, Cheng Zhu, and Zhixiang Chen*

Three related NAI2-interacting proteins from Arabidopsis are critical components in the biogenesis of ER bodies found only in the Brassicales and related structures likely also present in other plants. 212

[^{CC-BY}] Dual Sites for SEC11 on the SNARE SYP121 Implicate a Binding Exchange during Secretory Traffic. *Ben Zhang, Rucha Karnik, Jonas Alvim, Naomi Donald, and Michael R. Blatt*

The regulatory protein SEC11 binds differentially via two motifs on the Qa-SNARE SYP121 N-terminus, only one of which overlaps with the K⁺ channel-binding motif, to facilitate a binding exchange during SNARE complex assembly. 228

GENES, DEVELOPMENT AND EVOLUTION

[^{OPEN}] Senescence and Defense Pathways Contribute to Heterosis. *Rebeca Gonzalez-Bayon, Yifei Shen, Michael Groszmann, Anyu Zhu, Aihua Wang, Annapurna D. Allu, Elizabeth S. Dennis, W. James Peacock, and Ian K. Greaves*

Reduction of levels of salicylic acid in Arabidopsis C24/Ler hybrids results in a decrease in the expression of defense response genes and a delay in senescence. 240

ABC Transporter-Mediated Transport of Glutathione Conjugates Enhances Seed Yield and Quality in Chickpea. *Udita Basu, Hari D. Upadhyaya, Rishi Srivastava, Anurag Daware, Naveen Malik, Akash Sharma, Deepak Bajaj, Laxmi Narnoliya, Virevol Thakro, Alice Kujur, Shailesh Tripathi, Chellapilla Bharadwaj, V.S. Hegde, Ajay K. Pandey, Ashok K. Singh, Akhilesh K. Tyagi, and Swarup K. Parida*

An integrated genomic strategy delineated superior alleles of an ABC transporter gene, whose marker-assisted introgression enhanced the yield of chickpea by modulating glutathione-conjugate transport. 253

[^{OPEN}] Histone Deacetylase HDA19 Affects Root Cortical Cell Fate by Interacting with SCARECROW. *Wen-Qian Chen, Colleen Drapek, Dong-Xu Li, Zhi-Hong Xu, Philip N. Benfey, and Shu-Nong Bai*

Histone deacetylase HDA19 acts through SCARECROW to regulate Arabidopsis root cortical cell fate and epidermal cell patterning. 276

The AAA-ATPase MIDASIN 1 Functions in Ribosome Biogenesis and Is Essential for Embryo and Root Development. *Peng-Cheng Li, Ke Li, Juan Wang, Chuan-Zhi Zhao, Shu-Zhen Zhao, Lei Hou, Han Xia, Chang-Le Ma, and Xing-Jun Wang*

AtMDN1 plays roles in embryogenesis and root meristem maintenance, and its mutation leads to defects in nuclear export of the pre-60S ribosome and in pre-rRNA processing. 289

[^{OPEN}] Control of Cognate Sense mRNA Translation by cis-Natural Antisense RNAs. *Jules Deforges, Rodrigo S. Reis, Philippe Jacquet, Shaoline Sheppard, Veerendra P. Gadekar, Gene Hart-Smith, Andrea Tanzer, Ivo L. Hofacker, Christian Iseli, Ioannis Xenarios, and Yves Poirier*

A robust pipeline identifies and experimentally validates cis-natural antisense transcripts controlling cognate sense mRNA translation. 305

[^{OPEN}] Extrachloroplastic PP7L Functions in Chloroplast Development and Abiotic Stress Tolerance. *Duorong Xu, Giada Marino, Andreas Klingl, Beatrix Enderle, Elena Monte, Joachim Kurth, Andreas Hiltbrunner, Dario Leister, and Tatjana Kleine*

Arabidopsis protein phosphatase PP7-like (PP7L) is a predominantly nuclear-localized protein that promotes chloroplast translation and development, thus contributing to abiotic stress tolerance. 323

[^{OPEN}] HDA9-PWR-HOS15 Is a Core Histone Deacetylase Complex Regulating Transcription and Development. *Kevin S. Mayer, Xiangsong Chen, Dean Sanders, Jiani Chen, Jianjun Jiang, Phu Nguyen, Mark Scalf, Lloyd M. Smith, and Xuehua Zhong*

Arabidopsis HISTONE DEACETYLASE9 requires both PWR and HOS15 to regulate histone modifications, gene expression, and plant development. 342

- ESA1 Is Involved in Embryo Sac Abortion in Interspecific Hybrid Progeny of Rice.* Jingjing Hou, Caihong Cao, Yini Ruan, Yanyan Deng, Yaxin Liu, Kun Zhang, Lubin Tan, Zuofeng Zhu, Hongwei Cai, Fengxia Liu, Hongying Sun, Ping Gu, Chuanqing Sun, and Yongcai Fu
- ESA1 is involved in embryo sac abortion in hybrid backcross progeny derived from common wild rice and cultivated rice.* 356
- ^[OPEN]*FT Modulates Genome-Wide DNA-Binding of the bZIP Transcription Factor FD.* Silvio Collani, Manuela Neumann, Levi Yant, and Markus Schmid
- Genomic and biochemical analyses identify targets of the flowering time regulator FD at the genome-wide scale and shed light on the requirement for interaction with the florigen FLOWERING LOCUS T.* 367
- ^[OPEN]*Ubiquitin Specific Protease 15 Has an Important Role in Regulating Grain Width and Size in Rice.* Cuilan Shi, Yulong Ren, Linglong Liu, Fan Wang, Huan Zhang, Peng Tian, Tian Pan, Yongfei Wang, Ruonan Jing, Tianzhen Liu, Fuqing Wu, Qibing Lin, Cailin Lei, Xin Zhang, Shanshan Zhu, Xiuping Guo, Jiulin Wang, Zhichao Zhao, Jie Wang, Huqu Zhai, Zhijun Cheng, and Jianmin Wan
- OsUBP15, encoding a ubiquitin-specific protease15, regulates grain width and size in rice.* 381
- Antagonistic Actions of FPA and IBM2 Regulate Transcript Processing from Genes Containing Heterochromatin.* Aurélie Deremetz, Clémentine Le Roux, Yassir Idir, Cécile Brousse, Astrid Agorio, Isabelle Gy, Jane E. Parker, and Nicolas Bouché
- Intronic heterochromatic marks, associated with alternative polyadenylation sites, are decoded by RNA-binding proteins like FPA and IBM2, to tune the expression of key regulator genes such as IBM1 or RPP7.* 392
- ^[OPEN]*Evolution of Cold Acclimation and Its Role in Niche Transition in the Temperate Grass Subfamily Pooideae.* Marian Schubert, Lars Grønvoold, Simen R. Sandve, Torgeir R. Hvidsten, and Siri Fjellheim
- Comparative analysis of cold response in Pooideae species shows that cold acclimation is common in Pooideae but that this adaptation has largely evolved independently in different tribes.* 404
- Genetics of Variable Disease Expression Conferred by Inverse Gene-For-Gene Interactions in the Wheat-Parastagonospora nodorum Pathosystem.* Amanda R. Peters Haugrud, Zengcui Zhang, Jonathan K. Richards, Timothy L. Friesen, and Justin D. Faris
- Host gene-effector interactions in the wheat-P. nodorum pathosystem display additivity, epistasis, differential effector gene expression, and complex interplay of genetic regulation.* 420
- ^[OPEN]*Comparative Transcriptomic Analysis Provides Insight into the Domestication and Improvement of Pear (P. pyrifolia) Fruit.* Xiaolong Li, Lun Liu, Meiling Ming, Hongju Hu, Mingyue Zhang, Jing Fan, Bobo Song, Shaoling Zhang, and Jun Wu
- Genes associated with stone cell formation, fruit size, and sugar content underwent directional selection during pear domestication and improvement, which contributed to drastic changes of cultivated pears.* 435
- ^[OPEN]*An Essential Role for miRNA167 in Maternal Control of Embryonic and Seed Development.* Xiaozhen Yao, Jilin Chen, Jie Zhou, Hanchuanzhi Yu, Chennan Ge, Min Zhang, Xiuhua Gao, Xinhua Dai, Zhong-Nan Yang, and Yunde Zhao
- Complete deletion of the four MIR167 genes in Arabidopsis using CRISPR/Cas9 leads to late flowering and defects in flower and seed development.* 453
- MEMBRANES, TRANSPORT AND BIOENERGETICS**
- ^[OPEN]*The Potassium Transporter SIHAK10 Is Involved in Mycorrhizal Potassium Uptake.* Jianjian Liu, Junli Liu, Jinhui Liu, Miaomiao Cui, Yujuan Huang, Yuan Tian, Aiqun Chen, and Guohua Xu
- The gene SIHAK10 encodes a mycorrhiza-specific potassium transporter of the KT/KUP/HAK family that mediates K⁺ uptake at the intraradical symbiotic interface in tomato.* 465

SIGNALLING AND RESPONSE

[OPEN] Identification of Novel Inhibitors of Auxin-Induced Ca²⁺ Signaling via a Plant-Based Chemical Screen. *Kjell De Vriese, Ellie Himschoot, Kai Dünser, Long Nguyen, Andrzej Drozdzecki, Alex Costa, Moritz K. Nowack, Jürgen Kleine-Vehn, Dominique Audenaert, Tom Beeckman, and Steffen Vanneste*

Screening of an annotated library of biologically active molecules for inhibitory effects on auxin-induced Ca²⁺ entry in BY-2 cells yielded several new inhibitors for investigating Ca²⁺ signaling. 480

B-Box Containing Proteins BBX30 and BBX31, Acting Downstream of HY5, Negatively Regulate Photomorphogenesis in *Arabidopsis*. *Yueqin Heng, Fang Lin, Yan Jiang, Mingquan Ding, Tingting Yan, Hongxia Lan, Hua Zhou, Xianhai Zhao, Dongqing Xu, and Xing Wang Deng*

Two repressors of light signaling, BBX30 and BBX31, are transcriptionally and negatively regulated by the transcription factor HY5 through direct binding to the G-box cis-element present in their promoters. 497

Mesorhizobium huakuii HtpG Interaction with nsLTP AsE246 Is Required for Symbiotic Nitrogen Fixation. *Donglai Zhou, Yanan Li, Xuting Wang, Fuli Xie, Dasong Chen, Binguang Ma, and Youguo Li*

The Mesorhizobium huakuii molecular chaperone HtpG interacts with the lipid transfer protein AsE246, which plays an essential role in effective root nodule development and symbiotic nitrogen fixation. 509

The R2R3-MYB Transcription Factor MYB49 Regulates Cadmium Accumulation. *Ping Zhang, Ruling Wang, Qiong Ju, Weiqiang Li, Lam-Son Phan Tran, and Jin Xu*

ABI5 interacts with MYB49 and represses its function by preventing its binding to the downstream genes bHLH38, bHLH101, HIPP22, and HIPP44, resulting in inactivation of IRT1 and reduced Cd uptake. 529

Proteolytic Processing of SERK3/BAK1 Regulates Plant Immunity, Development, and Cell Death. *Jinggeng Zhou, Ping Wang, Lucas A.N. Claus, Daniel V. Savatin, Guangyuan Xu, Shujing Wu, Xiangzong Meng, Eugenia Russinova, Ping He, and Libo Shan*

Arabidopsis BAK1, a co-receptor of multiple receptor-like kinases, undergoes a proteolytic cleavage process that is essential for its functions in plant immunity, growth, and cell death control. 543

Independent Regulation of Symbiotic Nodulation by the SUNN Negative and CRA2 Positive Systemic Pathways. *Carole Laffont, Emeline Huault, Pierre Gautrat, Gabriella Endre, Peter Kalo, Virginie Bourion, Gérard Duc, and Florian Frugier*

SUNN and CRA2 negative and positive systemic pathways independently regulate symbiotic nodulation in Medicago truncatula. 559

[CC-BY] AVR2 Targets BSL Family Members, Which Act as Susceptibility Factors to Suppress Host Immunity. *Dionne Turnbull, Haixia Wang, Susan Breen, Marek Malec, Shaista Naqvi, Lina Yang, Lydia Welsh, Piers Hemsley, Tian Zhendong, Frederic Brunner, Eleanor M. Gilroy, and Paul R.J. Birch*

Late blight pathogen Phytophthora infestans uses effector Avr2 to target BSL phosphatases to suppress plant immunity. 571

The Chromatin Factor HNI9 and ELONGATED HYPOCOTYL5 Maintain ROS Homeostasis under High Nitrogen Provision. *Fanny Bellegarde, Amel Maghiaoui, Jossia Boucherez, Gabriel Krouk, Laurence Lejay, Liên Bach, Alain Gojon, and Antoine Martin*

Excessive nitrogen supply leads to reactive oxygen species accumulation and requires the function of major transcriptional regulators to maintain physiological balance. 582

SULTR3s Function in Chloroplast Sulfate Uptake and Affect ABA Biosynthesis and the Stress Response. *Zhen Chen, Ping-Xia Zhao, Zi-Qing Miao, Guo-Feng Qi, Zhen Wang, Yang Yuan, Nisar Ahmad, Min-Jie Cao, Ruediger Hell, Markus Wirtz, and Cheng-Bin Xiang*

The sultr3 quintuple mutant unequivocally demonstrates that sulfate transporter subfamily 3 is responsible for more than half of the chloroplast sulfate uptake and influences downstream sulfate assimilation and ABA biosynthesis as well as sulfate-induced stomatal closure. 593

Continued from preceding page

[OPEN] BES/BZR Transcription Factor TaBZR2 Positively Regulates Drought Responses by Activation of *TaGST1*.
Xiao-Yu Cui, Yuan Gao, Jun Guo, Tai-Fei Yu, Wei-Jun Zheng, Yong-Wei Liu, Jun Chen, Zhao-Shi Xu, and You-Zhi Ma

A BES/BZR-type transcription factor, TaBZR2, activates TaGST1 to scavenge reactive oxygen species and mediates crosstalk between brassinosteroids and drought signaling pathways.

605

High-CO₂/Hypoxia-Responsive Transcription Factors DkERF24 and DkWRKY1 Interact and Activate *DkPDC2* Promoter. Qing-gang Zhu, Zi-yuan Gong, Jingwen Huang, Donald Grierson, Kun-song Chen, and Xue-ren Yin

Two high-CO₂/hypoxia responsive transcription factors from persimmon fruit, DkERF24 and DkWRKY1, form a complex and synergistically transactivate the promoter of the hypoxia-responsive gene DkPDC2.

621

[OPEN] ANAC017 Coordinates Organellar Functions and Stress Responses by Reprogramming Retrograde Signaling. Xiangxiang Meng, Lu Li, Inge De Clercq, Reena Narsai, Yue Xu, Andreas Hartmann, Diego Lozano Claros, Eddie Custovic, Mathew G. Lewsey, James Whelan, and Oliver Berkowitz

Overexpression of ANAC017 in Arabidopsis leads to restricted growth, cell death, and early senescence due to impaired organellar retrograde signaling and transcriptomic reprogramming.

634

SYSTEMS AND SYNTHETIC BIOLOGY

[OPEN] Highly Resolved Systems Biology to Dissect the Etioplast-to-Chloroplast Transition in Tobacco Leaves. Tegan Armarego-Marriott, Łucja Kowalewska, Asdrubal Burgos, Axel Fischer, Wolfram Thiele, Alexander Erban, Deserah Strand, Sabine Kahlau, Alexander Hertle, Joachim Kopka, Dirk Walther, Ziv Reich, Mark Aurel Schöttler, and Ralph Bock

A comprehensive, high-resolution overview is presented of the molecular, ultrastructural, and physiological events involved in chloroplast differentiation and thylakoid biogenesis.

654

[CC-BY] Article free via Creative Commons CC-BY 4.0 license.

[OPEN] Articles can be viewed online without a subscription.