

On the Cover: The cover image illustrates the *Medicago truncatula* plant and the chromosomal distribution of its genes encoding small secreted peptides (SSPs, scale in Mbp) that are important regulatory components in plants. Among SSP activities are regulatory roles in growth, development, reproduction, stress tolerance, nodulation, and pathogen defense. These roles imply potential applications in agriculture to improve fertilizer-use efficiency and promote growth. Cover image credits: Clarissa Boschiero, Patrick X. Zhao, Amie Stearns, and Peter K. Lundquist.

ON THE INSIDE

Peter V. Minorsky 1

NEWS AND VIEWS

EPSIN1 Contributes to Plant Immunity by Modulating the Abundance of Pattern Recognition Receptors at the Plasma Membrane. *Wei Zhang* 3

Gibberellin Signaling in Plants: Entry of a new MicroRNA Player. *Raimund Nagel* 5

Hot on the Trail of DREB2A Protein Stability. *Madeleine Seale* 7

Modifying Ripening through Modular Transcription. *Sophia G. Zebell* 9

What Are the Roles for Dehydroascorbate Reductases and Glutathione in Sustaining Ascorbate Accumulation? *Elisa Dell'Aglio and Amna Mhamdi* 11

Sulfotransferase1 Is the Enzymatic Hub of Sulfated Salicinoids in Poplar. *Konstantinos Koudounas* 13

Overcoming Algal Vitamin B12 Auxotrophy by Experimental Evolution. *Elisa Dell'Aglio* 15

Phosphoregulation within the Photorespiratory Cycle: Regulate Smarter, Adapt Better? *Marc-Sven Roell* 17

AT the Onset of DNA Replication in Arabidopsis. *Anne-Sophie Fiorucci* 19

CBL2-CIPK6-TST2-Mediated Regulation of Sugar Homeostasis. *Kerri Hunter* 21

From Fuzz to Fiber: Identification of Genes Involved in Cotton Fiber Elongation. *Kim L Johnson* 23

NIT Proteins Regulate Rice Root Plasticity in Response to Nitrate and Ammonium. *Lena Maria Müller* 25

Location, Location, Location: Phototropin 2 Action at the Chloroplast Membrane. *Scott Hayes* 27

Improving Crop Water-Use Efficiency Requires Optimizing the Circadian Clock. *Meisha Holloway-Phillips* 29

MtSSPdb: A New Database for the Small Secreted Peptide Research Community. *Eva Hellmann* 31

A Novel Role for a Phospholipase D in Plant Immunity. *Marcelo Lattarulo Campos* 33

Starting Off Right: N-Terminal Acetylation Stabilizes an Immune-Activating Protein. *Charles Copeland* 35

LETTER

[OPEN]SCF^{TIR1/AFB} Auxin Signaling for Bending Termination during Shoot Gravitropism. *Huibin Han, Hana Rakusová, Inge Verstraeten, Yuzhou Zhang, and Jiří Friml*
TIR1/AFB signaling is required and sufficient for auxin-mediated PIN3 re-polarization and shoot gravitropic bending termination. 37

RESEARCH REPORT

SUMOylation Stabilizes the Transcription Factor DREB2A to Improve Plant Thermotolerance. *Feige Wang, Yiyang Liu, Yaqiao Shi, Danlu Han, Yuanyuan Wu, Weixian Ye, Huanling Yang, Guowei Li, Feng Cui, Shubo Wan, Jianbin Lai, and Chengwei Yang*
SUMOylation increases the protein stability of DREB2A, a key transcription factor in the response to heat stress in plant cells. 41

Continued on next page

RESEARCH ARTICLES

BIOCHEMISTRY AND METABOLISM

Sensory-Directed Genetic and Biochemical Characterization of Volatile Terpene Production in Kiwifruit. Yunliu Zeng, Mindy Y. Wang, Denise C. Hunter, Adam J. Matich, Peter A. McAtee, Mareike Knäbel, Cyril Hamiaux, Elizabeth A. Popowski, Sara R. Jaeger, Niels J. Nieuwenhuizen, Yar-Khing Yauk, and Ross G. Atkinson

The signature flavor in ripe kiwifruit is due to the terpene 1,8-cineole, which is produced via Actinidiachinensis terpene synthase 1b (AcTPS1b).

51

^[OPEN]Specific Lhc Proteins Are Bound to PSI or PSII Supercomplexes in the Diatom *Thalassiosira pseudonana*. Claudio Calvaruso, Anne Rokka, Eva-Mari Aro, and Claudia Büchel

The diatom Thalassiosira pseudonana shows a specific organization of the antenna complexes linked to PSI and PSII, whereby higher oligomeric antenna complexes are only weakly connected.

67

Allelic Mutations in the Ripening-Inhibitor Locus Generate Extensive Variation in Tomato Ripening. Yasuhiro Ito, Yasuyo Sekiyama, Hiroko Nakayama, Ayako Nishizawa-Yokoi, Masaki Endo, Yoko Shima, Nobutaka Nakamura, Eiichi Kotake-Nara, Susumu Kawasaki, Sakiko Hirose, and Seiichi Toki

A series of alleles in the Ripening-inhibitor (RIN) locus, encoding a transcription activator, repressor, transcriptionally neutral form, or a null allele, confer distinct ripening phenotypes.

80

^[OPEN]Transcription Factors BLH2 and BLH4 Regulate Demethylesterification of Homogalacturonan in Seed Mucilage. Yan Xu, Yiping Wang, Xiaoyu Wang, Shengqiang Pei, Yingzhen Kong, Ruibo Hu, and Gongke Zhou

Redundant transcription factors directly modulate expression of a pectin metabolism gene to regulate mucilage composition during seed coat development.

96

^[OPEN]Dehydroascorbate Reductases and Glutathione Set a Threshold for High-Light-Induced Ascorbate Accumulation. Yusuke Terai, Hiromi Ueno, Takahisa Ogawa, Yoshihiro Sawa, Atsuko Miyagi, Maki Kawai-Yamada, Takahiro Ishikawa, and Takanori Maruta

Cooperation of dehydroascorbate reductases and glutathione sets a threshold for ascorbate accumulation under high-light stress in Arabidopsis.

112

^[OPEN]Compensatory Guaiacyl Lignin Biosynthesis at the Expense of Syringyl Lignin in 4CL1-Knockout Poplar.

Chung-Jui Tsai, Peng Xu, Liang-Jiao Xue, Hao Hu, Batbayar Nyamdari, Radnaa Naran, Xiaohong Zhou, Geert Goeminne, Ruili Gao, Erica Gjersing, Joseph Dahlen, Sivakumar Pattathil, Michael G. Hahn, Mark F. Davis, John Ralph, Wout Boerjan, and Scott A. Harding

Knocking out a lignin-associated gene in poplar reveals a compensatory pathway for lignification with links to thiol redox balance and sulfur assimilation.

123

^[OPEN]The Occurrence of Sulfated Salicinoids in Poplar and Their Formation by Sulfotransferase1. Nathalie D. Lackus, Andrea Müller, Tabea D. U. Kröber, Michael Reichelt, Axel Schmidt, Yoko Nakamura, Christian Paetz, Katrin Luck, Richard L. Lindroth, C. Peter Constabel, Sybille B. Unsicker, Jonathan Gershenzon, and Tobias G. Köllner

Sulfated salicinoids in poplar are produced by sulfotransferase1 and may function in sulfur homeostasis and plant defense.

137

^[OPEN]The Four Arabidopsis Choline/Ethanolamine Kinase Isozymes Play Distinct Roles in Metabolism and Development. Ying-Chen Lin, Galileo Estopare Araguirang, Anh H. Ngo, Kui-Ting Lin, Artik Elisa Angkawijaya, and Yuki Nakamura

Four choline/ethanolamine kinases have distinct enzymatic functions in plant development and the metabolism of choline/ethanolamine in Arabidopsis.

152

^[CC-BY]Responses of a Newly Evolved Auxotroph of *Chlamydomonas* to B₁₂ Deprivation. Freddy Bunbury, Katherine E. Helliwell, Payam Mehrshahi, Matthew P. Davey, Deborah L. Salmon, Andre Holzer, Nicholas Smirnof, and Alison G. Smith

An artificially evolved strain of the model alga Chlamydomonas provides insight into how algal B₁₂ auxotrophy has arisen naturally but also reveals the challenges to becoming B₁₂ dependent.

167

Continued on next page

[OPEN] ONE-HELIX PROTEIN1 and 2 Form Heterodimers to Bind Chlorophyll in Photosystem II Biogenesis. *Daniel Hey and Bernhard Grimm*
Chlorophyll binding to the ONE-HELIX PROTEINS1/2 depends on heterodimer formation, mediated by conserved amino acids in their chlorophyll-binding motifs, and is essential for PSII biogenesis. 179

Phosphomimetic T335D Mutation of Hydroxypyruvate Reductase 1 Modifies Cofactor Specificity and Impacts Arabidopsis Growth in Air. *Yanpei Liu, Florence Guérard, Michael Hodges, and Mathieu Jossier*
Phosphomimetic NADH-DEPENDENT HYDROXYPYRUVATE REDUCTASE1 shows a modified cofactor preference towards NADPH. 194

CELL BIOLOGY

[OPEN] Arabidopsis DNA Replication Initiates in Intergenic, AT-Rich Open Chromatin. *Emily Wheeler, Ashley M. Brooks, Lorenzo Concia, Daniel L. Vera, Emily E. Wear, Chantal LeBlanc, Umamaheswari Ramu, Matthew W. Vaughn, Hank W. Bass, Robert A. Martienssen, William F. Thompson, and Linda Hanley-Bowdoin*
DNA replication initiation sites in plants associate most strongly with AT-rich and highly accessible chromatin, and not with genes or a particular epigenetic signature. 206

[OPEN] The Unfolded Protein Response Modulates a Phosphoinositide-Binding Protein through the IRE1-bZIP60 Pathway. *Chao-Yuan Yu and Kazue Kanehara*
A plant-specific phosphoinositide-binding protein plays a role in endoplasmic reticulum stress tolerance through the IRE1-bZIP60 pathway in Arabidopsis. 221

The Calcium Sensor CBL2 and Its Interacting Kinase CIPK6 Are Involved in Plant Sugar Homeostasis via Interacting with Tonoplast Sugar Transporter TST2. *Jinwu Deng, Xiyang Yang, Weinan Sun, Yuhuan Miao, Liangrong He, and Xianlong Zhang*
Calcineurin B-like protein interacting protein kinase 6 (CIPK6) is recruited to the tonoplast by Calcineurin B-like protein 2 (CBL2) and interacts with the tonoplast-localized sugar transporter TST2 to promote sugar accumulation, in particular glucose, in cotton. 236

GENES, DEVELOPMENT AND EVOLUTION

CASEIN KINASE2-Dependent Phosphorylation of PHOSPHATE2 Fine-tunes Phosphate Homeostasis in Rice. *Fei Wang, Meiju Deng, Jieyu Chen, Qiuju He, Xinye Jia, Huaxing Guo, Jiming Xu, Yidong Liu, Shuqun Zhang, Huixia Shou, and Chuanzao Mao*
The casein kinase II phosphorylates PHO2, causing it to be degraded more quickly, which affects its role in targeting phosphate transporter for degradation, thereby maintaining phosphate homeostasis in rice. 250

[OPEN] The Functions of Chloroplast Glutamyl-tRNA in Translation and Tetrapyrrole Biosynthesis. *Shreya Agrawal, Daniel Karcher, Stephanie Ruf, and Ralph Bock*
The generation and characterization of transplastomic tobacco plants provide insight into the dual role of the chloroplast glutamyl-tRNA in protein biosynthesis and tetrapyrrole biosynthesis. 263

The Ligonlintless-2 Short Fiber Mutation Is Located within a Terminal Deletion of Chromosome 18 in Cotton. *Jinesh D. Patel, Xianzhong Huang, Lifeng Lin, Sayan Das, Rahul Chandnani, Sameer Khanal, Jeevan Adhikari, Tariq Shehzad, Hui Guo, Eileen M Roy-Zokan, Junkang Rong, and Andrew H. Paterson*
Fine mapping and functional validation of the Ligon lintless-2 (Li₂) fiber mutant elucidates the genetics and development of one of the longest single-celled fibers. 277

MEMBRANES, TRANSPORT AND BIOENERGETICS

[OPEN] OsNAR2.1 Interaction with OsNIT1 and OsNIT2 Functions in Root-growth Responses to Nitrate and Ammonium. *Miaoquan Song, Xiaorong Fan, Jingguang Chen, Hongye Qu, Le Luo, and Guohua Xu*
A nitrate transport accessory protein interacts with two nitrilase proteins to regulate root-growth responses to nitrate and ammonium supply. 289

SIGNALLING AND RESPONSE

- [CC-BY] Phototropin2 Contributes to the Chloroplast Avoidance Response at the Chloroplast-Plasma Membrane Interface. Kazuhiro Ishishita, Takeshi Higa, Hidekazu Tanaka, Shin-ichiro Inoue, Aeri Chung, Tomokazu Ushijima, Tomonao Matsushita, Toshinori Kinoshita, Masato Nakai, Masamitsu Wada, Noriyuki Suetsugu, and Eiji Gotoh
Chloroplast-localized phototropin2 regulates the chloroplast avoidance response and possibly stomatal opening. 304
- [OPEN] The Circadian Clock Influences the Long-Term Water Use Efficiency of Arabidopsis. Noriane M. L. Simon, Calum A. Graham, Nicholas E. Comben, Alistair M. Hetherington, and Antony N. Dodd
Circadian rhythms play an important role in plant water use efficiency. 317
- [OPEN] RNA Interference-Based Screen Reveals Concerted Functions of MEKK2 and CRCK3 in Plant Cell Death Regulation. Yong Yang, Jun Liu, Chuanchun Yin, Luciano de Souza Vespoli, Dongdong Ge, Yanyan Huang, Baomin Feng, Guangyuan Xu, Ana Marcia E. de A. Manhães, Shijuan Dou, Cameron Criswell, Libo Shan, Xiaofeng Wang, and Ping He
A transient RNAi-based genetic screen helps elucidate the mechanisms of plant cell death control. 331
- WRKY13 Enhances Cadmium Tolerance by Promoting D-CYSTEINE DESULFHYDRASE and Hydrogen Sulfide Production. Qing Zhang, Wei Cai, Tong-Tong Ji, Ling Ye, Ying-Tang Lu, and Ting-Ting Yuan
A WRKY transcription factor activates the expression of a Cys desulfhydrase to induce the production of hydrogen sulfide in Arabidopsis, thus enhancing plant tolerance to cadmium. 345
- N-Terminal Acetylation Stabilizes SIGMA FACTOR BINDING PROTEIN1 Involved in Salicylic Acid-Primed Cell Death. Zihao Li, Vivek Dogra, Keun Pyo Lee, Rongxia Li, Mingyue Li, Mengping Li, and Chanhong Kim
N-terminal acetylation regulates SIB1 protein stability to ensure its function in salicylic acid-primed cell death. 358
- [OPEN] The BIR2/BIR3-Associated Phospholipase D γ 1 Negatively Regulates Plant Immunity. Maria A. Schlöffel, Andrea Salzer, Wei-Lin Wan, Ringo van Wijk, Raffaele Del Corvo, Maja Šemanjski, Efthymia Symeonidi, Peter Slaby, Joachim Kilian, Boris Maček, Teun Munnik, and Andrea A. Gust
A phospholipase D is a negative regulator of plant immunity and forms complexes with regulatory receptor-like kinases. 371
- Mode of Action of the Catalytic Site in the N-Terminal Ribosome-Inactivating Domain of JIP60. Michal Przydacz, Rhian Jones, Helen G. Pennington, Gerard Belmans, Maya Bruderer, Rachel Greenhill, Tia Salter, Peter A.D. Wellham, Ernesto Cota, and Pietro D. Spanu
The ribosome inactivating domain of barley JIP60 is a potent inducer of plant cell death dependent on four conserved active site amino acid residues for its deadly activity that inhibits translation. 385

SYSTEMS AND SYNTHETIC BIOLOGY

- [OPEN] MtSSPdb: The *Medicago truncatula* Small Secreted Peptide Database. Clarissa Boschiero, Xinbin Dai, Peter Knut Lundquist, Sonali Roy, Thomas Christian de Bang, Shulan Zhang, Zhaohong Zhuang, Ivone Torres-Jerez, Michael K. Udvardi, Wolf-Rüdiger Scheible, and Patrick Xuechun Zhao
*MtSSPdb hosts a compendium of small secreted peptide sequences with annotations and an RNA-seq-based gene expression atlas for *Medicago truncatula*, a plant small secreted peptide prediction tool, and phenotyping data from synthetic peptide screens in planta.* 399

[CC-BY] Article free via Creative Commons CC-BY 4.0 license.

[OPEN] Articles can be viewed online without a subscription.