

**On the Cover:** The cover image shows flowering of the Pooideae grass *Phalaris arundinaceae* from Ridabu in Norway. Flowering of temperate Pooideae is often triggered by short day photoperiods and an extended period of above freezing cold. It is not well understood whether Pooideae in highland tropical regions respond to the same signals to induce flowering, or use cues similar to Pooideae's lowland tropical ancestors. Cover image credit: Ane Charlotte Hjertaas, Norwegian University of Life Sciences, Ås, Norway.

## ON THE INSIDE

Peter V. Minorsky 808

## NEWS AND VIEWS

Tomatoes Turn Pale in the Heat: High Temperature Reduces Red and Green Pigmentation via Phytochromes. *Martin Balcerowicz* 810

An Alternative Route for Astaxanthin Biosynthesis in Green Algae. *Tianhu Sun* 812

TPLATE Complex Subunits Are Recruited Simultaneously to the Plasma Membrane. *Kerri Hunter* 814

Timing Is Everything: *MND1* Regulates Meristem Phase Change in Barley. *Lena Maria Müller* 816

Translational Regulation of PHOSPHATE1 Affects Shoot Phosphate Levels and Biomass. *Eva Hellmann* 818

A Novel Regulator of Stomatal Immunity in Tomato. *Marcelo Lattarulo Campos* 820

## TOPICAL REVIEW

[OPEN] Understanding Past, and Predicting Future, Niche Transitions based on Grass Flowering Time Variation. *Jill C. Preston and Siri Fjellheim*

*Major grass clades show strong conservation in how their flowering time is controlled, despite apparently low evolutionary constraints at the genetic level, and a dearth of data for tropical taxa.* 822

## RESEARCH ARTICLES

### BIOCHEMISTRY AND METABOLISM

[OPEN] Maize *Glossy2* and *Glossy2-like* Genes Have Overlapping and Distinct Functions in Cuticular Lipid Deposition. *Liza Esther Alexander, Yozo Okazaki, Michael A. Schelling, Aerial Davis, Xiaobin Zheng, Ludmila Rizhsky, Marna D. Yandea-Nelson, Kazuki Saito, and Basil J. Nikolau*

*Transgenesis dissection of the functional roles of maize Glossy2 and Glossy2-like genes in cuticular lipid deposition reveals both common and distinct functions.* 840

SIMYB72 Regulates the Metabolism of Chlorophylls, Carotenoids, and Flavonoids in Tomato Fruit. *Mengbo Wu, Xin Xu, Xiaowei Hu, Yudong Liu, Haohao Cao, Helen Chan, Zehao Gong, Yujin Yuan, Yingqing Luo, Bihong Feng, Zhengguo Li, and Wei Deng*

*A single transcription factor regulates the accumulation of several color compounds in tomato fruit.* 854

Phytochrome-Dependent Temperature Perception Modulates Isoprenoid Metabolism. *Ricardo Bianchetti, Belen De Luca, Luis A. de Haro, Daniele Rosado, Diego Demarco, Mariana Conte, Luisa Bermudez, Luciano Freschi, Alisdair R. Fernie, Louise V. Michaelson, Richard P. Haslam, Magdalena Rossi, and Fernando Carrari*

*Phytochrome-mediated temperature perception compromises plastidial development and function, impairing isoprenoid metabolism in tomato leaves and fruits.* 869

Astaxanthin Is Ketolated from Zeaxanthin Independent of Fatty Acid Synthesis in *Chromochloris zofingiensis*. *Yu Zhang, Ying Ye, Wei Ding, Xuemei Mao, Yantao Li, Henri Gerken, and Jin Liu*

*Astaxanthin, derived from the ketolation of zeaxanthin, is esterified mainly with oleic acid, and accumulates independently of de novo fatty acid synthesis in the green alga Chromochloris zofingiensis.* 883

Continued on next page

- [OPEN] Melatonin Represses Oil and Anthocyanin Accumulation in Seeds. Dong Li, Yuan Guo, Da Zhang, Shuangcheng He, Jingyun Gong, Haoli Ma, Xin Gao, Zhonghua Wang, Lixi Jiang, Xiaoling Dun, Shengwu Hu, and Mingxun Chen  
*Melatonin functions as a negative regulatory signal of seed oil and anthocyanin accumulation during the maturation of A. thaliana seeds.* 898
- [OPEN] Specialized Metabolism in a Nonmodel Nightshade: Trichome Acylinositol Biosynthesis. Bryan J. Leong, Steven M. Hurney, Paul D. Fiesel, Gaurav D. Moghe, A. Daniel Jones, and Robert L. Last  
*Parallels between naranjilla acylinositol and tomato acylsucrose biosynthesis suggest a common evolutionary origin.* 915
- [OPEN] Matrix-Assisted Laser Desorption/Ionization-Mass Spectrometry Imaging of Metabolites during Sorghum Germination. Lucia Montini, Christoph Crocoll, Roslyn M. Gleadow, Mohammed Saddik Motawia, Christian Janfelt, and Nanna Bjarnholt  
*Matrix-assisted laser desorption/ionization-mass spectrometry imaging reveals that biosynthesis and recycling of the plant defense compound dhurrin are key events during sorghum grain germination.* 925
- [OPEN] Suppressed Methionine  $\gamma$ -Lyase Expression Causes Hyperaccumulation of S-Methylmethionine in Soybean Seeds. Takuya Teshima, Naohiro Yamada, Yuko Yokota, Takashi Sayama, Kenji Inagaki, Takao Koeduka, Masayoshi Uefune, Masao Ishimoto, and Kenji Matsui  
*Suppression of Met catabolism in soybean seeds causes hyperaccumulation of S-methylmethionine.* 943
- Structure and Function of the Cytochrome P450 Monooxygenase Cinnamate 4-hydroxylase from *Sorghum bicolor*. Bixia Zhang, Kevin M. Lewis, Alejandra Abril, Dmitri R. Davydov, Wilfred Vermerris, Scott E. Sattler, and ChulHee Kang  
*A high-resolution crystal structure of the sorghum cinnamate 4-hydroxylase (SbC4H) has implications for the functional properties of this protein in phenylpropanoid metabolism.* 957
- CELL BIOLOGY**
- [OPEN] COPII Components Sar1b and Sar1c Play Distinct Yet Interchangeable Roles in Pollen Development. Xin Liang, Shan-Wei Li, Li-Min Gong, Sha Li, and Yan Zhang  
*The small GTPase Sar1, a component of the COPII complex that mediates vesicular trafficking from ER to Golgi, participates in sporophytic and gametophytic control of pollen development in Arabidopsis, with different isoforms playing distinct or redundant roles.* 974
- [OPEN] High Temporal Resolution Reveals Simultaneous Plasma Membrane Recruitment of TPLATE Complex Subunits. Jie Wang, Evelien Mylle, Alexander Johnson, Nienke Besbrugge, Geert De Jaeger, Jiří Friml, Roman Pleskot, and Daniel Van Damme  
*Lowering the temperature increases the temporal resolution of visualizing protein recruitment at the plasma membrane, demonstrated by dual-color live-cell imaging of endocytic complex subunits.* 986
- ECOPHYSIOLOGY AND SUSTAINABILITY**
- [OPEN] The Root Foraging Response under Low Nitrogen Depends on DWARF1-Mediated Brassinosteroid Biosynthesis. Zhongtao Jia, Ricardo F.H. Giehl, and Nicolaus von Wirén  
*Mild nitrogen deficiency upregulates key genes involved in brassinosteroid biosynthesis to stimulate root elongation.* 998
- [OPEN] Multiple Integrated Root Phenotypes Are Associated with Improved Drought Tolerance. Stephanie P. Klein, Hannah M. Schneider, Alden C. Perkins, Kathleen M. Brown, and Jonathan P. Lynch  
*Integrated root phenotypes that optimize water use efficiency and overcome mechanical impedance from dry, hard soils are beneficial for drought tolerance in maize.* 1011
- Building a Robust Chromatin Immunoprecipitation Method with Substantially Improved Efficiency. Huimin Zhao, Hongyan Li, Yaqi Jia, Xuejing Wen, Huiyan Guo, Hongyun Xu, and Yucheng Wang  
*Optimizing various steps in ChIP methods increases ChIP enrichment by an average of 14-fold in birch and is suitable for both woody and herbaceous plants.* 1026

## GENES, DEVELOPMENT AND EVOLUTION

- [OPEN] A Chromodomain-Helicase-DNA-Binding Factor Functions in Chromatin Modification and Gene Regulation. Yue Lu, Feng Tan, Yu Zhao, Shaoli Zhou, Xiangsong Chen, Yongfeng Hu, and Dao-Xiu Zhou  
*The chromodomain-helicase-DNA-binding factor CHR729 regulates nucleosome positioning and histone modification to control the expression of poised genes in rice.* 1035
- OsATM Safeguards Accurate Repair of Meiotic Double-Strand Breaks in Rice. Chao Zhang, Fanfan Zhang, Xinjie Cheng, Kangwei Liu, Jiaqi Tang, Yafei Li, Ding Tang, Zhukuan Cheng, and Hengxiu Yu  
*OsATM functions in parallel with homologous recombination to promote accurate meiotic double-strand break repair.* 1047
- Landscape of the Noncoding Transcriptome Response of Two Arabidopsis Ecotypes to Phosphate Starvation. Thomas Blein, Coline Balzergue, Thomas Roulé, Marc Gabriel, Laetitia Scalisi, Tracy François, Céline Sorin, Aurélie Christ, Christian Godon, Etienne Delannoy, Marie-Laure Martin-Magniette, Laurent Nussaume, Caroline Hartmann, Daniel Gautheret, Thierry Desnos, and Martin Crespi  
*Thousands of lncRNAs with ecotype-specific expression, including two that likely regulate primary root growth, are potentially linked to the evolution of regulatory mechanisms among ecotypes.* 1058
- Mitochondria-Associated Pyruvate Kinase Complexes Regulate Grain Filling in Rice. Li Hu, Bin Tu, Wen Yang, Hua Yuan, Jialian Li, Lianan Guo, Ling Zheng, Weilan Chen, Xiaobo Zhu, Yuping Wang, Peng Qin, Bingtian Ma, and Shigui Li  
*Mitochondrial complexes regulate grain filling by fine-tuning sucrose translocation in rice.* 1073
- [OPEN] An Acyl-CoA N-Acyltransferase Regulates Meristem Phase Change and Plant Architecture in Barley. Agatha Walla, G. Wilma van Esse, Gwendolyn K. Kirschner, Ganggang Guo, Annika Brünje, Iris Finkemeier, Rüdiger Simon, and Maria von Korff  
*MANY NODED DWARF1 encodes an acyl-CoA N-acyltransferase that acts as a major regulator of meristem phase change, promotes reproductive growth, and controls overall shoot architecture in barley.* 1088
- [OPEN] Structure-Function Analysis of Interallelic Complementation in ROOTY Transheterozygotes. Javier Brumos, Benjamin G. Bobay, Cierra A. Clark, Jose M. Alonso, and Anna N. Stepanova  
*Structure-function modeling explores the basis of interallelic complementation in multiple combinations of rooty alleles.* 1110
- [OPEN] Cell Wall Invertase Is Essential for Ovule Development through Sugar Signaling Rather Than Provision of Carbon Nutrients. Shengjin Liao, Lu Wang, Jun Li, and Yong-Ling Ruan  
*Ovule-specific silencing of cell wall invertase inhibits initiation of the ovule by disrupting sugar signaling but without inducing carbon starvation.* 1126
- [OPEN] Modulation of Shoot Phosphate Level and Growth by PHOSPHATE1 Upstream Open Reading Frame. Rodrigo S. Reis, Jules Deforges, Tatiana Sokoloff, and Yves Poirier  
*An upstream open reading frame in the 5' untranslated region of a phosphate transporter gene inhibits its translation in a phosphate-independent manner, and this regulation can be harnessed to increase shoot phosphate accumulation and improve growth under phosphate deficiency.* 1145
- Rice GERMIN-LIKE PROTEIN 2-1 Functions in Seed Dormancy under the Control of Abscisic Acid and Gibberellic Acid Signaling Pathways. Haiting Wang, Yuman Zhang, Na Xiao, Ge Zhang, Fang Wang, Xiaoying Chen, and Rongxiang Fang  
*Expression of a gene in seed scutellum is antagonistically regulated by abscisic acid and gibberellic acid signaling pathways to control seed dormancy.* 1157
- Decreased Protein Abundance of Lycopene  $\beta$ -Cyclase Contributes to Red Flesh in Domesticated Watermelon. Jie Zhang, Honghe Sun, Shaogui Guo, Yi Ren, Maoying Li, Jinfang Wang, Haiying Zhang, Guoyi Gong, and Yong Xu  
*The gene CILCYB contributes to the red flesh color of domesticated watermelon through decreased protein level to convert lycopene to  $\beta$ -carotene in the carotenogenesis pathway.* 1171

- Maize Plant Architecture Is Regulated by the Ethylene Biosynthetic Gene *ZmACS7*. *Hongchao Li, Lijing Wang, Meishan Liu, Zhaobin Dong, Qifang Li, Shulang Fei, Hongtu Xiang, Baoshen Liu, and Weiwei Jin*  
*Alteration of the C terminus of 1-aminocyclopropane-1-carboxylic acid (ACC) Synthase 7 (ZmACS7), an ethylene biosynthetic protein, enhances the stability of this enzyme, thus increasing ACC and ethylene production, which in turn alters plant architecture in maize.* 1184
- MEMBRANES, TRANSPORT AND BIOENERGETICS**
- [OPEN] Seasonal Zinc Storage and a Strategy for Its Use in Buds of Fruit Trees. *Ruohan Xie, Jianqi Zhao, Lingli Lu, Patrick Brown, Xianyong Lin, Samuel M. Webb, Jun Ge, Olga Antipova, Luxi Li, and Shengke Tian*  
*Zn is deposited in stem nodes subtending axillary buds and then sequestered, probably as Zn-phytate, prior to dormancy break in spring, when Zn is released for use in budbreak and growth.* 1200
- [OPEN] Rethinking the Influence of Chloroplast Movements on Non-photochemical Quenching and Photoprotection. *Sam Wilson and Alexander V. Ruban*  
*Chloroplast movement-related optical alterations in plant cell ultrastructure do not impact non-photochemical quenching nor act as a photoprotective mechanism.* 1213
- [OPEN] The ZIP Transporter Family Member OsZIP9 Contributes To Root Zinc Uptake in Rice under Zinc-Limited Conditions. *Sheng Huang, Akimasa Sasaki, Naoki Yamaji, Haruka Okada, Namiki Mitani-Ueno, and Jian Feng Ma*  
*The zinc transporter OsZIP9, expressed at the exodermis and endodermis of root mature region, contributes to Zn uptake from soil in rice.* 1224
- ZINC TRANSPORTER5 and ZINC TRANSPORTER9 Function Synergistically in Zinc/Cadmium Uptake. *Longtao Tan, Mengmeng Qu, Yuxing Zhu, Can Peng, Jiurong Wang, Dongying Gao, and Caiyan Chen*  
*Zinc influx transport proteins function synergistically in rice to maintain zinc homeostasis.* 1235
- SIGNALLING AND RESPONSE**
- [OPEN] Epigenetic Activation of *Enoyl-CoA Reductase* By An Acetyltransferase Complex Triggers Wheat Wax Biosynthesis. *Lingyao Kong, Pengfei Zhi, Jiao Liu, Haoyu Li, Xiaona Zhang, Jie Xu, Jiaqi Zhou, Xiaoyu Wang, and Cheng Chang*  
*A transcriptional activator recruits a histone acetyltransferase complex that epigenetically regulates the biosynthesis of wheat cuticular wax and is essential for triggering the germination of the powdery mildew pathogen.* 1250
- [OPEN] Retrograde Induction of *phyB* Orchestrates Ethylene-Auxin Hierarchy to Regulate Growth. *Jishan Jiang, Yanmei Xiao, Hao Chen, Wei Hu, Liping Zeng, Haiyan Ke, Franck A. Ditengou, Upendra Devisetty, Klaus Palme, Julin Maloof, and Katayoon Dehesh*  
*The plastidial retrograde metabolite MEcPP orchestrates coordination of the light and hormonal signaling cascade by inducing phytochrome B abundance and modulating auxin and ethylene levels.* 1268
- [OPEN] Light and Abscisic Acid Coordinately Regulate Greening of Seedlings. *Di Xu, Di Wu, Xiao-han Li, Yu'e Jiang, Tian Tian, Qingshuai Chen, Lin Ma, Haiyang Wang, Xing Wang Deng, and Gang Li*  
*Light and abscisic acid pathways antagonistically regulate greening of seedlings by regulating the transcriptional activity of FHY3.* 1281
- [OPEN] The F-Box-Like Protein FBL17 Is a Regulator of DNA-Damage Response and Colocalizes with RETINOBLASTOMA RELATED1 at DNA Lesion Sites. *Naomie Gentric, Kinda Masoud, Robin P. Journot, Valérie Cognat, Marie-Edith Chabouté, Sandra Noir, and Pascal Genschik*  
*The mutation of FBL17 leads to a strong upregulation of genes involved in DNA damage and repair processes, and FBL17 protein is recruited at nuclear DNA lesion foci upon double-strand break induction.* 1295
- [OPEN] Tomato Atypical Receptor Kinase1 Is Involved in the Regulation of Preinvasion Defense. *Andrew R. Guzman, Jung-Gun Kim, Kyle W. Taylor, Daniel Lanver, and Mary Beth Mudgett*  
*A leucine rich repeat receptor-like pseudokinase is involved in the regulation of stomatal movement in response to bacteria and biotic elicitors, affecting resistance to bacterial invasion.* 1306

- A Cytokinin Signaling Type-B Response Regulator Transcription Factor Acting in Early Nodulation. *Sovanna Tan, Myriam Sanchez, Carole Laffont, Stéphane Boivin, Christine Le Signor, Richard Thompson, Florian Frugier, and Mathias Brault*  
*A Medicago truncatula cytokinin-signaling transcription factor is required for early symbiotic nodulation and to regulate key Nod factor signaling and endoreduplication genes.* 1319
- [OPEN] Calcium-Dependent Hydrogen Peroxide Mediates Hydrogen-Rich Water-Reduced Cadmium Uptake in Plant Roots. *Qi Wu, Liping Huang, Nana Su, Lana Shabala, Haiyang Wang, Xin Huang, Ruiyu Wen, Min Yu, Jin Cui, and Sergey Shabala*  
*Root cadmium uptake is influenced by hydrogen via modulation of a plasma membrane-based respiratory burst oxidase homolog that operates upstream of a cadmium transporter protein.* 1331
- Dual Activities of Receptor-Like Kinase OsWAKL21.2 Induce Immune Responses. *Kamal Kumar Malukani, Ashish Ranjan, Shiva Jyothi Hota, Hitendra Kumar Patel, and Ramesh V. Sonti*  
*A rice receptor-like kinase induces immune responses in rice and Arabidopsis by kinase and guanylate cyclase activities, respectively.* 1345
- [OPEN] Arabidopsis Lectin Receptor Kinase P2K2 Is a Second Plant Receptor for Extracellular ATP and Contributes to Innate Immunity. *An Quoc Pham, Sung-Hwan Cho, Cuong The Nguyen, and Gary Stacey*  
*A receptor kinase that can bind to ATP with high affinity plays a partially redundant role during plant immunity.* 1364
- [OPEN] Diverse Allyl Glucosinolate Catabolites Independently Influence Root Growth and Development. *Ella Katz, Rammyani Bagchi, Verena Jeschke, Alycia R. M. Rasmussen, Aleshia Hopper, Meike Burow, Mark Estelle, and Daniel J. Kliebenstein*  
*Allyl-glucosinolate and its catabolites use multiple mechanisms to affect plant growth and development through specific responses that are optimal to any given environment.* 1376
- [CC-BY] Self-Incompatibility Triggers Irreversible Oxidative Modification of Proteins in Incompatible Pollen. *Tamanna Haque, Deborah J. Eaves, Zongcheng Lin, Cleidiane G. Zampronio, Helen J. Cooper, Maurice Bosch, Nicholas Smirnov, and Veronica E. Franklin-Tong*  
*In the self-incompatibility response, incompatible pollen stimulates irreversible oxidative protein modifications that affect crucial protein functions.* 1391
- [CC-BY] Article free via Creative Commons CC-BY 4.0 license.  
[OPEN] Articles can be viewed online without a subscription.