

## Magnesium Uptake by Soybeans

J. E. Leggett and W. A. Gilbert<sup>1</sup>

Mineral Nutrition Laboratory, USDA, ARS, SWCRD, Beltsville, Maryland 20705

Received February 4, 1969.

**Abstract.** Magnesium contents of soybean (*Glycine max*) roots increase and the K and Ca contents decrease with increased  $MgCl_2$  concentrations in ambient solutions. The Mg uptake is inhibited when both Ca and K are present in the solution, but not by K or Ca alone. Chloride uptake, which is very low from the  $MgCl_2$  solution, is greatly enhanced by the presence of K. The selectivity against Mg imparted by  $K^+ + Ca$  appears to be at an external barrier for cation uptake as shown by its dependence on the presence of Ca in the external solution. The Ca content of roots is influenced only slightly by changes in external Ca concentrations from  $10^{-4}$  to  $10^{-2}$  M, but that of shoots is greatly enhanced as the Ca concentration is increased or the K concentration is decreased. These effects on Ca contents are explained as arising from transport to the shoot without involvement of vacuoles of root cells.

Salt uptake by plants involves interactions between the major nutrient cations,  $K^+$ ,  $Ca^{2+}$ , and  $Mg^{2+}$ , as well as  $Na^+$  and  $H^+$  under experimental and natural conditions. Predominant attention has been focused on the singly charged cations,  $K^+$  and  $Na^+$ , and to a lesser extent, effects of  $Ca^{2+}$  presence on them. Collander (2) included doubly charged cations in his study of cation uptake by many species. Moore *et al.* (17) followed the uptake of  $Mg^{2+}$  by excised barley roots as influenced by  $Ca^{2+}$ . Attention to interactions of several cations, however, is very limited and the interactions are not understood (6, 10).

We have shown previously that  $Mg^{2+}$  uptake from single-salt solutions by soybean roots is accompanied by loss of  $K^+$ ,  $Ca^{2+}$ , and  $Cl^-$  (13). Observations are reported here on the 3-fold interactions among  $Mg^{2+}$ ,  $Ca^{2+}$ , and  $K^+$ .

### Materials and Methods

**Excised Roots.** Roots were obtained from soybean seedlings (*Glycine max* L. Merr. var. Hawkeye) (13). Briefly, seeds were soaked in water 6 hr, then planted on a stainless steel screen suspended over a continuously aerated 0.2 mM  $CaSO_4$  solution. The plants were grown in a dark chamber maintained at a temperature of 27°. At the end of 4 days, the roots were excised, cut to 6-cm length from the apex and suspended in a 0.2 mM  $CaSO_4$  solution. The elapsed time between root excision and experiment initiation was less than 2 hr.

Excised root samples weighing 0.5 g were used in each liter of test solution. The initial solution pH of 5.6 was not adjusted during the 24-hr uptake

experiment. It was more convenient to adjust the root to solution ratios based upon previous experience such that the final pH values would be in the range of 5.1 to 5.5.

At the end of the experimental period, the roots were rinsed, transferred to beakers, and heated in a muffle oven to 480° for 2 hr. The ash was dissolved in 20 ml of 0.1  $HNO_3$  and 10% acetic acid solution and reserved for analysis.

Chloride analysis of plant roots after the above dry-ashing procedure was compared with the analysis after several hot water extractions. The  $Cl^-$  value was the same for both procedures. However,  $Cl^-$  will be lost if the ash is maintained at 480° for more than 8 hr. Selection of the 2-hr period was made on the basis of complete ashing and no  $Cl^-$  loss. We did not attempt to relate  $Cl^-$  loss more precisely to the ashing period.

**Intact Plants.** Lots of 5 dark-germinated seeds were placed on gauze in a holder with their roots extending into a 0.2 mM  $CaSO_4$  solution. These lots were grown on a 15-hr day at 27° for 7 days when the first trifoliolate leaf was expanding. At this time, the 5 plants had produced about 2 g of roots and 3 to 4 g wet weight of shoots. The lots were transferred after rinsing to experimental solutions as desired. Roots and shoots were separated at the end of the experimental period and prepared for analysis as described for excised roots.

**Inorganic Analysis.** Cation contents of the plants and solutions were determined by atomic absorption spectrophotometry (4). Chloride contents of the samples were assayed with an Aminco-Cotlove automatic chloride titrator (3).

### Results and Discussion

**Uptake by Excised Roots.** Magnesium uptake by soybean roots from  $MgCl_2$  solutions was observed

<sup>1</sup> Present address: Department of Biology, Miami Dade Junior College, North Campus, Miami, Florida.




FIG. 1. Influence of the  $MgCl_2$  concentration of absorption solutions on the Mg, K, and Ca contents of excised soybean roots. Absorption time, 24 hr.

over a concentration range of 0.1 to 10 mM (Fig. 1). These roots decreased in  $K^+$  and  $Ca^{2+}$  content with increase in  $Mg^{2+}$  concentration. The attained chloride level was below  $1 \mu eq g^{-1}$  fresh weight of roots for the 24-hr uptake at all  $MgCl_2$  concentrations. Although the  $Mg^{2+}$  uptake approximates the loss of  $K^+$  plus  $Ca^{2+}$ , we hesitate to infer that  $Mg^{2+}$  uptake is dependent only on an equivalent  $K^+$  plus  $Ca^{2+}$  exchange because the charge balance of  $K^+$  plus  $Ca^{2+}$ , and  $Cl^-$  loss is less than the  $Mg^{2+}$  uptake for roots previously loaded with KCl (cf. Fig. 4 ref 13). Nevertheless, in the absence of  $Cl^-$  entry,  $Mg^{2+}$  uptake must occur in exchange for other cations.

If the loss of  $K^+$  and  $Ca^{2+}$  represented a general cellular "leakiness," a decreased respiration rate might be expected to result from simultaneous loss of substrates. Oxygen consumption and  $CO_2$  production, however, remained constant, independent of the  $Mg^{2+}$  treatment (13). The results rather indicate increased membrane permeability only with respect to the inorganic cations.

A further test for the integrity of the cellular structures is afforded by results with 2,4-dinitrophenol (DNP) (table I) which was used as a

Table I. Influence of DNP on Ion Retention by the Soybean Root

Absorption time was 24 hr.

Treatment	K	Tissue content	
		Mg	Ca
	$\mu eq/g$ fresh wt		
$CaSO_4$ 0.2 mM	55	3	3
$MgCl_2$ 10 mM	5	46	<1
DNP 0.01 mM	0	2	2
$MgCl_2$ 10 mM			
DNP + 0.01 mM	0	18	<1
$H_2O$	50	3	2

metabolic inhibitor without regard to its exact mode of action. The observed reduction by DNP of the Mg and K contents in the tissue indicated that the ion levels depended on metabolism. Barker and Koontz (1) in studying DNP and  $Ca^{2+}$  uptake concluded that DNP altered the barrier to both DNP and  $Ca^{2+}$  transport. In the present report, DNP induced loss of  $Mg^{2+}$ ,  $Ca^{2+}$ , and  $K^+$ , whereas the high  $Mg^{2+}$  accumulation was correlated only with the loss of  $Ca^{2+}$  and  $K^+$ . Magnesium accumulation must result from an energy requiring step because of the observed DNP inhibition.

Some of the influences of cation interactions are shown by results in Fig. 2. The solution used initially contained 10 mM  $MgCl_2$  at various pH's. The roots lost both  $K^+$  and  $Ca^{2+}$  at each acidity. Potassium losses were greatest under the less acid conditions. Magnesium uptake increased with increasing pH, with the greatest change occurring up to pH 5.




FIG. 2. Influence of H ion concentration on the changes in Mg, K, and Ca contents of excised soybean roots in a 10 mM  $MgCl_2$  solution for 24 hr. The roots were grown in 0.2 mM  $CaSO_4$ .

Attempts to determine the amount of total root  $Ca^{2+}$  necessary to induce a change in ion preference have not been successful. The evidence points to the necessity for  $Ca^{2+}$  in the external solution to maximize this ion preference (5, 8, 9, 12, 18), although the presence of external  $Ca^{2+}$  does not prevent  $Ca^{2+}$  loss from the root (13). Anion uptake can also be maximized by addition of  $Ca^{2+}$  to the external solution (12, 13, 14). A case for increased prefer-




FIG. 3. Influence of  $\text{CaCl}_2$  concentration of absorption solutions on the Mg, K, Ca, and Cl contents in excised soybean roots. Solution composition: 5 mM KCl + 2.5 mM  $\text{MgCl}_2$  + 0 to 5 mM  $\text{CaCl}_2$ . Absorption period, 24 hr.

ence in uptake of 1 anion over another is not as obvious as for cations. The increased uptake in the presence of  $\text{Ca}^{2+}$  reported by Viets (18) seemingly applies to either cation or anion without a dependent coupling of the uptake for each ion.

Calcium in the external solution increased  $\text{K}^+$  uptake (5, 8, 9, 12, 14, 18) and decreased  $\text{Mg}^{2+}$  uptake (10, 17) by barley roots. Although  $\text{K}^+$  uptake by soybean roots was increased, the  $\text{Mg}^{2+}$  uptake was only slightly reduced in the presence of  $\text{Ca}^{2+}$ . Potassium was also an ineffective inhibitor of  $\text{Mg}^{2+}$  uptake. Addition of both  $\text{K}^+$  and  $\text{Ca}^{2+}$  to the  $\text{Mg}^{2+}$  solution drastically reduced  $\text{Mg}^{2+}$  uptake (Fig. 3). A  $\text{Ca}^{2+}$  concentration of 0.1 meq/liter was sufficient to invert the  $\text{Mg}^{2+}$  -  $\text{K}^+$  ratio value in the root when equivalent amounts of  $\text{K}^+$  and  $\text{Mg}^{2+}$  at a total concentration of 10 meq/liter were in the external solution. The time course of these interactions is shown in Fig. 4. The  $\text{Mg}^{2+}$  uptake is essentially complete within 1 hr. Likewise the loss of  $\text{Ca}^{2+}$ , presumably in exchange for  $\text{Mg}^{2+}$ , occurred during the first hr of the uptake period. Potassium uptake was positive for every sampling time in contrast to the observed loss in the absence of  $\text{Ca}^{2+}$ . The maximum  $\text{Ca}^{2+}$  effect near 1 meq/liter (Fig. 3) was also observed in excised barley roots for the uptake of  $\text{K}^+$  (5, 8),  $\text{SO}_4^{2-}$  (11), and  $\text{PO}_4^{3-}$  (12).

The interaction of  $\text{Mg}^{2+}$ ,  $\text{K}^+$ , and  $\text{Ca}^{2+}$  leading to the greater preference for  $\text{K}^+$  and a restriction of

$\text{Mg}^{2+}$  uptake occurs at the external solution-root surface interface. We have previously proposed that the interaction takes place at the epidermal cell layer (13), while others consider that intercellular interfaces are involved (9, 15). We find the following features of ion uptake by soybean roots placed in a  $\text{K}^+$  +  $\text{Ca}^{2+}$  +  $\text{Mg}^{2+}$  chloride solution (A) further  $\text{Ca}^{2+}$  accumulation in the roots is small (Fig. 3), (B) some changes in cation content take place quickly (Fig. 4), (C) the  $\text{Cl}^-$  increases parallel those of  $\text{K}^+$  (Fig. 3), and (D) the  $\text{Mg}^{2+}$  accumulation stops after 1 hr (Fig. 4). In these experiments,  $\text{Ca}^{2+}$  affected the influx of  $\text{Mg}^{2+}$ ,  $\text{K}^+$ , and  $\text{Cl}^-$  more than their efflux since the absorbed Mg could not be removed by incubation of Mg-loaded roots in a  $\text{Ca}^{2+}$  +  $\text{K}^+$  solution. Mengel and Helal (14) concluded that  $\text{Ca}^{2+}$  reduces the  $\text{K}^+$  efflux rather than increasing the influx thereby providing a greater net accumulation. The  $\text{Ca}^{2+}$  -  $\text{K}^+$  mediated effects on selectivity for  $\text{K}^+$  uptake relative to  $\text{Mg}^{2+}$  are not readily explainable by a change in relative flux rates for  $\text{K}^+$  uptake alone. Rather, calcium has some distinct influences on membrane properties upon which  $\text{K}^+$  effects are superimposed.




FIG. 4. Cation contents of roots incubated in Ca + Mg + K salt solutions as a function of time. Solution composition: 1 mM  $\text{CaCl}_2$  + 5 mM KCl + 2.5 mM  $\text{MgCl}_2$ .

Regulation of  $\text{Mg}^{2+}$  uptake by  $\text{Ca}^{2+}$  +  $\text{K}^+$  is clearly evident in the case of solutions of varying  $\text{Ca}^{2+}$  -  $\text{Mg}^{2+}$  ratios and constant ionic strength. In a solution with a total ( $\text{Mg}^{2+}$  +  $\text{Ca}^{2+}$ ) salt concentration of 10 meq/liter (Fig. 5) roots maintained their initial  $\text{K}^+$  content and increased their  $\text{Cl}^-$  content in the presence of calcium. In the absence of  $\text{Ca}^{2+}$ , the  $\text{K}^+$  and  $\text{Cl}^-$  contents approached zero. Calcium contents decreased, as would be expected from completion, as the  $\text{Ca}^{2+}$  -  $\text{Mg}^{2+}$  ratio decreased from 1 to 0. The inhibition of  $\text{Mg}^{2+}$  uptake by  $\text{Ca}^{2+}$ , however, was negligible as can be determined by comparing  $\text{Mg}^{2+}$  uptake at equal concentrations in Fig. 1 and 5, and as shown by the slight change in




FIG. 5. Influence of Ca/Mg on the Mg, K, Ca, and Cl contents of excised soybean roots in 5 mM (Ca + Mg)Cl<sub>2</sub> solutions for 24 hr.

Mg<sup>2+</sup> content at intermediate Ca<sup>2+</sup> - Mg<sup>2+</sup> ratios (Fig. 5). The Cl<sup>-</sup> content of the root remained below 20 meq/kg fresh weight in these experiments in which the Cl<sup>-</sup> concentration in the solution was 10 meq/liter and K<sup>+</sup> was not an initial constituent.

Potassium at 1 meq/liter in the 10-meq/liter Ca<sup>2+</sup> + Mg<sup>2+</sup> solutions strikingly shifted the root ion contents in the 24-hr experimental period (Fig. 6). The Ca<sup>2+</sup> contents remained essentially unaffected by the addition of KCl (*cf.* Fig. 5 and 6). The K<sup>+</sup> and Cl<sup>-</sup> contents approximately doubled over the range of 20- to 80-mole percent Ca<sup>2+</sup> in which range K<sup>+</sup>:Cl<sup>-</sup>:Ca<sup>2+</sup> ratios within each experiment were relatively constant (*cf.* Fig. 5 and 6). The Mg<sup>2+</sup> content in the presence of Ca<sup>2+</sup> and K<sup>+</sup>, on the other hand, decreased about 5-fold relative to the level in the absence of K<sup>+</sup>. With a Ca<sup>2+</sup> to Mg<sup>2+</sup> ratio of 1.0, the Ca<sup>2+</sup> and Mg<sup>2+</sup> contents were equal in the presence of K<sup>+</sup> (Fig. 6), but in the absence of K<sup>+</sup>, the Mg<sup>2+</sup> content of the roots exceeded the Ca<sup>2+</sup> level by about 5-fold (Fig. 5). In the absence of Ca<sup>2+</sup>, the Mg<sup>2+</sup> content exceeded K<sup>+</sup> by about




FIG. 6. Influence of Ca/Mg on the Mg, K, Ca, and Cl contents of excised soybean roots after 24 hr in 1 mM KCl and 5 mM (Ca + Mg)Cl<sub>2</sub> solutions. (Note the change in ordinate scale after Fig. 5.)

2- and 5-fold (Fig. 3 and 6). For the conditions represented in Fig. 6, the roots actually lost 11 µeq/g of their initial K<sup>+</sup> to the solution. When the Ca<sup>2+</sup> concentration in the solution was 0.1 meq/liter in the presence of 5 mM KCl + 2.5 mM MgCl<sub>2</sub>, that is, Ca<sup>2+</sup>/Mg<sup>2+</sup> = 0.02, the K<sup>+</sup> and Mg<sup>2+</sup> contents were about equal (Fig. 3). At Ca<sup>2+</sup>/Mg<sup>2+</sup> = 0.1, the K<sup>+</sup> exceeded Mg<sup>2+</sup> by 7-fold (Fig. 3).

Thus, it is evident that the predominance of K<sup>+</sup> over Mg<sup>2+</sup> uptake by soybean roots depends on the presence of Ca<sup>2+</sup> in the external solution. Also, restricted Mg uptake depends on the presence of both K<sup>+</sup> and Ca<sup>2+</sup> in the external solutions. These controlling interactions appear to be exerted on the solution side of the rate-limiting barrier for uptake, which might be considered as the solution side of a limiting membrane. Time courses of adjustments to initial conditions are correspondingly rapid as shown in Fig. 4. There is a symmetrical exchange of Ca<sup>2+</sup> and Mg<sup>2+</sup> which is essentially complete in 30 min, after which the uptake of Mg<sup>2+</sup> is very low and symmetrical with a small Ca<sup>2+</sup> loss (Fig. 4). In this 30-min period, the rapid initial change of K<sup>+</sup> content shifts to a steady increase, that is, a constant uptake rate.

*Uptake by Intact Plants.* Ion uptake by intact plants is affected by growth and transport from the root to the shoot. Kinetics of uptake are best followed on initially low-salt tissue in order to maximize rates of accumulation. Soybean plants can be grown on CaSO<sub>4</sub> solutions for about 2 weeks before mineral-deficiency symptoms appear. Roots obtained from either etiolated or green soybean plants cultivated in CaSO<sub>4</sub>, however, have relatively high salt contents compared with low-salt barley roots. The high level of K<sup>+</sup> in the tissue is due to the high K<sup>+</sup> content of the soybean seed.

Results in table II are for an uptake period of 24 hr from the indicated solutions for plants grown on 0.2 mM CaSO<sub>4</sub>, with the initial composition as shown in line 1 of the table. The several test solutions contained either nitrate (soln 1), nitrate + sulfate (soln 2), sulfate (soln 3), or chloride as predominant anions. The cation concentrations of both the root and shoot tissues increased by 1.2- to 2.0-fold, indicating that the plants shifted from a relatively low- to a high-salt state. Ion accumulation and transport are reflected rather than growth.

Effects of K<sup>+</sup> + Ca<sup>2+</sup> on restriction of Mg<sup>2+</sup> uptake (table II), although greatest in the root tissue, are also clearly evident in the shoot composition. The shoots, which were initially higher in Mg<sup>2+</sup> than the roots, probably did not approach a steady state for salt accumulation in the 24-hr test period. The Ca concentrations of the roots of the intact plants were similar to the excised ones in being relatively constant in the several solutions. The Ca<sup>2+</sup> concentrations in the shoot, on the other hand, increased greatly in soln 1 (0.1 Hoagland) and soln 2 (-K<sup>+</sup>) relative to the initial values. Continued uptake by and

Table II. Cation Content of  $\text{CaSO}_4$ -grown Plants After 24-Hr Uptake Period From Various Salt Solutions

Treatment	Tissue content							
	Root				Shoot			
	K	Mg	Ca	$\Sigma$	K	Mg	Ca	$\Sigma$
	<i>μeq/g fresh wt</i>							
	Initial content, prior to 24-hr uptake period							
	Content after 24-hr uptake period							
$\text{CaSO}_4$ , 0.2 mM	41	4	13	58	64	13	32	109
Soln 1 <sup>1</sup>								
[0.1 Hoagland (7)]	65	13	16	94	99	19	47	165
Soln 2 <sup>2</sup> (-K)	29	56	19	104	64	27	88	179
Soln 3 <sup>3</sup> (-N)	82	6	10	98	109	23	43	175
$\text{MgCl}_2$ , 10 mM	40	39	7	86	78	19	35	132
KCl, 10 mM	105	4	9	118	81	17	33	131

<sup>1</sup> Soln 1 = 0.5 mM  $\text{KNO}_3$ ; 0.1 mM  $\text{KH}_2\text{PO}_4$ ; 0.5 mM  $\text{Ca}(\text{NO}_3)_2$ ; 0.2 mM  $\text{MgSO}_4$ .

<sup>2</sup> Soln 2 = 0.05 mM  $\text{Ca}(\text{H}_2\text{PO}_4)_2$ ; 0.5 mM  $\text{Ca}(\text{NO}_3)_2$ ; 0.2 mM  $\text{MgSO}_4$ .

<sup>3</sup> Soln 3 = 0.25 mM  $\text{K}_2\text{SO}_4$ ; 0.05 mM  $\text{Ca}(\text{H}_2\text{PO}_4)_2$ ; 0.2 mM  $\text{CaSO}_4$ ; 0.2 mM  $\text{MgSO}_4$ .

movement of  $\text{Ca}^{2+}$  through the root as would be expected if  $\text{Ca}^{2+}$  is essentially excluded from the root vacuoles. This is based on the assumption that most of the  $\text{Cl}^-$  accumulates in the vacuoles and from the observed negligible  $\text{Cl}^-$  uptake by soybean roots from a  $\text{CaCl}_2$  solution.

A similar, more direct observation of  $\text{Ca}^{2+}$  uptake and movement of  $\text{Ca}^{2+}$  through the root was made by Moore, Mason, and Maas (16). They observed an initial rapid uptake of  $^{45}\text{Ca}$  by excised barley roots with a negligible accumulation, while  $\text{Ca}^{2+}$  in the exudate was 3- to 5-fold the external solution  $\text{Ca}^{2+}$  concentration. The epidermal and cortical root cells apparently do not accumulate  $\text{Ca}^{2+}$ . They suggest that the barrier to  $\text{Ca}^{2+}$  uptake and the concentrating step occurs at the endodermis upon transfer into the conducting vessels. Their results suggest that the accumulation of Ca found in plant shoots probably takes place in the root. One must add that the stem and leaf cells also accumulate  $\text{Ca}^{2+}$  in contrast to the negligible accumulation by the root cells.

Omission of  $\text{K}^+$  (soln 2), while greatly enhancing  $\text{Mg}^{2+}$  uptake relative to the complete nutrient (soln 1) did not significantly change the total cation content ( $\Sigma$ ). Omission of nitrate (soln 3), leaving sulfate as the predominant anion, reduced Mg uptake without essentially changing  $\Sigma$  of cations in roots or shoots. The balance in  $\Sigma$  of cations in the root was effected by the reciprocal changes in  $\text{K}^+$  and  $\text{Mg}^{2+}$ , with the effect of  $\text{Ca}^{2+}$  being important in the shoot. Absence of nitrate, however, enhanced  $\text{K}^+$  contents in both tissues.

We cannot fully assess the anion and cation interactions, which are incidental to the main purpose of the work. There is little doubt, however, that the  $\text{K}^+$ ,  $\text{Mg}^{2+}$ , and  $\text{Ca}^{2+}$  uptakes are interdependent and that these interactions are shown both by the excised roots and the

## Literature Cited

1. BAREER, D. A. AND H. V. KOONTZ. 1963. Uptake of dinitrophenol and its effect on translocation and calcium accumulation in barley. *Plant Physiol.* 38: 60-65.
2. COLLANDER, R. 1941. Selective absorption of cations by higher plants. *Plant Physiol.* 16: 691-720.
3. COTLOVE, E., H. V. TRANHAM, AND R. L. BOWMAN. 1958. An instrument and method for automatic, rapid, accurate, and sensitive titration of chloride in biological samples. *J. Lab. Clin. Med.* 51: 461.
4. DAVID, D. J. 1960. The determination of exchangeable sodium, potassium, calcium, and magnesium in soils by atomic-absorption spectrophotometry. *Analyst* 85: 495-503.
5. EPSTEIN, E. 1961. The essential role of calcium in selective cation transport by plant cells. *Plant Physiol.* 36: 437-45.
6. EPSTEIN, E. AND J. E. LEGGETT. 1954. The absorption of alkaline earth cations by barley roots: kinetics and mechanism. *Am. J. Botany* 41: 785-91.
7. HOAGLAND, D. R. AND D. I. ARNON. 1950. The water-culture method for growing plants without soil. *California Agr. Expt. Sta. Cir.* 347.
8. JACOBSON, L., R. J. HANNAPEL, D. P. MOORE, AND M. SCHAEDELE. 1961. Influence of calcium on selectivity of ion absorption process. *Plant Physiol.* 36: 58-65.
9. JACOBSON, L., D. P. MOORE, AND R. J. HANNAPEL. 1960. Role of calcium in absorption of monovalent cations. *Plant Physiol.* 35: 352-58.
10. LAZAROFF, N. AND M. G. PITMAN. 1966. Calcium and magnesium uptake by barley seedlings. *Australian J. Biol. Sci.* 19: 991-1005.
11. LEGGETT, J. E. AND E. EPSTEIN. 1956. Kinetics of sulfate absorption by barley roots. *Plant Physiol.* 35: 326-32.
12. LEGGETT, J. E., R. A. GALLOWAY, AND H. G. GAUCH. 1965. Calcium activation of orthophosphate by barley roots. *Plant Physiol.* 40: 897-902.
13. LEGGETT, J. E. AND W. A. GILBERT. 1967. Localization of the Ca-mediated apparent ion selectivity in the cross-sectional volume of soybean roots. *Plant Physiol.* 42: 1658-64.
14. MENGEL, K. AND M. HELAL. 1967. Der Einfluss des austauschbaren  $\text{Ca}^{2+}$  junger Gerstenwurzeln auf den Flux von  $\text{K}^+$  und Phosphat—eine Interpretation des Viets-Effektes. *Z. Pflanzenern., Dung., Boden* 57: 223-34.
15. MOORE, D. P., L. JACOBSON, AND R. OVERSTREET. 1961. Uptake of calcium by excised barley roots. *Plant Physiol.* 36: 53-58.
16. MOORE, D. P., B. J. MASON, AND E. V. MAAS. 1965. Accumulation of calcium in exudate of individual barley roots. *Plant Physiol.* 40: 641-44.
17. MOORE, D. P., R. OVERSTREET, AND L. JACOBSON. 1961. Uptake of magnesium and its interaction with calcium in excised barley roots. *Plant Physiol.* 36: 290-96.
18. VIETS, F. G., JR. 1944. Calcium and other polyvalent cations as accelerators of ion accumulation by excised barley roots. *Plant Physiol.* 19: