

Characterization of Photosynthetic Rhythms in Marine Dinoflagellates

I. PIGMENTATION, PHOTOSYNTHETIC CAPACITY AND RESPIRATION¹

Received for publication January 26, 1977 and in revised form May 12, 1977

BARBARA B. PRÉZELIN, BLANCHE W. MEESON, AND BEATRICE M. SWEENEY

Department of Biological Sciences, University of California, Santa Barbara, California 93106

ABSTRACT

Circadian rhythms in photosynthesis were defined for the first time in the dinoflagellates *Glenodinium sp.* (M. Bernard strain) and *Ceratium furca* Ehrenberg (B. Meeson strain) and compared with that in *Gonyaulax polyedra* Stein. All three phytoplankton species had photosynthetic rhythms with daily amplitudes ranging from 3 to 5 and maxima displayed about midday. The photosynthetic pigment content and absorption properties of the cells were constant over the circadian cycle. Diurnal periodicities in respiration never accounted for more than 30% of the photosynthetic rhythm and did not persist under constant conditions. There was sufficient similarity between the circadian rhythms of these three dinoflagellates to suggest the mechanism of regulation may be the same for each of them.

yaulax polyedra (8), the photosynthetic rhythm has been shown to be controlled by a biological clock. In these cases, rhythmicity is an endogenous characteristic of the cell and occurs independently of environmental variables (21, 23). Therefore, photosynthesis not only exhibits rhythmicity when cells are cultured under light-dark cycles (LD), but this rhythmicity persists with a period of about 24 hr when cells are transferred to constant conditions of continuous dim illumination (LL). Unfortunately, few measurements have been made to test whether the diurnal periodicity of photosynthesis observed in natural populations persists under constant conditions.

In the present laboratory study, the circadian natures of the photosynthetic rhythm in three marine dinoflagellates are compared and demonstrated for the first time in two of them. The results indicate that the photosynthetic rhythm of the type previously documented in *G. polyedra* may be widespread among the dinoflagellates and that further work on the regulation of the photosynthesis rhythm in *G. polyedra* should be broadly significant.

Diurnal periodicity in photosynthetic capacity (P_{max})² has been documented during the last 20 years for several groups of algae and mixed populations of phytoplankton (6, 12, 13, 21). In these organisms, maximal productivity is restricted to a short time each day, and thus the periodic nature of photosynthesis complicates field measurements of primary production. It appears clear from the extensive data now available that environmental variables significantly affect the magnitude of photosynthesis, but have little or no effect on the timing of the photosynthetic maximum each day. Recently, Stross and his co-workers (2, 22) demonstrated a coincidence between the daily maximum in photosynthesis in *Euglena* and that in the rates of phosphate uptake in phosphorous-sufficient cultures. It appears unlikely that phosphorous availability serves to regulate the timing of the photosynthetic periodicity, since the maximal phosphate uptake rates occurred 6 hr after that in photosynthesis, when cultures were phosphorous-limited (3).

A few marine algae have been examined extensively for endogenous features which might account for their photosynthetic rhythm (4, 8, 9, 14, 18, 19, 21, 25-28). Although several of these studies demonstrate a daily periodicity in various photosynthetic parameters correlated with the cell cycle of synchronized cultures (4, 17, 18, 28), it is not clear that these organisms divide either once a day or in synchrony in natural populations. In the case of the green alga *Acetabularia* (9, 26), the diatom *Phaeodactylum tricornutum* (14), and the dinoflagellate *Gon-*

MATERIALS AND METHODS

Algae used in this study and their origins were as follows: *Glenodinium sp.* (L. Provasoli, M. Bernard strain, UCSB code No. 5M29), *G. polyedra* Stein clone 70A (USCB code No. 5M20), and *Ceratium furca* Ehrenberg (B. Meeson strain, UCSB code No. 5M30). All dinoflagellates were grown in unialgal batch cultures in 2.8-liter Fernbach flasks containing 1 to 1.5 liters of half-strength *f* medium (*f*/2) (7). In addition, medium used for culturing *C. furca* contained 65 μM sodium glycerophosphate, in place of 47 μM NaH_2PO_4 , and 200 μM nitrilotriacetic acid (NTA). Overhead illumination was provided by banks of cool-white fluorescent lamps. Irradiance was measured with a United Detector Technology model 40A light meter. Cells were cultured on an alternating 12-hr light-dark schedule (LD) at $21 \text{ C} \pm 1 \text{ C}$ and $1,000 \mu\text{w}/\text{cm}^2$ (70 ft-c). In experiments designed to test the circadian rhythmicity of a cell parameter, cultures were transferred at the beginning of the light period (0000 "circadian time" or ct 0) to continuous light (LL) at an irradiance between 400 and 500 $\mu\text{w}/\text{cm}^2$. Growth was assessed by cell counts made with an AO Bright Line hemocytometer or Hauser Scientific counting chamber.

Pigments were extracted after the methods of Jeffrey (10). Details of extraction, TLC, and extinction coefficients have been described previously (15).

For photosynthetic measurements, freshly harvested cells (3 min, 1,300 rpm, room temperature) were suspended in a small volume of fresh *f*/2 medium with the addition of 5 to 10 mM NaHCO_3 (pH 8). Dilute suspensions used for O_2 production measurements contained between 2.5 and 5 μg Chl *a*/5-ml sample. Prior to photosynthetic measurements, O_2 levels in the

¹ This investigation was supported by National Science Foundation Grant BMS 72-02221A03.

² Abbreviations: P_{max} : photosynthetic capacity; LD: light-dark cycles; LL: continuous illumination; ct: circadian time; UCSB: University of California, Santa Barbara algal culture collection.

sample were reduced by passing N_2 gas over the surface of the cell suspension. The sample was stoppered and kept at 20 C in the dark for 15 to 30 min. Photosynthetic O_2 evolution was measured polarographically with a Delieu and Walker (5) arrangement of the Clark-type O_2 electrode by the experimental procedure previously described (15). Temperature was controlled in the reaction vessel at 20 C by a constant temperature water bath (Forma Scientific Inc.). Collimated light was supplied by a cooled 500 w tungsten bulb from a Viewlex model V-25 slide projector and was filtered through 5 cm of 1% cupric sulfate solution. The irradiance of the incident beam was varied by placing wire mesh screens between the lamp and the reaction chamber.

RESULTS

Circadian rhythms of photosynthesis were defined for the first time in the dinoflagellates *Glenodinium sp.* and *C. furca* and compared with the well defined system in *G. polyedra*. Diurnal periodicity of P_{max} was established in LD cultures (Figs. 1a, 2a, 3a) with measurements made at 6-hr intervals, corrected for dark respiration, and expressed both on a cellular and Chl *a* basis. Each dinoflagellate displayed maximum photosynthetic capability at midday, followed by a 2- to 6-fold decline in the rates of O_2 evolution to a minimum during the subsequent dark period. Increased rates of O_2 evolution were reinitiated shortly prior to the onset of the next light period. The circadian nature of these fluctuations in P_{max} was established in the same LD cultures by transferring them to LL. Rates of O_2 evolution continued to demonstrate a rhythmicity with a period of approximately 24 hr and were maintained for several days under continuous dim illumination (Figs. 1a, 2a, 3a). During this time and




FIG. 1. a: Rhythms of photosynthetic capacity in *Glenodinium sp.*, expressed as $\mu\text{mol } O_2/\text{cell}\cdot\text{hr}$ (●—●) or as $\mu\text{mol } O_2/\mu\text{mol Chl } a\cdot\text{hr}$ (○—○). Light-dark cycle was 12:12 hr; irradiance of light phase was $1,000 \mu\text{w}/\text{cm}^2$. Dark periods are indicated by hatched bars on abscissa. Cultures were transferred to continuous dim illumination ($500 \mu\text{w}/\text{cm}^2$) at circadian hr 00. Cell division occurred at beginning of light period of light-dark cycle and ceased upon transfer to constant dim light. Cells were in midexponential phase of growth. b: Whole cell pigmentation of *Glenodinium sp.* over circadian time. Concentrations of peridinin (●), Chl *a* (Δ), and Chl *c*₂ (▲) are expressed as $\mu\text{mol} \times 10^{-10}/\text{cell}$.




FIG. 2. a: Rhythms of photosynthetic capacity in *G. polyedra*, expressed as $\mu\text{mol } O_2/\text{cell}\cdot\text{hr}$ (●—●) or as $\mu\text{mol } O_2/\mu\text{mol Chl } a\cdot\text{hr}$ (○—○). Light-dark cycle was 12:12 hr; irradiance of light phase was $800 \mu\text{w}/\text{cm}^2$. Dark periods are indicated by hatched bars on abscissa. Cultures were transferred to continuous dim illumination ($500 \mu\text{w}/\text{cm}^2$) at circadian hr 00. Cell division occurred at beginning of light period and ceased upon transfer to constant dim light. Cells were in late exponential phase of growth. b: Whole cell pigmentation of *G. polyedra* over circadian time. Concentrations of peridinin (●), Chl *a* (Δ), and Chl *c*₂ (▲) are expressed as $\mu\text{mol} \times 10^{-7}/\text{cell}$.

under the conditions of continuous dim illumination, cell division ceased. Although cells were viable and photosynthetically active, the photosynthetic rhythm of these dinoflagellates is not closely linked to their cell cycle. Unlike O_2 evolution, dark respiration rates displayed diurnal periodicity in LD cultures which appeared to damp out upon transfer of the cultures to constant illumination (Fig. 3a).

Although a difference of almost two orders of magnitude was noted in the cellular photosynthetic capacities of the three species of dinoflagellates examined, the amplitudes of the rhythms were approximately the same for *Glenodinium sp.*, *C. furca*, and *G. polyedra* cultured under similar growth conditions averaging at about 3.4, 3, and 5.2, respectively. Apparently unrelated to circadian control, the differences in P_{max} rates between the dinoflagellate species were correlated with differences in such cell parameters as cell size, cell volume, photosynthetic pigment content, and perhaps vacuolar space (inversely related to cell density) (Table I).

P_{max} within a single species of dinoflagellate was dependent on irradiances during growth. While photosynthetic rhythms persisted with roughly the same amplitude, the magnitude of the photosynthetic capacity was lower in cultures of *G. polyedra* grown at irradiances below $1,000 \mu\text{w}/\text{cm}^2$. Light-saturated photosynthetic rates were reduced by a factor of 5, from 200 to $40 \times 10^{-7} \mu\text{mol } O_2/\text{cell}\cdot\text{hr}$, when cells were cultured at $800 \mu\text{w}/\text{cm}^2$ rather than at $1,200 \mu\text{w}/\text{cm}^2$ at ct 6 on the same day (Table I). The photosynthetic stress response of *G. polyedra* to lowered irradiance has been further defined and will be described elsewhere (B. Prézélin and B. Sweeney, in preparation).

At the same time that photosynthetic capacity was being measured, the photosynthetic pigment content of an aliquot of the same culture was analyzed by acetone-extraction and TLC.

Although some alterations with time were observed, the cellular concentrations of Chl *a*, Chl *c*₂, and peridinin did not fluctuate with a period of 24 hr (Figs. 1b, 2b, 3b). The only significant fluctuations in pigment content occurred as an irradiance-induced response when cultures were transferred to constant dim illumination. Since incident irradiance of LL was about half that of LD, the total irradiance per day was approximately equivalent under both LD and LL conditions. Cellular pigment content of *G. polyedra* and *C. furca* was most sensitive to the transfer and began to increase markedly 6 to 12 hr after being placed in LL. Pigment concentrations reached a maximum after 12 to 24 hr, and returned to values comparable to those measured in the LD cultures after 24 to 48 hr in LL. Additional experimental measurements on similarly cultured dinoflagellates indicated that the pigment content of all three genera was stable and nonperiodic

in LD cultures and in LL cultures which had been in constant illumination for a day or 2.

DISCUSSION

The presence of circadian rhythms in photosynthesis has been established for three genera of marine dinoflagellates, *G. polyedra* and now *Glenodinium sp.* and *Ceratium furca*. There is sufficient similarity between the rhythms of these three dinoflagellates to suggest that the mechanisms of circadian regulation of photosynthesis may be the same for each of them. The maximal rates of photosynthetic O₂ evolution occurs about midday, *C. furca* displaying a maximum during the late morning (ct 5) and a few hr before the occurrence of the maxima of *Glenodinium sp.* (ct 7) and *G. polyedra* (ct 7) (B. Meeson, unpublished). Although the magnitudes of the rhythms differ in relation to cell size and culture condition, each of the dinoflagellates examined shows a 3- to 5-fold change in magnitude over the day and a free running periodicity of about 24 hr. In no species does rhythmicity depend on fluctuations in cell absorption (16) or photosynthetic pigment content of the cells (Figs. 1b, 2b, 3b), these cell parameters being found to be constant in photoadapted log phase cultures. The photosynthetic rhythms are detected when photosynthetic O₂ evolution is calculated either on a cellular or Chl basis (Figs. 1a, 2a, 3a). These characteristics are consistent with the circadian rhythmicity of photosynthesis described for *Phaeodactylum tricoratum* (14) and *Acetabularia crenulata* (9, 26) and the diurnal fluctuations in photosynthesis observed in *Chlorella pyrenoidosa* (20) and *Scenedesmus obliquus* (17-19).

In addition, the three dinoflagellates show diurnal periodicities in respiration which can never account for more than 30% of the photosynthetic rhythm and which did not persist under constant conditions. It is clear that variations in the rate of respiration cannot account for the characteristic rhythm observed in photosynthesis. Like *G. polyedra* (24), *C. furca* also displays a circadian rhythm in cell division, the maximum number of early division pairs appearing just prior to the beginning of the light period (B. Meeson, unpublished). The greatest number of dividing pairs of *Glenodinium sp.* also occurs at the beginning of the light period in cultures entrained to a 12:12 hr LD cycle, but the circadian rhythmicity of its cell division has not been tested.

The present studies establish *G. polyedra* as a test organism representative of those dinoflagellates which display a daily rhythmicity in their photosynthetic capacity. Extended studies on *G. polyedra* are being conducted to examine more fully how the regulation of photosynthetic capacity is mediated by a biological clock. Some of those results are presented in a following study (16). Finally, increasing documentation of *in situ* diurnal periodicity in photosynthesis and Chl fluorescence (11) requires a closer examination of natural populations under constant conditions, to define what role biological clocks may play in regulating primary production.




FIG. 3. a: Rhythms of photosynthetic capacity in *C. furca*, expressed as $\mu\text{mol O}_2/\text{cell}\cdot\text{hr}$ (●—●) or as $\mu\text{mol O}_2/\mu\text{mol Chl } a/\text{hr}$ (○—○). Respiration is expressed as $\mu\text{mol O}_2/\text{cell}\cdot\text{hr}$ (■—■). Light-dark cycle was 12:12 hr; irradiance of light phase was $1,000 \mu\text{w}/\text{cm}^2$. Dark periods are indicated by hatched bars on abscissa. Cultures were transferred to continuous dim illumination ($500 \mu\text{w}/\text{cm}^2$) at circadian hr 00. Cell division occurred at beginning of light period of light-dark cycle and ceased upon transfer to constant dim light. Cells were in midexponential phase of growth. b: Whole cell pigmentation of *C. furca* over circadian time. Concentrations of peridinin (●), Chl *a* (△), and Chl *c*₂ (▲) are expressed as $\mu\text{mol} \times 10^{-8}/\text{cell}$.

Table I. Comparison of cell parameters and photosynthetic rates of three marine dinoflagellates.

Species Growth Irradiance	Cell Size	Cell Volume ($\text{cm}^3/\text{cell} \times 10^{10}$)	Cell Density ($\text{gm}/\text{cell} \times 10^8$)	chl <i>a</i> /cell ($\mu\text{mol} \times 10^{10}$)	P _{max} /cell ¹ ($\times 10^7$)	P _{max} /cm ³	P _{max} /chl <i>a</i>
<i>Glenodinium sp.</i> 1200 $\mu\text{w}/\text{cm}^2$	16 μ X 11 μ	60	5.0 - 8.3	15	3.5	58	23
<i>G. polyedra</i> 1200 $\mu\text{w}/\text{cm}^2$	60 μ X 40 μ	540	2.22	760	175 - 225	32 - 42	23 - 30
800 $\mu\text{w}/\text{cm}^2$		410		500	40	10	8
<i>C. furca</i> 1200 $\mu\text{w}/\text{cm}^2$	130 μ X 135 μ	800	1.37	500	300	38	60

¹P_{max} is measured as $\mu\text{moles oxygen/hr}$ at $5000 \mu\text{w}/\text{cm}^2$ and is corrected for dark respiration.

LITERATURE CITED

1. BUTLER WL 1972 Absorption spectra of biological materials. *Methods Enzymol* 23: 3-25
2. CHISHOLM SW, RG STROSS 1976 Phosphate uptake kinetics in *Euglena gracilis* (Z) (Euglenophyceae) grown on light/dark cycles. I. Synchronized batch cultures. *J Phycol* 12: 210-217
3. CHISHOLM SW, RG STROSS 1976 Phosphate uptake kinetics in *Euglena gracilis* (Z) (Euglenophyceae) grown on light/dark cycles. II. Phased PO_4 -limited cultures. *J Phycol* 12: 217-222
4. CODD GA, MJ MENRET 1971 Photosynthetic products of division synchronized cultures of *Euglena*. *Plant Physiol* 47: 635-639
5. DELIEU T, DA WALKER 1972 An improved cathode for the measurement of photosynthetic oxygen evolution by isolated chloroplasts. *New Phytol* 71: 201-225
6. DOTY MS, M OGURI 1957 Evidence for a photosynthetic daily periodicity. *Limnol Oceanogr* 2: 37-40
7. GUILLARD RRL, JH RHYTHER 1962 Studies of marine planktonic diatoms. I. *Cyclotella nana* Hustedt and *Detonula confervaceae* (Cleve). *Gran Can J Microbiol* 8: 229-239
8. HASTINGS JW, L ASTRACHAN, BM SWEENEY 1961 A persistent daily rhythm in photosynthesis. *J Cell Physiol* 45: 69-76
9. HELLEBUST JA, J TERBORGH, GC MCLEOD 1967 The photosynthetic rhythm of *Acetabularia crenulata*. II. Measurements of photoassimilation of carbon dioxide and the activities of enzymes of the reductive pentose cycle. *Biol Bull* 133: 670-678
10. JEFFREY SW 1968 Quantitative thin-layer chromatography of chlorophylls and carotenoids from marine algae. *Biochim Biophys Acta* 162: 271-285
11. KIEFER DA 1973 Fluorescence properties of natural phytoplankton populations. *Mar Biol* 22: 263-269
12. LORENZEN CJ 1963 Diurnal variation in photosynthetic activity of natural phytoplankton populations. *Limnol Oceanogr* 8: 56-62
13. MALONE TC 1971 Diurnal rhythms in net plankton and nannoplankton assimilation ratios. *Mar Biol* 10: 285-289
14. PALMER JD, L LIVINGSTON, FD ZUSY 1964 A persistent diurnal rhythm in photosynthetic capacity. *Nature* 203: 1087-1088
15. PRÉZELIN BB 1976 The role of peridinin-chlorophyll *a*-proteins in the photosynthetic light adaptation of the marine dinoflagellate *Glenodinium* sp. *Planta* 128: 133-141
16. PRÉZELIN BB, BM SWEENEY 1977 Characterization of photosynthetic rhythms in marine dinoflagellates. II. Photosynthesis-irradiance curves and *in vivo* chlorophyll *a* fluorescence. *Plant Physiol* 60: 388-392
17. SENGER H 1970 Characterization of a synchronous culture of *Scenedesmus obliquus*: its potential photosynthetic capacity and its photosynthetic quotient during the life cycle. *Planta* 90: 243-266
18. SENGER H 1970 Quantum yield and variable behavior of the two photosystems of the photosynthetic apparatus during the life cycle of *Scenedesmus obliquus* in synchronous culture. *Planta* 92: 327-346
19. SENGER H, NI BISHOP 1967 Quantum yield of photosynthesis in synchronous *Scenedesmus* culture. *Nature* 214: 140-143
20. SOROKIN C 1960 Photosynthetic activity in synchronized cultures of algae and its dependence on light intensity. *Arch Mikrobiol* 37: 151-160
21. SOURNIA A 1974 Circadian periodicities in natural populations of marine phytoplankton. *Adv Mar Biol* 12: 325-389
22. STROSS RG, SW CHISHOLM, TA DOWNING 1963 Causes of daily rhythm in photosynthetic rates of phytoplankton. *Biol Bull* 145: 200-209
23. SWEENEY BM 1960 The photosynthetic rhythm in single cells of *G. polyedra*. *Cold Spring Harbor Symp Quant Biol* 25: 145-148
24. SWEENEY BM, HASTINGS JW 1958 Rhythmic cell division in populations of *Gonyaulax polyedra*. *J Protozool* 5: 217-221
25. SWEENEY BM, FT HAXO 1961 Persistence of a photosynthetic rhythm in enucleated *Acetabularia*. *Science* 134: 1361-1363
26. TERBORGH J, GC MCLEOD 1967 The photosynthetic rhythm of *Acetabularia crenulata*. I. Continuous measurements of oxygen exchange in alternating light-dark regimes and in constant light at different intensities. *Biol Bull* 133: 659-669
27. VANDEN DRIESSCHE T 1966 Circadian rhythms in *Acetabularia*. Photosynthetic capacity and chloroplast shape. *Exp Cell Res* 42: 18-30
28. WALTHER WG, LN EDMUNDS JR 1973 Studies on the control of the rhythm of photosynthetic capacity in synchronized cultures of *Euglena gracilis*. *Plant Physiol* 51: 250-258