

**Short Communication**

# Evidence for Control of Carbon Partitioning by Fructose 2,6-Bisphosphate in Spinach Leaves<sup>1</sup>

Received for publication October 4, 1983 and in revised form December 5, 1983

STEVEN C. HUBER\* AND D. MARK BICKETT

*United States Department of Agriculture, Agricultural Research Service, Departments of Botany and Crop Science, North Carolina State University, Raleigh, North Carolina 27650*

## ABSTRACT

Excision of spinach (*Spinacia oleracea* L.) leaves had no effect on photosynthetic rates, but altered normal carbon partitioning to favor increased formation of starch and decreased formation of sucrose. The changes were evident within 2 hours after excision. Concurrently, leaf fructose-2,6-bisphosphate content increased about 5-fold (from 0.1 to 0.5 nanomoles per gram fresh weight). The activities of sucrose-P synthase and cytoplasmic fructose 1,6-bisphosphatase in leaf extracts remained constant during the time period tested. It is postulated that the rise in fructose 2,6-bisphosphate was responsible for the change in carbon partitioning.

## MATERIALS AND METHODS

**Materials.** Fresh beef liver for preparation of phosphofructokinase was obtained locally. All other reagents and chemicals were obtained from Sigma.

**Plant Growth and Experimental Protocol.** Spinach (*Spinacia oleracea* L.) plants were grown in the greenhouse during the months of April and May. Fully expanded leaves were excised at 0900 h on clear, sunny days. The petioles were recut while submerged in degassed distilled H<sub>2</sub>O, placed in tubes of the same water, and positioned into full sunlight. Several leaves were harvested just after excision and at various times over the 4.5-h experimental period. Prior to each harvest, leaf CER was measured. After the CER measurement, four leaf punches (2.0 cm<sup>2</sup> total) were taken for starch and soluble sugar analysis. The midrib and petiole were removed from the leaf and the remaining tissue weighed and sliced. A subsample (0.5 g) of tissue was taken for F26BP determinations, and the rest was frozen at -80°C for SPS and FBPase measurements.

**Photosynthesis Measurements.** CER was measured using an Anarad<sup>3</sup> differential IR CO<sub>2</sub> analyzer equipped with a clamp-on plexiglass cuvette that enclosed 5 cm<sup>2</sup> of the upper and lower leaf surfaces. Ambient air was passed through the cuvette at a flow rate of 1.5 l/min, and the CO<sub>2</sub> differential of incoming and outgoing air was used to determine the CER.

**Starch and Sugar Analysis.** Leaf discs were extracted with hot 80% ethanol until the tissue was pigment free. The particulate fraction was analyzed for starch after digestion with amyloglucosidase (8). The supernatants were evaporated to dryness. The residues were resuspended in water and assayed for hexose and sucrose (8).

**Leaf F26BP Determination.** Immediately after harvest, 0.5 g of tissue was ground in a mortar with 2 ml of ice cold buffer containing 50 mM HEPES-NaOH (pH 7.5), 10 mM KF, 2.5 mM DTT, and 2 mM EDTA. The extract was placed in a boiling water bath for 2 min and then centrifuged at 32,000g for 5 min. The supernatant was used for F26BP analysis. Using these procedures, recovery of authentic F26BP added to the grind medium before homogenization of leaf tissue averaged 70%.

The determination of F26BP content was based on the ability of the extract to relieve the inhibition of phosphofructokinase by ATP. Phosphofructokinase was partially purified from fresh beef liver, and the F26BP assay was performed essentially as described by Furuya and Uyeda (4). The assay mixture contained 50 mM

Partitioning of photosynthetically fixed carbon between starch and sucrose can be altered by 'source-sink' manipulations (5). When 'sink demand' for sucrose was decreased by girdling (9), fruit removal (1), or leaf excision (8), photosynthetic rates were ultimately reduced while starch accumulation either increased or was similar to that in control plants. Hence, relative partitioning of carbon into starch was increased.

In short-term leaf excision experiments for soybean plants (8), the decreased rate of sucrose accumulation was attributed to a reduction in the maximum activity of SPS<sup>2</sup>, the enzyme responsible for photosynthetic sucrose formation. Within 2.5 h after leaf excision, SPS activity was reduced to only 26% of the activity at the time of detachment (8).

In preliminary experiments with detached spinach leaves, changes in carbon partitioning were observed, but the maximum activity of SPS was not decreased. This is documented in the present report and, in addition, we show that the change in partitioning in spinach leaves was associated with a dramatic increase in F26BP content. F26BP is a potent inhibitor of cytoplasmic FBPase (2, 11), and the results obtained are consistent with a regulatory role for this metabolite in the sucrose formation pathway.

<sup>1</sup> Cooperative investigations of the North Carolina Agricultural Research Service and the Agricultural Research Service, United States Department of Agriculture, Raleigh, NC. Paper No. 9017 of the Journal Series of the North Carolina Agricultural Research Service, Raleigh, NC.

<sup>2</sup> Abbreviations: SPS, sucrose-P synthase; F26BP, fructose 2,6-bisphosphate; FBPase, fructose 1,6-bisphosphatase; CER, carbon exchange rate (mg CO<sub>2</sub>/dm<sup>2</sup>·h); FBP, fructose 1,6-bisphosphate.

<sup>3</sup> Mention of a trademark or proprietary product does not constitute a guarantee or warranty of the product by the United States Department of Agriculture and does not imply its approval to the exclusion of other products that may also be suitable.

Hepes (pH 7.4), 1 mM EDTA, 5 mM MgCl<sub>2</sub>, 1 mM NH<sub>4</sub>Cl, 0.16 mM NADH, 2.5 mM DTT, 0.5 units aldolase, 5.8 units triose-P-isomerase, 0.5 units  $\alpha$ -glycerol-P-dehydrogenase, 20 units of phosphofructokinase, and 20  $\mu$ l of spinach extract. After 2 min, the reaction was initiated by addition of 1.2 mM ATP and monitored by following the decrease in absorbance at A<sub>340</sub>. All rate measurements were taken 4 min after initiation, and F26BP concentration was determined with respect to a standard curve. Acid treatment of samples (4) completely eliminated the apparent F26BP activity.

**SPS and FBPase Assays.** Frozen tissue was ground (1 g fresh wt/8 ml medium) with a mortar and pestle in 50 mM Hepes-NaOH (pH 7.5), 5 mM MgCl<sub>2</sub>, 1 mM EDTA, 2.5 mM DTT, and 0.5% BSA. The extracts were filtered through eight layers of cheese cloth and clarified by centrifugation at 32,000g for 5 min. SPS was assayed by measuring fructose-6-P-dependent formation of sucrose (+ sucrose-P) from UDP-glucose as previously described (7). Cytoplasmic FBPase was assayed by a continuous spectrophotometric assay. The 1.0-ml reaction mixture contained 50 mM Hepes-NaOH (pH 7.0), 5 mM MgCl<sub>2</sub>, 10  $\mu$ M FBP, 0.2 mM NADP, 2 units each of phosphoglucosomerase and glucose-6-P dehydrogenase, and an aliquot of leaf extract. Under these assay conditions, activity of chloroplast FBPase is negligible and there was virtually no activity in the absence of Mg<sup>2+</sup> (*i.e.* nonspecific phosphatase).

## RESULTS

Detached spinach leaves continued to photosynthesize at rates similar to those of attached (control) leaves during the 4-h test period. CER was constant at about 9 mg CO<sub>2</sub>/g fresh wt · h. In the attached leaves, there was a gradual increase with time in leaf sucrose concentration (Fig. 1A). Excised leaves, in contrast, accumulated sucrose at an accelerated rate for the first 2 h. Thereafter, sucrose continued to accumulate, but at a reduced rate (Fig. 1A), which corresponded with a marked increase in the rate of starch accumulation (Fig. 1B). There was no accumulation of carbohydrates in the midrib or petioles of detached leaves (data not shown). Hence, translocation was effectively blocked by leaf excision. The results clearly indicated that, after 2 h, carbon partitioning was altered to favor increased formation of starch at the expense of sucrose formation.

The decrease in sucrose formation was not the result of changes in the maximum activities of SPS or cytoplasmic FBPase, as attached and detached leaves had similar enzyme activities in leaf extracts (Table I). However, leaf excision resulted in a dramatic increase in leaf F26BP content (Fig. 2). Attached leaves maintained a relatively low and constant level of F26BP (about 0.15 nmol/g fresh weight). In detached leaves, the F26BP content increased rapidly so that, after 2 h, leaves contained about 0.5 nmol F26BP/g fresh weight.

## DISCUSSION

Excision of spinach leaves resulted in an initial accumulation of sucrose that preceded the change in carbon partitioning. Similarly, Foyer *et al.* (3) reported that sucrose pretreatment of protoplasts from mature spinach leaves inhibited subsequent sucrose formation. However, with the protoplast system, the sucrose pretreatment also inhibited CO<sub>2</sub> fixation whereas, in the present study with intact leaves, the change in carbon partitioning occurred while CER remained constant.

It is unlikely that in excised leaves the reduction in sucrose accumulation after 2 h was caused by direct feedback inhibition by sucrose of enzymes in the sucrose pathway (3, 7). Similarly, the effect cannot be ascribed to reduced activities (measured *in vitro*) of SPS or cytoplasmic FBPase (Table I), as was the case with detached soybean leaves (8). It is likely that the increase in




FIG. 1. Changes in (A) leaf sucrose content and (B) leaf starch accumulation of excised (○) and attached, control (▲) leaves of spinach plants. Rates of carbohydrate accumulation, expressed as mg/g fresh wt · h, are shown parenthetically.

Table I. Effect of Excision on Activity of Certain Enzymes in Spinach Leaf Extracts

Tissue	Elapsed Time <sup>a</sup>	Enzyme	
		SPS	cyto FBPase
	h	$\mu$ mol product/g fresh wt · h	
Attached	0	16.1 ± 2.5	24.7 ± 1.6
Attached	4	13.8 ± 2.0	23.6 ± 1.5
Detached	4	14.7 ± 2.3	27.5 ± 3.4

<sup>a</sup> The experiment was started at time zero, which was about 0900 h, and lasted 4 h.

F26BP concentration (Fig. 2) was responsible for the reduction in sucrose formation. F26BP is a potent inhibitor of cytoplasmic FBPase (2, 11). Using conversion factors developed by Stitt *et al.* (12), and assuming strict compartmentation, the concentration of F26BP in the cytoplasm of detached leaves increased from about 1  $\mu$ M to a maximum level of about 5  $\mu$ M. This change in F26BP concentration would likely cause a substantial inhibition of FBPase activity (2).

It is intriguing to note that apparent species differences may exist in the mechanisms employed to reduce the rate of sucrose formation in response to a block in export. The results presented




FIG. 2. Changes in F26BP content of excised (O) and attached, control (▲) leaves for spinach.

herein suggest that spinach plants modulate the level of the regulatory metabolite F26BP. In contrast, the maximum activity of SPS is regulated in soybean leaves (8). It is not known whether there are changes in the concentration of F26BP in soybean leaves under these conditions (we have encountered technical difficulties in extracting F26BP from soybean leaves).

F26BP plays an important role in regulating glycolysis and gluconeogenesis in the liver cell, and its concentration is hormonally controlled (6). It appears that F26BP may play a similar role in the leaf cell. To date, it is known that F26BP is present

in leaves and that it is a potent inhibitor of cytoplasmic FBPAse (2, 11). In addition, Stitt *et al.* (10) observed a slow accumulation of F26BP in attached spinach leaves during the course of the photoperiod that was correlated with accumulation of sucrose and increased starch formation. Floating leaf discs on solutions containing sucrose or glucose resulted in increased levels of F26BP (10), which suggested that the F26BP level may be regulated by the soluble carbohydrate pool. The results obtained in the present study show that F26BP increases in intact leaves in response to a decrease in export, which is consistent with a regulatory role for F26BP in carbon metabolism.

#### LITERATURE CITED

1. CIHA AJ, WA BRUN 1978 Effect of pod removal on nonstructural carbohydrate concentration in soybean tissue. *Crop Sci* 18: 773-776
2. CSÉKE C, NF WEEDEN, BB BUCHANAN, K UYEDA 1982 A special fructose bisphosphate functions as a cytoplasmic regulatory metabolite in green leaves. *Proc Natl Acad Sci USA* 79: 4322-4326
3. FOYER C, J ROWELL, D WALKER 1983 The effect of sucrose on the rate of *de novo* sucrose biosynthesis in leaf protoplasts from spinach wheat and barley. *Arch Biochem Biophys* 220: 232-238
4. FURUYA E, K UYEDA 1980 An activation factor of liver phosphofructokinase. *Proc Natl Acad Sci USA* 77: 5861-5864
5. HEROLD A 1980 Regulation of photosynthesis by sink activity—the missing link. *New Phytol* 86: 131-144
6. HERS HG, E VAN SCHAFTINGEN 1982 Fructose 2,6-bisphosphate 2 years after its discovery. *Biochem J* 206: 1-12
7. HUBER SC 1981 Interspecific variation in activity and regulation of leaf sucrose phosphate synthetase. *Z Pflanzenphysiol* 102: 443-450
8. RUFTY JR TR, SC HUBER 1983 Changes in starch formation and activities of sucrose phosphate synthase and cytoplasmic fructose 1,6-bisphosphatase in response to source-sink alterations. *Plant Physiol* 72: 474-480
9. SETTER TL, WA BRUN, ML BRENNER 1980 Stomatal closure and photosynthetic inhibition in soybean leaves induced by petiole girdling and pod removal. *Plant Physiol* 65: 884-887
10. STITT M, R GERHARDT, B KÜRZEL, HW HELDT 1983 A Role for fructose 2,6-bisphosphate in the regulation of sucrose synthesis in spinach leaves. *Plant Physiol* 72: 1139-1141
11. STITT M, G MIESKES, HD SOLING, HW HELDT 1982 On a possible role of fructose 2,6-bisphosphate in regulating photosynthetic metabolism in leaves. *FEBS Lett* 145: 217-222
12. STITT M, W WIRTZ, HW HELDT 1980 Metabolite levels in the chloroplast and extrachloroplast compartments of spinach protoplasts. *Biochim Biophys Acta* 593: 85-102