

# Isotope Ratios of Cellulose from Plants Having Different Photosynthetic Pathways

Received for publication August 16, 1983 and in revised form October 31, 1983

LEONEL O. STERNBERG, MICHAEL J. DENIRO, AND HYRUM B. JOHNSON\*

*Department of Earth and Space Sciences (L. O. S.), and Department of Earth and Space Sciences and Archaeology Program (M. J. D.), University of California, Los Angeles, California 90024; and United States Department of Agriculture, Agricultural Research Service (H. B. J.), Grassland, Soil and Water, Research Laboratory, Temple, Texas 76503*

## ABSTRACT

Hydrogen and carbon isotope ratios of cellulose nitrate and oxygen isotope ratios of cellulose from C<sub>3</sub>, C<sub>4</sub>, and Crassulacean acid metabolism (CAM) plants were determined for plants growing within a small area in Val Verde County, Texas. Plants having CAM had distinctly higher deuterium/hydrogen (D/H) ratios than plants having C<sub>3</sub> and C<sub>4</sub> metabolism. When hydrogen isotope ratios are plotted against carbon isotope ratios, each photosynthetic mode separates into a distinct cluster of points. C<sub>4</sub> plants had many D/H ratios similar to those of C<sub>3</sub> plants, so that hydrogen isotope ratios cannot be used to distinguish between these two photosynthetic modes. *Portulaca mundula*, which may have a modified photosynthetic mode between C<sub>4</sub> and CAM, had a hydrogen isotope ratio between those of the C<sub>4</sub> and CAM plants. When oxygen isotope ratios are plotted against carbon isotope ratios, no distinct clustering of the C<sub>4</sub> and CAM plants occurs. Thus, oxygen isotope ratios are not useful in distinguishing between these metabolic modes. A plot of hydrogen isotope ratios versus oxygen isotope ratios for this sample set shows considerable overlap between oxygen isotope ratios of the different photosynthetic modes without a concomitant overlap in the hydrogen isotope ratios of CAM and the other two photosynthetic modes. This observation is consistent with the hypothesis that higher D/H ratios in CAM plants relative to C<sub>3</sub> and C<sub>4</sub> plants are due to isotopic fractionations occurring during biochemical reactions.

A large body of research has demonstrated that stable carbon isotope ratios of plants are related to photosynthetic pathways (1, 2, 19). C<sub>3</sub> plants, which fix CO<sub>2</sub> via ribulose biphosphate carboxylase (1), have  $\delta^{13}\text{C}$  values (see "Materials and Methods" section for definition of  $\delta$ ) that average about  $-25\text{‰}$ , while C<sub>4</sub> and CAM plants, which fix CO<sub>2</sub> via P-enolpyruvate carboxylase (1), have average  $\delta^{13}\text{C}$  values of about  $-13\text{‰}$  (1). Thus, carbon isotope ratios can be used to distinguish C<sub>3</sub> plants from CAM and C<sub>4</sub> plants. Differences of  $\delta^{13}\text{C}$  values among species within a given photosynthetic group are caused at least in part by gas exchange efficiency and factors associated with the leaf surface (10).

It has been demonstrated that hydrogen isotope ratios can be used to distinguish CAM plants from C<sub>3</sub> and C<sub>4</sub> plants (20, 21, 25). In one study, cellulose nitrate prepared from CAM plants collected in Riverside County, California, had average  $\delta\text{D}$  values of  $+56 \pm 30\text{‰}$  ( $n = 11$ ) while cellulose nitrate of C<sub>3</sub> and C<sub>4</sub> plants growing at the same site had average  $\delta\text{D}$  values of  $-69 \pm 35$  ( $n = 9$ ) and  $-27 \pm 4\text{‰}$  ( $n = 2$ ), respectively (20). In a study of greenhouse-grown plants, the average  $\delta\text{D}$  value of cellulose

nitrate of CAM plants was  $+30 \pm 19\text{‰}$  ( $n = 20$ ), while C<sub>3</sub> and C<sub>4</sub> plants had average  $\delta\text{D}$  values of  $-80 \pm 35$  ( $n = 10$ ) and  $-25 \pm 2\text{‰}$  ( $n = 2$ ), respectively (21). In both cases,  $\delta\text{D}$  values of CAM plants did not overlap with those of C<sub>3</sub> and C<sub>4</sub> plants. Oxygen isotope ratios, however, could not be used to distinguish CAM from C<sub>3</sub> and C<sub>4</sub> plants in either of these studies.

Previous reports on differences of hydrogen isotope ratios between C<sub>3</sub> and C<sub>4</sub> plants are conflicting (18, 25). Ziegler *et al.* (25) reported large differences in hydrogen isotope ratios between C<sub>3</sub> and C<sub>4</sub> plants, with C<sub>4</sub> plants being enriched in deuterium relative to C<sub>3</sub> plants. Smith and Epstein (18), however, observed no differences between hydrogen isotope ratios of C<sub>3</sub> and C<sub>4</sub> plants. One of the problems with both of these earlier reports was that isotope analysis was done on total plant matter. Different chemical components of plant matter can have different isotope ratios (16). Thus, differences in chemical composition between plants might contribute to differences in  $\delta\text{D}$  values and  $\delta^{18}\text{O}$  values. Furthermore, a large proportion of hydrogen in organic matter is exchangeable with water (9), and thus, some routine treatments of samples, such as rinsing with water, can alter their hydrogen isotope ratios. Isotopic analysis of plant cellulose and cellulose nitrate eliminates these variables because measurements are made on only one chemical component and exchangeable hydroxyl hydrogens are eliminated by nitration (9).

Another complicating factor in the study of oxygen and hydrogen isotopes in plants is that oxygen and hydrogen isotope ratios of plants are not influenced only by plant physiological behavior but also by climate and the  $^{18}\text{O}/^{16}\text{O}$  and D/H ratios of meteoric water. Variations in the isotopic composition of meteoric water ranging from about  $-40$  to  $+6\text{‰}$  for  $\delta^{18}\text{O}$  values and from about  $-320\text{‰}$  to  $+50\text{‰}$  for  $\delta\text{D}$  values have been reported (5). In general, meteoric water from warmer and drier climates has higher  $\delta^{18}\text{O}$  and  $\delta\text{D}$  values than meteoric water from cooler and more humid regions (12). Plant oxygen and hydrogen isotope ratios are also influenced by the effects of climate on transpiration. Leaf water from plants growing in drier climates has higher  $\delta\text{D}$  and  $\delta^{18}\text{O}$  values relative to local meteoric water than leaf water from plants growing in more humid climates (11, 23). To eliminate the effect of variability of isotope ratios of meteoric water and the effect of different climates on the plant oxygen and hydrogen isotope ratios, plants must be sampled within a restricted area, thus making it more likely that meteoric water and climate are the same for all plants being compared. Only then can differences in D/H and  $^{18}\text{O}/^{16}\text{O}$  ratios among groups of plants representing the different photosynthetic modes be ascribed to distinctive physiological and biochemical processes.

Recent analysis of cellulose nitrate and cellulose from C<sub>3</sub> and C<sub>4</sub> plants growing at one site in Riverside County, California and in a greenhouse showed that, in each case, hydrogen and oxygen

isotope ratios of C<sub>4</sub> plants were slightly higher than that of C<sub>3</sub> plants (20, 21). However, in these reports only two C<sub>4</sub> species were represented in each of the two sample sets, and thus, no firm conclusions could be drawn. In this study we report results of isotopic analysis of cellulose nitrate and cellulose from a sample set of 35 species composed of 13 C<sub>4</sub>, 12 C<sub>3</sub>, and 10 CAM species, all growing in close proximity in a well defined small drainage system.

## MATERIALS AND METHODS

Plants were collected near the Highway 90 bridge over the Pecos River in Val Verde County, Texas. The plants sampled were growing close to one another to maximize the probability that they had access to meteoric water with the same isotopic signature. Stem and leaf material were collected from three separate individual plants or plant populations (in the case of small plants) for each of the species selected. The triplicate samples provided opportunity for replicating isotope measurement of individual species. After the plants were identified, they were dried at 50°C for several days.

Dried plant material was ground to fine powder in a Wiley mill. Cellulose was extracted by the method of Wise (24). Oxygen isotope ratios of cellulose were determined by the method of Rittenberg and Ponticorvo (17) as modified by Burk (4). Carbon and hydrogen isotope ratios of cellulose nitrate prepared as described elsewhere (6) were determined by a modified version of the Stump and Frazer method (16, 22). Isotope ratios are expressed as  $\delta$  values:

$$\delta(\text{‰}) = \left[ \frac{R_{\text{sample}}}{R_{\text{standard}}} - 1 \right] \times 1000$$

where  $R$  represents <sup>18</sup>O/<sup>16</sup>O for oxygen, D/H for hydrogen, and <sup>13</sup>C/<sup>12</sup>C for carbon. The standards were SMOW<sup>1</sup> for oxygen and hydrogen and the PDB for carbon. The precisions of the isotopic analyses were  $\pm 2\text{‰}$  for  $\delta\text{D}$ ,  $\pm 0.5\text{‰}$  for  $\delta^{18}\text{O}$  and  $\pm 0.2\text{‰}$  for  $\delta^{13}\text{C}$ . A limited number of replicate measurements were made for selected species from each of the photosynthetic types.

## RESULTS

Carbon isotope ratios of plants sampled in this study are similar to those expected based on previously reported measurements for plants representing different photosynthetic modes (1, 2, 19), with C<sub>3</sub> plants having  $\delta^{13}\text{C}$  values ranging from  $-22.2$  to  $-26.1\text{‰}$  and CAM and C<sub>4</sub> plants having  $\delta^{13}\text{C}$  values between  $-10.6$  to  $-13.8\text{‰}$  (Table I). Thus, all CAM plants sampled in this study were operating in the CAM mode. The relative distribution of hydrogen isotope ratios among the plants representing the different photosynthetic types is also similar to previous reports (20, 21). Values of  $\delta\text{D}$  ranged from  $-71$  to  $-9\text{‰}$  for C<sub>3</sub> plants,  $-53$  to  $-20\text{‰}$  for C<sub>4</sub> plants, and  $+34$  to  $+66\text{‰}$  for CAM (Table I).  $\delta\text{D}$  values for CAMs ( $51 \pm 10\text{‰}$ ,  $n = 12$ ) are discrete from those of C<sub>3</sub> and C<sub>4</sub> plants, while those for C<sub>3</sub> ( $-42 \pm 16\text{‰}$ ,  $n = 16$ ) and C<sub>4</sub> ( $-33 \pm 12\text{‰}$ ,  $n = 16$ ) plants show considerable overlap (Table II and Fig. 1). Thus, it is unlikely that hydrogen isotope ratios will be useful to distinguish between these two photosynthetic pathways. The oxygen isotope ratios of the plants sampled showed more distinct patterns with less variability than previously reported for plants representing the different photosynthetic pathways (20, 21). Values of  $\delta^{18}\text{O}$  ranged from  $+23.5$  to  $+29.2\text{‰}$  for C<sub>3</sub> plants,  $+32.5$  to  $+37.0\text{‰}$  for C<sub>4</sub> plants, and  $+22.6$  to  $+35.4\text{‰}$  for CAMs (Table I). The average values were highest for C<sub>4</sub> plants ( $+34.7 \pm 1.4\text{‰}$ ,  $n = 16$ ), followed by CAM

<sup>1</sup> Abbreviations: SMOW, standard mean ocean water; PDB, belemnite from the Peedee formation of South Carolina.

Table I. Species, Photosynthetic Mode, Carbon and Hydrogen Isotope Ratios of Cellulose Nitrate and Oxygen Isotope Ratios of Cellulose for each Species Sampled

In some cases, more than one specimen was analyzed. CAM plants were identified by their succulence, C<sub>4</sub> plants by Kranz anatomy, and C<sub>3</sub> plants by their lack of succulence and Kranz anatomy.

Species	Mode	$\delta^{13}\text{C}$	$\delta\text{D}$	$\delta^{18}\text{O}$
<i>Acacia belandieri</i>	C <sub>3</sub>	-25.0	-49	+23.5
<i>Acacia rigidula</i>	C <sub>3</sub>	-24.2	-56	+23.5
<i>Cassia lindheimeriana</i>	C <sub>3</sub>	-25.7	-45	+28.3
<i>Fouquieria splendens</i>	C <sub>3</sub>	-26.1	-47	+27.0
<i>Fraxinus greggii</i>	C <sub>3</sub>	-23.3	-23	+26.6
		-22.9	-17	+25.0
		-23.9	-9	+26.2
<i>Jatropha dioica</i>	C <sub>3</sub>	-24.2	-52	+24.9
<i>Lippia graveolens</i>	C <sub>3</sub>	-26.1	-71	+28.8
<i>Penstemon bacharifolius</i>	C <sub>3</sub>	-24.0	-42	+27.6
		-25.6	-65	+28.2
		-25.8	-50	+29.2
<i>Prosopis glandulosa</i>	C <sub>3</sub>	-23.6	-44	+24.5
<i>Selagenella lepidophylla</i>	C <sub>3</sub>	-25.9	-31	+28.9
<i>Selagenella underwoodii</i>	C <sub>3</sub>	-25.7	-40	+27.8
<i>Dasyliirion texanum</i>	C <sub>3</sub>	-22.2	-28	+28.3
<i>Aristida wrightii</i>	C <sub>4</sub>	-12.6	-53	+32.5
		-11.8	-45	+32.6
<i>Bothriochloa sacchariodes</i>	C <sub>4</sub>	-11.4	-36	+35.1
<i>Bouteloua curtipendula</i>	C <sub>4</sub>	-12.1	-34	+32.5
<i>Bouteloua hirsuta</i>	C <sub>4</sub>	-12.4	-42	+37.0
		-12.5	-29	+35.2
<i>Digitaria californica</i>	C <sub>4</sub>	-12.1	-38	+34.5
<i>Erioneuron pilosum</i>	C <sub>4</sub>	-12.7	-25	+33.9
<i>Heteropogon contortus</i>	C <sub>4</sub>	-11.6	-43	+33.8
<i>Leptoloma cognatum</i>	C <sub>4</sub>	-11.0	-21	+36.7
<i>Panicum hallii</i>	C <sub>4</sub>	-12.1	-40	+35.8
<i>Pappophorum bicolor</i>	C <sub>4</sub>	-12.1	-42	+34.9
<i>Portulaca mundula</i>	C <sub>4</sub>	-11.9	-1	+33.9
<i>Sporobolus cryptandrus</i>	C <sub>4</sub>	-12.4	-33	+34.8
<i>Setaria viridis</i>	C <sub>4</sub>	-11.4	-20	+36.7
		-11.5	-34	+34.4
<i>Agave lecheguilla</i>	CAM	-11.0	+51	+22.6
<i>Echinocereus enneacanthus</i>	CAM	-10.9	+60	+31.1
		-12.5	+44	+35.4
<i>Echinocereus triglochidiatus</i>	CAM	-10.6	+34	+29.2
<i>Ferocactus hamataeanthus</i>	CAM	-12.8	+52	+29.7
<i>Opuntia edwardsii</i>	CAM	-13.8	+36	+28.7
<i>Opuntia leptocaulis</i>	CAM	-12.2	+57	+32.1
<i>Opuntia lindheimeri</i>	CAM	-11.5	+54	+27.9
<i>Opuntia phaeocantha</i>	CAM	-12.3	+39	+27.9
<i>Yucca torreyi</i>	CAM	-11.6	+63	+30.5
		-12.0	+66	+31.8
<i>Yucca baccata</i>	CAM	-13.5	+57	+29.0

Table II. Average  $\delta$  Values of each Photosynthetic Mode Sampled in this Study

Photosynthetic Mode	Average Isotope Ratios		
	$\delta^{13}\text{C}$	$\delta\text{D}$	$\delta^{18}\text{O}$
C <sub>3</sub>	$-24.6 \pm 1.3\text{‰}$	$-42 \pm 16\text{‰}$	$+26.8 \pm 1.9\text{‰}$
C <sub>4</sub>	$-12.0 \pm 0.5\text{‰}$	$-33 \pm 12\text{‰}$	$+34.6 \pm 1.4\text{‰}$
CAM	$-12.1 \pm 1.0\text{‰}$	$+51 \pm 10\text{‰}$	$+29.7 \pm 3.1\text{‰}$

( $+29.7 \pm 3.1\text{‰}$ ,  $n = 12$ ) and C<sub>3</sub> plants ( $+26.8 \pm 1.9\text{‰}$ ,  $n = 16$ ) (Table II). The  $\delta^{18}\text{O}$  values of the C<sub>4</sub> plants were distinctly separate from those of the C<sub>3</sub> plants, whereas the  $\delta^{18}\text{O}$  values of C<sub>4</sub> plants and CAM plants showed some overlap (Fig. 2).




FIG. 1. Relationship between  $\delta D$  and  $\delta^{13}C$  values of  $C_3$ ,  $C_4$ , and CAM plants. Points with bars indicate the mean and range of multiple measurements for different specimens of selected species.




FIG. 2. Relationship between  $\delta^{18}O$  and  $\delta^{13}C$  for the three photosynthetic modes. Points with bars indicate the mean and range of multiple measurements for different specimens of selected species.

### DISCUSSION

Clustering of each photosynthetic type relative to their  $\delta D$  and  $\delta^{13}C$  values (Fig. 1) is similar to but more pronounced than that reported previously (20, 21), with CAM plants in the upper right quadrant,  $C_4$  in the lower right quadrant, and  $C_3$  plants in the

lower left quadrant. The  $C_4$  group has one outlying point with an exceptionally high  $\delta D$  value of  $-1\text{‰}$  (Fig. 2). This value was obtained for cellulose nitrate of *Portulaca mundula* (Table I), which is a succulent  $C_4$  plant. Other species of this genus, namely *P. olearaceae* and *P. grandiflora*, have characteristics of CAM plants, such as acid flux and succulence, as well as characteristics of  $C_4$  plants, such as Kranz anatomy (13, 14). Because *P. mundula* is also a succulent  $C_4$  species with Kranz anatomy, it quite possibly has the same modified photosynthetic mode as *P. olearaceae* and *P. grandiflora*. This may have caused its  $\delta D$  value to be intermediate between those of the  $C_4$  and CAM groups. If this is the case, hydrogen isotope ratios may be sensitive to shifts from  $C_4$  to CAM. This type of shift would not affect the carbon isotope ratio, since the primary carboxylation reaction is mediated by P-enolpyruvate carboxylase for both photosynthetic pathways.

Oxygen isotope ratios of cellulose relative to the  $\delta^{13}C$  values of cellulose nitrate (Fig. 2) do not separate the photosynthetic types into separate clusters as completely as did the hydrogen and carbon isotope ratios (Fig. 1). Thus, oxygen isotope analysis cannot be used to distinguish CAM plants from  $C_4$  plants. One exceptional point (lower right quadrant) in Figure 2 deserves further discussion. It represents *Agave lecheguila*, a CAM species that had an unusually low  $\delta^{18}O$  value relative to the other CAM plants. We had observed previously that another *Agave* species, *A. deserti*, had low  $\delta^{18}O$  values relative to other CAM species growing in the same area (20). Although these two *Agave* species are leaf succulents, their low  $\delta^{18}O$  values probably are not due solely to their physiognomy, as suggested earlier (20), since two other leaf succulent species analyzed in the present study, *Yucca bacatta* and *Y. torreyi* (Table I) did not have low  $\delta^{18}O$  values. Some other hypothesis must be advanced to explain the low  $\delta^{18}O$  values of these *Agave* species.

Figure 3 shows the relationship between the  $\delta D$  values of cellulose nitrate and the  $\delta^{18}O$  values of cellulose extracted from plants sampled in this study. The data points are separated into




FIG. 3. Relationship between  $\delta D$  and  $\delta^{18}O$  for the three photosynthetic modes. Points with bars indicate the mean and range of multiple measurements for different specimens of selected species.

three clusters representing the C<sub>3</sub>, C<sub>4</sub>, and CAM photosynthetic types, as was observed in the  $\delta D$  versus  $\delta^{13}C$  plot (Fig. 1). Three factors influence the relationship between the  $\delta D$  value of cellulose nitrate and the  $\delta^{18}O$  value of cellulose in a plant. The first is the relationship between the  $\delta D$  and  $\delta^{18}O$  values of the ground-water absorbed by the plant through its roots, which is given by the equation  $\delta D = 8 \times \delta^{18}O + 10\%$  (the meteoric water line equation) (5). The second factor relates to the isotopic fractionation that occurs during evapotranspiration, in which the leaf water becomes enriched in deuterium and in  $^{18}O$  relative to groundwater (15). The relationship between the  $\delta D$  and  $\delta^{18}O$  values of leaf water from plants undergoing evapotranspiration is approximately linear with a slope of about 3 (15). The third factor which influences the relationship between the  $\delta D$  value of cellulose nitrate and  $\delta^{18}O$  value of cellulose are the biochemical isotopic fractionations associated with carbohydrate and cellulose synthesis (8). If the elevated hydrogen isotope ratios in the CAM group are due to evapotranspiration, then the oxygen isotope ratios of cellulose from CAM plants should also be distinctly elevated. This is because evapotranspiration, which causes enrichment of deuterium, also enriches plant water in  $^{18}O$  (11), with the  $^{18}O$ -enrichment subsequently being passed on to cellulose (7). Joint enrichment of D and  $^{18}O$  in CAM cellulose nitrate and cellulose was not observed in this study. The  $\delta D$  values of the CAM group are elevated relative to the rest of the samples measured without a concomitant increase in  $\delta^{18}O$  values of cellulose (Fig. 3). Therefore we conclude, as we did previously (20), that  $\delta D$  values in CAM plants are elevated due to biochemical reactions.

The C<sub>4</sub> group shows elevated  $\delta D$  and  $\delta^{18}O$  values relative to the C<sub>3</sub> plants. One explanation might be that C<sub>4</sub> plants produce their cellulose at a different time of the year and thus use meteoric water with a different isotopic composition than that used by C<sub>3</sub> plants. This explanation seems unlikely because an increase in the  $\delta^{18}O$  values of meteoric water will be accompanied by an increase in the  $\delta D$  values by a factor of 8 (5). In our sample set, however, the hydrogen isotope ratios of cellulose nitrate of C<sub>4</sub> plants are only slightly higher than those of the C<sub>3</sub> group, while the  $\delta^{18}O$  values of cellulose from the two groups differ by large amounts. Furthermore, all plants sampled showed evidence of being essentially opportunistic in exhibiting new growth in response to late summer rains just prior to collection. A second explanation is that differences in  $\delta D$  and  $\delta^{18}O$  values between C<sub>3</sub> and C<sub>4</sub> plants are related to evapotranspiration. A regression analysis between the  $\delta D$  values of cellulose nitrate and  $\delta^{18}O$  values of cellulose from C<sub>3</sub> and C<sub>4</sub> plants gives a line with a slope of 1.4 (intercept = -80,  $r = 0.43$ ) which approaches the slope of the  $\delta D$  versus  $\delta^{18}O$  line of leaf water from plants which are undergoing evapotranspiration (15). Regression characteristics from previously published results (20) were similar (slope = 1.5, intercept = -94.5,  $r = 0.50$ ). Thus, it seems likely that differences in oxygen and hydrogen isotopes in C<sub>4</sub> plants relative to C<sub>3</sub> plants are related to evapotranspiration and not biochemical reactions or differences in the water available for plant growth. Recent comparative field studies demonstrated that C<sub>4</sub> plants are less sensitive to a decrease in RH when compared to C<sub>3</sub> plants (3). C<sub>4</sub> plants did not show a decrease in photosynthesis with a reduction in RH whereas C<sub>3</sub> plants did. Thus, C<sub>4</sub> plants may have a considerable amount of cellulose synthesized under drier conditions when plant water would be more enriched in deuterium and oxygen relative to meteoric water; conversely, C<sub>3</sub> plants may not synthesize much cellulose unless conditions are more humid.

The differences in  $\delta^{18}O$  and  $\delta D$  values observed among species and/or individual plants within a given photosynthetic type may reflect processes associated with gaseous exchange, water use efficiency, and tolerance to moisture stress that are more subtle

than those that separate the major photosynthetic pathways (10, 11).

## CONCLUSIONS

Our measurements agree with previously published results in that a clear discrimination existed among C<sub>3</sub>, C<sub>4</sub>, and CAM photosynthetic pathways when hydrogen and carbon isotope ratios of cellulose nitrate were considered. These measurements might also be sensitive to plants which have modified photosynthetic modes between C<sub>4</sub> and CAM, as may be the case for *P. mundula*. A large overlap in oxygen isotope ratios existed between CAM and the other two photosynthetic pathways without an overlap in hydrogen isotope ratios of CAM plants relative to C<sub>3</sub> and C<sub>4</sub> plants. This indicates that deuterium enrichment in CAM plants as compared to C<sub>3</sub> and C<sub>4</sub> plants is due to isotopic fractionations occurring during biochemical reactions and not during evapotranspiration, as has been suggested (25). Cellulose from C<sub>4</sub> plants had distinctly different oxygen isotope ratios compared to cellulose from C<sub>3</sub> plants. Our data are consistent with the hypothesis that as conditions become drier, C<sub>4</sub> plants continue cellulose synthesis while cellulose synthesis for C<sub>3</sub> plants is inhibited. Thus, cellulose in C<sub>4</sub> plants would be synthesized under drier conditions when plant water would be enriched in deuterium and  $^{18}O$ . This hypothesis can be tested by growing C<sub>4</sub> plants and C<sub>3</sub> plants under controlled arid and humid conditions and then examining the relationships between their  $\delta D$  and  $\delta^{18}O$  values.

## LITERATURE CITED

- BENDER MM 1971 Variation in  $^{13}C/^{12}C$  ratios of plants in relation to the pathway of photosynthetic carbon dioxide fixation. *Phytochemistry* 10: 1239-1344
- BENDER MM, I RONHAM, HM VINES, CC BLACK 1973  $^{13}C/^{12}C$  ratio changes in Crassulacean acid metabolism plants. *Plant Physiol* 52: 427-430
- BUNCE JA 1983 Differential sensitivity to humidity of daily photosynthesis in the field in C<sub>3</sub> and C<sub>4</sub> species. *Oecologia* 57: 262-265
- BURK RL 1980 Factors affecting  $^{18}O/^{16}O$  ratios in cellulose. Ph.D. dissertation. University of Washington, Seattle
- CRAIG H 1961 Isotopic variations in meteoric water. *Science* 133: 1702-1703
- DENIRO MJ 1978 The effects of different methods of preparing cellulose nitrate on the determination of D/H ratios of nonexchangeable hydrogen of cellulose. *Earth Planet Sci Lett* 54: 177-185
- DENIRO MJ, S EPSTEIN 1979 Relationship between oxygen isotope ratios of terrestrial plant cellulose, carbon dioxide and water. *Science* 204: 51-53
- DENIRO MJ, S EPSTEIN 1981 Isotopic composition of cellulose from aquatic organisms. *Geochim Cosmochim Acta* 42: 495-506
- EPSTEIN S, CJ YAPP, JH HALL 1976 The determination of D/H ratio of non-exchangeable hydrogen in cellulose extracted from aquatic and land plants. *Earth Planet Sci Lett* 30: 241-251
- FARQUHAR GD, MH O'LEARY, JA BERRY 1982 On the relationship between carbon isotope discrimination and intercellular carbon dioxide concentration in leaves. *Aust J Plant Physiol* 9: 121-137
- FERHI A, R LETOLLE 1977 Transpiration and evaporation as the principal factors in oxygen isotope variations of organic matter in land plants. *Physiol Veg* 15: 363-370
- FRIEDMAN I, AC REDFIELD, B SCHOEN, J HARRIS 1964 The variation of the deuterium content of natural waters in the hydrological cycle. *Rev Geophys* 2: 177-224
- KOCH K, RA KENNEDY 1980 Characteristics of Crassulacean acid metabolism in the succulent C<sub>4</sub> dicot *Portulaca olearacea* L. *Plant Physiol* 65: 193-197
- KU SB, YJ SHIEH, BJ REGER, CC BLACK 1981 Photosynthetic characteristics of *Portulaca grandiflora*, a succulent C<sub>4</sub> dicot. *Plant Physiol* 68: 1073-1080
- LESAINTE C, L MERLIVAT, J BRICOUT, J-C FONTES, R GAUTHERET 1974 Sur la composition en isotopes stables de l'eau de la tomate et du maïs. *C R Acad Sci Paris* 278D: 2925-2930
- NORTHFELT DW, MJ DENIRO, S EPSTEIN 1981 Hydrogen and carbon isotope ratios of the cellulose nitrate and saponifiable lipid fractions prepared from annual growth rings of a California redwood. *Geochim Cosmochim Acta* 45: 1895-1898
- RITTENBERG D, L PONTICORVO 1956 A method of determination of the  $^{18}O$  concentrations of the oxygen of organic compounds. *J Appl Radiat Isotopes* 1: 208-214
- SMITH BN, S EPSTEIN 1970 Biogeochemistry of the stable isotopes of hydrogen and carbon in salt marsh. *Plant Physiol* 46: 738-742
- SMITH BN, S EPSTEIN 1970 Two categories of  $^{13}C/^{12}C$  ratios for higher plants.

- Plant Physiol 47: 380-384
20. STERNBERG L, MJ DENIRO 1983 Isotopic composition of cellulose from C<sub>3</sub>, C<sub>4</sub> and CAM plants growing in the vicinity of one another. *Science* 220: 947-948
  21. STERNBERG L, MJ DENIRO, IP TING 1983 Carbon, hydrogen and oxygen isotope ratios of plants having intermediary photosynthetic modes. *Plant Physiol* 74: 104-107
  22. STUMP RK, JW FRAZER 1971 Simultaneous determination of carbon, hydrogen, and nitrogen in organic compounds. *Nucl Sci Abstr* 28: 746
  23. WERSHAW RL, I FRIEDMAN, SJ HELLER 1966 Hydrogen isotope fractionation of water passing through trees. In GD Hobson, GC Spears, eds, *Advances in Organic Geochemistry*. Pergamon Press, New York, pp 55-67
  24. WISE LE 1944 *Wood Chemistry*. Reinhold, New York, 1-21
  25. ZIEGLER H, CB OSMOND, W STICKLER, D TRIMBORN 1976 Hydrogen isotope discrimination in higher plants: correlation with photosynthetic pathway and environment. *Planta* 128: 85-92