

Communication

Lethal Hydroxyl Radical Production in Paraquat-Treated Plants¹

Charles F. Babbs*, Jo Ann Pham, and Ronald C. Coolbaugh

Biomedical Engineering Center (C.F.B., J.A.P.) and Department of Botany and Plant Pathology (R.C.C.), Purdue University, West Lafayette, Indiana 47907

ABSTRACT


Bipyridinium herbicides, including paraquat and diquat, are believed to act by generating highly reactive, oxygen-centered free radicals within chloroplasts when treated plants are exposed to sunlight. This hypothesis has not yet been confirmed by direct chemical measurements of specific free radicals. We studied paraquat-treated plants using a new method able to detect and quantify formation of highly reactive and deleterious hydroxyl radicals (HO*), in which dimethyl sulfoxide (DMSO) is used as a molecular probe. DMSO is oxidized by HO* to form the stable, nonradical compound, methane sulfinic acid, which can be easily extracted from plant tissue and measured spectrophotometrically. Initial experiments revealed formation of extraordinary numbers of hydroxyl radicals in light-exposed, paraquat + DMSO-treated plants, equivalent at least to the cumulative number of HO* radicals per gram of fresh tissue that would be produced by 10,000 rads of gamma irradiation. This appears to be the greatest production of hydroxyl radicals yet observed in a biological system and is quite sufficient to explain the rapid death of top growth in paraquat-treated plants.

Bipyridinium herbicides are widely used in agriculture because they are inexpensive and highly effective against a wide variety of weeds. Since these herbicides are quickly rendered inert by absorption to clays in soil and by bacterial action, weeds can be sprayed and useful crops planted with minimal tilling of the soil and minimal soil erosion. The currently accepted mechanism of action of bipyridinium herbicides, for which paraquat (1,1'-dimethyl-4,4'-bipyridinium dichloride) is the prototypic compound, involves light-catalyzed generation of oxygen containing free radicals within chloroplasts (11, 16). Free radicals are highly reactive chemical species having an unpaired valence electron, herein denoted as*. Of late, oxygen-centered free radicals, including the superoxide anion, O₂⁻, and the highly reactive hydroxyl radical, HO*, have been proposed as a new class of pathogens in biology and medicine (3, 8, 13, 15, 21, 24) that can be placed in the sequence—parasites, bacteria, viruses, auto-antibodies, free radicals—in order of decreasing physical size and roughly increasing dif-

ficulty of detection. Hydroxyl radicals, in particular, are among the most highly reactive molecular species known. They are generally considered to initiate a major fraction of radiation-induced damage to living tissues when high energy γ rays split the hydrogen-oxygen bond in intracellular water: HOH \rightarrow H* + *OH (25). In the presence of suitable chelated transitional metals, particularly iron, hydroxyl radicals can also be formed from superoxide and hydrogen peroxide at ambient temperatures and neutral pH by the iron-catalyzed, superoxide-driven Fenton reaction (Table IC) (2, 13, 14, 24).

According to the free radical hypothesis explaining the effect of paraquat upon green plants (12, 16, 27), the divalent paraquat cation, paraquat²⁺, accepts an electron from PSI in chloroplasts to produce the blue, monocationic paraquat radical, paraquat^{•+}, which in turn is rapidly oxidized by molecular oxygen to regenerate paraquat²⁺ with concomitant reduction of oxygen to superoxide, O₂⁻ (Table IA). The paraquat molecule is then ready for another cycle of reduction and oxidation. A portion of the excess superoxide, either spontaneously or in the presence of SOD,² is converted to hydrogen peroxide. Superoxide and hydrogen peroxide that escape detoxification (Table IB) by superoxide dismutases, catalase (in peroxisomes), and peroxidases may then participate in two pathological pathways leading to the generation of highly toxic HO* radicals. The first is the superoxide-driven Fenton reaction (Table IC), involving trace amounts of iron. The second is a Fenton-like reaction, proposed by Winterbourn (Table ID) (30), in which the reduced paraquat radical, paraquat^{•+}, reacts with hydrogen peroxide either directly (30) or in the presence of very small, catalytic concentrations of iron (C Winterbourn, personal communication, February 1989).

Having recently developed a new method to trap and quantify HO* radicals in living systems (4, 5), we conducted experiments to measure HO* generation in paraquat-treated leaves and thus confirm or deny free radical hypotheses regarding paraquat's effect. We used DMSO as a molecular probe for HO* and assayed MSA, the immediate nonradical product of the trapping reaction.



DMSO hydroxyl MSA methyl
MSA radical

¹ This work was supported by a Focused Giving Grant from Johnson & Johnson, and by grant HL-35996 from the National Heart, Lung, and Blood Institute, U.S. Public Health Service, Bethesda, MD.

² Abbreviations: SOD, superoxide dismutase; MSA, methane sulfinic acid.

Table I. Model of Paraquat-Induced Free Radical Generation

A. Paraquat-induced oxidative stress	
PSI	
paraquat ²⁺	→ paraquat ^{•+} ,
O ₂ + paraquat ^{•+}	→ O ₂ ^{-•} + paraquat ²⁺
B. Normal detoxification of superoxide	
2 H ⁺ + 2 O ₂ ^{-•}	→ H ₂ O ₂ + O ₂ (SOD reaction)
2 H ₂ O ₂	→ 2 H ₂ O + O ₂ (catalase reaction)
C. Superoxide-driven Fenton reaction	
O ₂ ^{-•} + Fe ³⁺	→ O ₂ + Fe ²⁺
2 O ₂ ^{-•} + 2 H ⁺	→ H ₂ O ₂ + O ₂
H ⁺ + Fe ²⁺ + H ₂ O ₂	→ HO [•] + Fe ³⁺ + H ₂ O
D. Winterbourn's reaction	
paraquat ^{•+} + H ₂ O ₂	→ paraquat ²⁺ + OH ⁻ + HO [•] (original version)
paraquat ^{•+} + Fe ³⁺	→ paraquat ²⁺ + Fe ²⁺ (trace Fe catalyzed)
H ⁺ + Fe ²⁺ + H ₂ O ₂	→ HO [•] + Fe ³⁺ + H ₂ O

of DMSO with hydroxyl radicals (10, 17, 18) as a marker for HO[•] generation.

Because DMSO is exceedingly nontoxic and can be tolerated by living systems in up to 1 M concentrations (1, 7, 19, 20, 28, 29), because it is rapidly absorbed and distributes to all tissue compartments (9), and because it yields a single, stable product (MSA) upon oxidation by HO[•], DMSO is a nearly ideal molecular probe for trapping HO[•] radicals. A further advantage in using DMSO to trap hydroxyl radicals is that one can estimate actual numbers of HO[•] radicals produced—as opposed to simply detecting the presence of HO[•] radicals without being able to specify how many were generated. Hydroxyl radicals are so reactive that they readily combine with normal cellular constituents at frequencies within one or two orders of magnitude of their collision frequency (23). Unless the molecular probe is present in high concentrations, all but a tiny percentage of HO[•] radicals will react with endogenous biomolecules rather than the introduced molecular probe. One therefore has no way of knowing what fraction of them was actually trapped, or in turn the total number of HO[•] radicals generated. Thus, to quantify the numbers of HO[•] radicals generated in living tissue, the concentration of the trapping agent must be relatively high, in practice about 0.1 M or greater for DMSO (6, 22), so that the probability that an HO[•] will react with the trapping agent, rather than a host biomolecule is large. In the present studies, we used 5% (0.7 M) DMSO as a molecular probe for HO[•] generation in paraquat-treated plants.

MATERIALS AND METHODS

Treatment of Plant Material

To estimate HO[•] generation *in vivo*, we measured MSA produced in the small aquatic plant, duckweed (*Lemna minor*), and in perennial ryegrass (*Lolium perenne* L.) treated with DMSO and paraquat, compared to controls treated with paraquat only, DMSO only, or neither agent. Flats of ryegrass were exposed to natural sunlight (about 2000 μE m⁻² s⁻¹) for 1 to 4 d during July to August 1988, until paraquat-treated leaves became brown. In similar experiments, cultures of duckweed were exposed to fluorescent light (830 μE m⁻² s⁻¹) for 24 h in an environmental control growth chamber at 20

to 22°C. In the studies of ryegrass, DMSO was applied as a 5% solution in water so that it thoroughly soaked the sod. Paraquat (0.1% of the commercial concentrate, Gramoxone, from ICI Americas) was applied once as a mist spray until dripping to the foliage prior to light exposure. In studies of duckweed the small aquatic plants were suspended in 5% DMSO and 0.1% paraquat was sprayed liberally over exposed surface leaves. After light exposure, 2 to 5 g of leaves were harvested for analysis of MSA content by a modification of a colorimetric assay previously described (4) after quick-freezing with liquid N₂, grinding with a mortar and pestle, aqueous extraction of the resulting powder with 15 to 20 mL distilled water, centrifugation to remove solid materials, and concentration of the aqueous supernatant by lyophilization. The sulfinic acid assay included two stages: removal of interference with the aid of Sep-Pak columns and the color reaction with fast blue BB dye.

Removal of Interference with Sep-Pak Columns

Principle

Many potentially interfering species are anionic detergents, such as free fatty acids, which form salt-like complexes with the fast blue BB dye used for the color reaction in the subsequent colorimetric assay (5). Most interfering anions are much less hydrophilic than MSA itself. To remove a portion of such interfering species, a preliminary extraction of the sample with toluene/*n*-butanol (3/1) at neutral pH may be used. Further removal of detergent-like interference can be obtained by adding fast blue BB dye to the sample and applying the mixture to a lipophilic (C₁₈ Sep-Pak) column. When the column is subsequently eluted with water, the complexes of dye and lipophilic anions remain on the column, while sulfinic acid is readily eluted and collected. In this way detergent-like anions that are specifically capable of complexing with fast blue BB dye to form lipophilic salts are sequestered on the column. The color reaction with additional fast blue BB dye and toluene/*n*-butanol extraction is then run on the effluent fraction containing the sulfinic acid.

Procedure

Lyophilized samples were resuspended in 1 mL of distilled water and extracted twice with 2 mL of toluene/*n*-butanol (3/1). The pH of the aqueous phase was adjusted to 2.5 by addition of HCl. Then 100 μL of 30 mM fast blue BB salt was added to the aqueous sample with mixing, which was applied to a Sep-Pak C₁₈ column (Cat. No. 51910, Waters Associates, Milford, MA 01757) that had been preeluted with 2 mL methanol and 2 mL water. The methane sulfinate anion was then eluted from the column with distilled water. The first 1.3 mL of effluent was discarded, and the next 1.5 mL collected for assay by the color reaction.

Color Reaction

The 1.5 mL aqueous volume collected from the column was transferred to a test tube, the pH was adjusted to 2.5 by the addition of HCl, and the color reaction was begun by

addition of 0.1 mL of 30 mM fast blue BB salt (freshly prepared and kept in the dark). Ten min were allowed for product development at room temperature in the dark. Then 1.5 mL of toluene/*n*-butanol (3/1) was added and mixed thoroughly with the aqueous phase for 60 s on a Vortex mixer to extract the yellow, hydrophobic reaction product. After low speed centrifugation to separate the phases, the lower phase, containing unreacted diazonium salt, was removed by aspiration and discarded. The toluene/*n*-butanol phase was washed with 2 mL of *n*-butanol-saturated water for 30 s to remove remaining unreacted diazonium salt. The tubes were centrifuged at 500*g* for 3 min, and the upper phase, containing the diazo-sulfones, was transferred to a cuvet. One-tenth mL of pyridine/glacial acetic acid (95/5) was added to stabilize the color, which otherwise fades gradually at acid pH. The absorbance as a function of the wavelength from 340 to 520 nm was recorded on a stripchart recorder, using a blank prepared beginning with 1.5 mL of distilled water carried through the same procedure. The sulfonic acid content was then calculated from the absorbance at $\lambda_{\text{max}} = 425$ nm, with reference to a standard curve.

RESULTS AND DISCUSSION

The results of a typical experiment using *Lemna minor* are presented in Figure 1 as visible absorbance spectra from the sulfonic acid assay of whole plants exposed to the various treatments. Standard, authentic MSA produces a characteristic absorbance peak at 420 nm (dotted curve). Plants treated with both paraquat and DMSO (D) produced a large, similarly shaped peak. The sharp peak for DMSO + paraquat-treated leaves was characteristic of all experiments. The smaller peak for paraquat-only-treated leaves (C) was common for duck-




Figure 1. Data from typical experiment with duckweed (*Lemna minor*), in which hydroxyl radicals were trapped and measured as methane sulfonic acid. Visible absorbance spectra of colorimetric assay of untreated leaves (A), DMSO-only-treated leaves (B), paraquat-only-treated leaves (C), and paraquat + DMSO-treated leaves (D). Dotted line represents 300 nmol MSA standard.

weed but not for ryegrass. Leaves treated with DMSO, without paraquat (B) or either agent (A) generated no significant signal at 420 nm.

Quantitative yields of MSA per gram (fresh weight) of leaves, shown in Table II, were on the order of 100 nmol/g or 0.1 mM for ryegrass and 475 nmol/g or about 0.5 mM for duckweed. Although this measured production of MSA would represent oxidation of only about 0.01 percent of the DMSO present in the leaves, the cumulative effect of 0.1 mM hydroxyl radicals would be highly toxic to any living system. A revealing comparison can be made by calculating the dose of γ irradiation required to produce the same number of hydroxyl radicals as were generated biologically in the paraquat-treated leaves. This can be done by applying the physical formula for the cumulative number of hydroxyl radicals formed per unit volume of γ irradiated water

$$\text{Number of HO}^* \text{ radicals} = \text{radiation dose} \times G_{\text{HO}^*}$$

where G_{HO^*} is the radiochemical yield of hydroxyl radicals, a known constant equal to 2.9 molecules per 100 electron volts of absorbed energy (25). After converting units (using $1 \text{ eV} = 1.602 \times 10^{-19} \text{ J}$ and $1 \text{ rad} = 0.01 \text{ J/kg}$ absorbed energy), one obtains the expression for the net yield, y , of hydroxyl radicals in water irradiated by dose, D , of high energy γ rays as y (pmol/g) = $3.01 D$ (rads). Thus, a yield of 100 nanomoles of hydroxyl radicals per gram = 100,000 pmol/g is equivalent to that caused by a radiation dose of approximately 33,000 rads. For comparison, a dose of γ irradiation sufficient to kill 97% of cells grown in tissue culture is on the order of 10,000 rads (26). Thus, the cumulative numbers of hydroxyl radicals that we have detected in plants treated with normal, herbicidal concentrations of paraquat are very great indeed and are likely to represent a lethal event. The results provide, we believe for the first time, a direct chemical test of the role of a specific oxygen-centered free radical (HO*) in paraquat-

Table II. Methane Sulfonic Acid Production in DMSO + Paraquat Treated Plants—Results of Preliminary Studies

Quantitative yields of HO* trapped as MSA per g (fresh weight) of leaves calculated from the absorbance at 425 nm. To correct for false positive, background signal, the mean absorbance per gram for all ryegrass control groups, which did not differ significantly, was subtracted from the absorbance of experimental (paraquat + DMSO) treated ryegrass. Similarly, the absorbance for duckweed controls concomitantly treated with paraquat only, which was the largest control value, was subtracted from that of experimental (paraquat + DMSO) duckweed. Experimental and control plants were grown side by side under identical conditions. Mean control absorbance was about 20% of that for DMSO + paraquat treated plants. The generation of MSA was then calculated from the experimental-control absorbance difference with reference to a standard curve. Results expressed as MSA per gram fresh weight.

Experiment Number	MSA in Ryegrass	MSA in Duckweed
	<i>nmol/g fresh leaves</i>	
1	67	983
2	27	109
3	124	394
4	149	414

induced leaf kill, as well as the first direct, chemical demonstration of lethal numbers of oxygen radicals in a biological system that has not been exposed to ionizing radiation.

LITERATURE CITED

- Ashwood-Smith MJ (1975) Current concepts concerning radioprotective and cryoprotective properties of dimethyl sulfoxide in cellular systems. *Ann NY Acad Sci* **243**: 246–256
- Aust SD, Morehouse LA, Thomas CE (1985) Hypothesis paper—role of metals in oxygen radical reactions. *J Free Rad Biol Med* **1**: 3–25
- Babbs CF (1985) Role of iron ions in the genesis of reperfusion injury following successful cardiopulmonary resuscitation: preliminary data and a biochemical hypothesis. *Ann Emerg Med* **14**: 777–783
- Babbs CF, Gale MJ (1987) Colorimetric assay for methanesulfinic acid in biological samples. *Anal Biochem* **163**: 67–73
- Babbs CF, Gale MJ (1988) Methane sulfinic acid production from DMSO as an indicator of hydroxyl radical production in vivo. In *Free Radicals: Methodology and Concepts*. C Rice-Evans, B Halliwell, eds, Richelieu Press, London, pp 91–121
- Babbs CF, Griffin DW (1988) Scatchard analysis of methane sulfinic acid production from dimethyl sulfoxide: a method to quantify hydroxyl radical formation in physiologic systems. *Free Rad Biol Med* **6**: 493–503
- Benville PE, Smith CE, Shanks WE (1968) Some toxic effects of dimethylsulfoxide in salmon and trout. *Toxicol Appl Pharmacol* **12**: 156–187
- Bulkley GB (1983) The role of oxygen free radicals in human disease processes. *Surgery* **94**: 407–411
- Denko CW, Goodman RM, Miller R, Donovan T (1967) Distribution of dimethyl sulfoxide in the rat. *Ann NY Acad Sci* **141**: 77–84
- Dixon WT, Norman ROC, Buley AL (1964) Electron spin resonance studies of oxidation. Part II: Aliphatic acids and substituted acids. *J Chem Soc* 3625–3634
- Duke S (1985) *Weed Physiology*. CRC Press, Boca Raton, FL, pp 131–233
- Farrington JA, Ebert M, Land EJ, Fletcher K (1973) Bipyridylum quaternary salts and related compounds. V. Pulse radiolysis studies of the reaction of paraquat radical with oxygen, implications for the mode of action of bipyridyl herbicides. *Biochim Biophys Acta* **314**: 372–381
- Freeman BA, Crapo JD (1982) Biology of disease—free radicals and tissue injury. *Lab Invest* **47**: 412–426
- Fridovich I (1983) Superoxide radical: an endogenous toxicant. *Annu Rev Pharmacol Toxicol* **23**: 239–257
- Halliwell B, Gutteridge JMC (1984) Oxygen toxicity, oxygen radicals, transition metals and disease. *Biochem J* **219**: 1–14
- Halliwell B, Gutteridge JMC (1987) *Free Radicals in Biology and Medicine*. Oxford University Press, Oxford, pp 206–234
- Klein SM, Cohen G, Cederbaum AI (1981) Production of formaldehyde during metabolism of dimethyl sulfoxide by hydroxyl radical generating systems. *Biochemistry* **20**: 6006–6012
- Lagercrantz C, Forshult S (1969) Trapping of short-lived free radicals as nitroxide radicals detectable by ESR spectroscopy. The radicals formed in the reaction between OH-radicals and some sulphoxides and sulphones. *Acta Chem Scand* **23**: 811–817
- Mason MM (1971) Toxicology of DMSO in animals. In SW Jacob, EE Rosen, DC Wood, eds, *Dimethyl Sulfoxide*. Marcel Dekker, New York, pp 113–131
- Maurer RR (1977) Ciba Foundation Symposium 52: The Freezing of Mammalian Embryos. Elsevier, Amsterdam, p 116
- McCord JM (1985) Oxygen-derived free radicals in postischemic tissue injury. *N Engl J Med* **312**: 159–163
- Repine JE, Pfenninger OW, Talmage DW, Berger EM, Pettijohn DE (1981) Dimethylsulfoxide prevents DNA nicking mediated by ionizing radiation or iron/hydrogen peroxide-generated hydroxyl radical. *Proc Natl Acad Sci USA* **78**: 1001–1003
- Semenov NN (1958) *Some Problems of Chemical Kinetics and Reactivity, Vol 1*. Pergamon Press, New York
- Slater TF (1972) *Free Radical Mechanisms in Tissue Injury*. Pion Limited, London
- Spinks JWT, Woods RJ (1976) *An Introduction to Radiation Chemistry, Ed 2*. Wiley-Interscience, New York
- Suit HD (ed) (1973) *Radiation Biology: A Basis for Radiotherapy*. Textbook of Radiotherapy, Lea & Febiger, Philadelphia
- Vaughan KC, Duke SO (1983) In situ localization of the sites of paraquat action. *Plant Cell Environ* **6**: 13–20
- Willadsen SM (1977) Factors affecting the survival of sheep embryos during deep-freezing and thawing. Ciba Foundation Symposium 52: The Freezing of Mammalian Embryos. Elsevier, Amsterdam, pp 175–201
- Wilmot I, Rowson LEA (1973) Experiments on the low temperature preservation of cow embryos. *Vet Rec* **92**: 686–690
- Winterbourn CC (1981) Production of hydroxyl radicals by paraquat radicals and H₂O₂. *FEBS Lett* **128**: 2339–2342