

Communication

Overproduction of Petunia Chloroplastic Copper/Zinc Superoxide Dismutase Does Not Confer Ozone Tolerance in Transgenic Tobacco¹

Lynne H. Pitcher, Eileen Brennan, Arthur Hurley, Pamela Dunsmuir, James M. Tepperman, and Barbara A. Zilinskas*

Department of Biochemistry and Microbiology, Lipman Hall, Cook College, Rutgers University, New Brunswick, New Jersey 08903–0231 (L.H.P., B.A.Z.), Department of Plant Pathology, Martin Hall, Cook College, Rutgers University, New Brunswick, New Jersey 08903–0231 (E.B., A.H.), and DNA Plant Technology Corporation, 6701 San Pablo Avenue, Oakland, California 94608 (P.D., J.M.T.)

ABSTRACT

Transgenic tobacco (*Nicotiana tabacum* cultivar W38) plants that overproduce petunia chloroplastic Cu/Zn superoxide dismutase were exposed to ozone dosages that injure control tobacco plants. Based on foliar injury ratings, there was no consistent protection provided to the transgenic plants. These data indicate that an increase in the chloroplastic Cu/Zn superoxide dismutase alone is not sufficient to reduce ozone toxicity.

bacco. Tepperman and Dunsmuir (19) transformed “Wisconsin 38” tobacco with DNA encoding petunia chloroplastic Cu/Zn-SOD, which was expressed in the chloroplast such that the transgenic plants overproduced the chloroplastic SOD. Inasmuch as certain histological (20) and physiological evidence (22) identifies the chloroplast as a primary site of O₃-induced injury, we proposed to use the transformed tobacco to test the hypothesis that tobacco plants with elevated levels of chloroplastic SOD are rendered tolerant to O₃ at dosages that normally injure tobacco.

Ozone damage is thought to result from the action of free radicals and other oxidants produced by the interaction of O₃ and its degradation products with plant tissue constituents (9). Cellular antioxidant systems are a front-line defense against oxygen free radicals. The activity of SOD,² one such enzymatic antioxidant, has been shown to increase in response to O₃ (7, 8, 14). Whether this increase is part of a protective, defense response or is a secondary consequence of injury is subject to debate. Lee and Bennett (12) reported a substantial increase in SOD activity in bean plants treated with EDU, a compound capable of conferring tolerance on O₃-susceptible plants (4). They also attributed the “natural” tolerance of young *versus* old leaves in the O₃-susceptible BBL290 bean cultivar to a higher endogenous SOD content in young leaves. They suggested that SOD plays a major role in detoxifying O₃-induced superoxide radicals which are at least in part responsible for ozone injury.

We have been investigating the role of SOD in several O₃-stressed model systems including *Pisum sativum*, *Phaseolus vulgaris*, and *Nicotiana tabacum*. This report concerns to-

MATERIALS AND METHODS

Nicotiana tabacum (tobacco) cv Wisconsin 38 plants were grown from seeds produced by selfing original transformants overproducing petunia chloroplastic Cu/Zn-SOD (19). Plants for assay were selected by rooting on kanamycin (200 µg/mL). Three plant types were tested: the control W38.11, which had been transformed with the kanamycin resistance gene only, and independent transformants SOD 35/1.4 and SOD 13.22 (hereafter called S1 and S2, respectively), which overproduced the petunia chloroplastic Cu/Zn-SOD (19). Vegetative plants aged 3 to 4 months with 9 to 11 nonsenescent leaves were used, uniform plants being chosen within each experiment. The newest leaf over 2 cm long was designated as leaf 1. Phenotypically there were no distinguishing features between the control and the two SOD transformants.

Ozone fumigation tests were conducted with five plants of each genotype in a dynamic flow system contained within a greenhouse (13); an equal number were placed in a similar chamber minus O₃. Ozone was produced by a generator (OREC model 0341–0) supplied with high grade oxygen. Ozone concentration in the chamber was measured by the neutral potassium iodide method (10). A dosage of 0.30 ppm O₃ for 6 h was used because it was capable of eliciting a differential response from indicator tobacco cultivars Bel W3 and BelB, sensitive and tolerant, respectively. Forty-eight hours after ozonation, leaves 1 through 8 were rated for visible injury. The experiment was conducted three times. Injury data for leaves 5 through 8 were subjected to analysis of

¹ This is New Jersey Agricultural Experiment Station Publication D-01905–1–91, supported in part by state funds and by the United States Hatch Act. This work is also supported by the Cooperative State Research Service, U.S. Department of Agriculture, under Agreement No. 89–3471–4502, and by the DuPont Company.

² Abbreviations: SOD, superoxide dismutase; EDU, ethylene-diurea.




Figure 1. Quantitation of the chloroplastic Cu/Zn-SOD overexpression in transgenic tobacco on activity gels according to Beauchamp and Fridovich (1). SOD activity of W38.11 control tobacco (W) was compared with that of transgenic plants 35/1.4 (S1). Lanes were loaded with crude soluble leaf protein extract as follows: W, 1 to 3: 500, 250, and 100 μg ; S1, 1 to 9: 0.5, 1, 5, 10, 20, 30, 40, 50, and 60 μg .

variance; data from young leaves 1 through 4 were excluded from analysis inasmuch as these typically showed no injury.

The visible injury rating system integrated scores for intensity and extent of injury, the highest possible injury index being 100%. Seven categories of injury intensity were assigned the following values: (0) no injury; (0.05) slight silvering; (0.10) severe silvering; (0.25) small adaxial necrotic flecks; (0.50) larger adaxial flecks; (0.75) large bifacial necrotic areas; and (1.0) very large bifacial necrotic areas. The percentage of area/leaf exhibiting each range of severity was multiplied by the assigned value, and then the total sum per leaf was reported as injury index.

Immediately following ozonation in experiment 1, leaf 6 from nonozonated and ozonated plants was harvested, frozen in liquid N_2 , and stored at -70°C until prepared for analysis. Leaves were ground to a powder in liquid N_2 , then homogenized in ice-cold buffer (100 mM sodium phosphate, pH 7.8, 0.1 mM EDTA and 10 mM β -mercaptoethanol) at a ratio of 2 g leaf powder/10 mL buffer. After filtration through Miracloth (Calbiochem), the filtrate was centrifuged at 39,000g for 15 min at 4°C . Aliquots of the supernatant were frozen in liquid N_2 and stored at -70°C . Protein concentration was determined by the Bradford method (2). To assess the degree to which the SOD was overexpressed in the transformants, immunoblot and SOD activity gel analyses were employed as described earlier (1, 16). Immunoblots and activity gels were analyzed on an Ultrascan XL Laser Densitometer (LKB, Sweden), using GELSCAN software (Pharmacia).

RESULTS AND DISCUSSION

In transgenic tobacco, the overproduction of Cu/Zn-SOD was shown by immunoblot and gel activity analysis to be approximately 15 times that of the control genotype, with both transformed types showing similar levels of overexpression of the SOD, as measured by densitometric analysis of

the immunoblot and activity gels. There were no differences in SOD activity between nonexposed plants and O_3 -exposed plants sampled immediately after ozonation. Figure 1 shows SOD activity gel containing a dilution series of leaf extracts from transformant 35/1.4 (S1) as compared with the control tobacco transformant W38.11 (W). The fastest migrating band corresponds to the endogenous chloroplastic Cu/Zn-SOD of tobacco in the control samples (W) and the introduced petunia chloroplastic Cu/Zn-SOD together with the endogenous activity in the transgenic (S1) extracts as demonstrated earlier by Tepperman and Dunsmuir (19).

All genotypes exhibited visible foliar injury as a result of O_3 fumigation (Table I, Fig. 2). Leaves 1 through 4 of all three genotypes were nonsymptomatic, thus exhibiting O_3 resistance typical of young, developing leaves. In all genotypes, leaf 5 (which was nearly fully expanded) was intermediate in sensitivity, while maximum damage was seen in leaves 6 through 8 (which were fully expanded). Statistical analysis of injury data by leaf age (Table I) indicated a significant decrease in injury in only 4 of 24 possible comparisons between transgenic plants overexpressing SOD and control plants. Had SOD played a protective role, one would have expected, if not complete protection, at least a consistent decrease in injury in each comparison. The data demonstrate that a large amount of additional Cu/Zn-SOD in the chloroplast does not protect the ozonated plants.

We have also tested whether the acknowledged tolerance of Bel B tobacco relative to very sensitive Bel W3 involved elevated levels of SOD. Immunoblots and SOD activity gels showed the same level of SOD for the two cultivars, thus eliminating dissimilarities in SOD levels as a possible reason for their differing sensitivities (data not shown).

EDU is an O_3 protectant which, in our hands, completely protects a large number of plant species, including tobacco, pea, and beans from visible injury at the ozone dosage used in the present study. Lee and Bennett (12) proposed that the

Table I. Foliar Injury Index in Response to Ozone in Control and Transgenic Tobacco Overproducing SOD

Visible foliar injury index as a function of leaf age was measured in control (W38.11) and transgenic tobacco lines (S1 and S2) engineered to overproduce petunia chloroplastic Cu/Zn-SOD, 48 h subsequent to a 6 h exposure to 0.3 ppm ozone. The means and SE are presented for each of three experiments, with $n = 5$ for each individual genotype. Means in a horizontal row (within one experiment) followed by the same letter are not significantly different at the 0.05% level using an LSD test.

Leaf Number	Experiment 1			Experiment 2			Experiment 3		
	Control	S1	S2	Control	S1	S2	Control	S1	S2
5	11.6 ± 4.0a	26.2 ± 9.0a	24.0 ± 5.4a	7.8 ± 6.1a	5.6 ± 3.3a	1.0 ± 1.0a	27.6 ± 10a	18.2 ± 5.7a	5.2 ± 2.4b
6	31.2 ± 8.7a	48.4 ± 8.2a	42.8 ± 10.6a	29.0 ± 10b	6.2 ± 1.2a	9.0 ± 4.1a	47.0 ± 14.4a	33.8 ± 10.8a	10.6 ± 4.7b
7	26.2 ± 8.8a	44.2 ± 15.0a	61.2 ± 7.2b	13.8 ± 7.5a	32.6 ± 12.2a	29.8 ± 13.1a	44.0 ± 14.8a	55.5 ± 14.4b	32.8 ± 12.6a
8	6.6 ± 6.6b	43.0 ± 14.0a	44.4 ± 19.0a	5.6 ± 5.6a	19.0 ± 6.0a	9.2 ± 6.5a	40.7 ± 13.3a	56.5 ± 18.3a	42.2 ± 15.6a

protective action of EDU is the result of an increase in SOD activity. This idea has been refuted in two published studies (6, 7) and in our own work (L.H. Pitcher, E. Brennan, and B.A. Zilinskas, unpublished results). Neither does the present work with transgenic tobacco support the idea that higher SOD levels protect plants from O₃ damage.

The conclusion drawn from the present study that Cu/Zn-SOD overexpression does not *per se* guarantee O₃ tolerance extends a principle developed in other laboratories in which stress from paraquat-generated superoxide radicals was not ameliorated by increased SOD. Tepperman and Dunsmuir (19) showed that elevated chloroplastic Cu/Zn-SOD in tobacco was not able to reduce paraquat's toxicity. Studies with "engineered" prokaryotic and eukaryotic systems have pro-

vided examples in which higher levels of SOD activity promote greater sensitivity to paraquat (see refs. 17 and 18 and citations therein). Two related explanations have been offered. Elevated SOD activity results in increased H₂O₂ production when superoxide radicals undergo dismutation. The enzymes capable of removing H₂O₂ are not likely to be elevated as a consequence of the introduction of the SOD genes. Therefore, the H₂O₂ itself presents a new threat. More importantly, the likelihood of production of very toxic hydroxyl radicals is greatly increased via the interaction of the superoxide radicals and H₂O₂ in the Haber-Weiss reaction. In a complementary vein, Yim *et al.* (23) recently reported that Cu/Zn-SOD itself is able to catalyze hydroxyl radical production directly from H₂O₂. In our study, we report some cases (Table I) in which SOD-overproducing plants had more O₃-induced injury than the control plants, but we cannot provide anything but speculation at this time as to why this occurred.

While overexpression of SOD may not necessarily be beneficial, numerous studies with bacteria, protists, animals, and plants indicate that SOD activity is required for life in oxygen-containing atmospheres. Thus, a double mutant of *Escherichia coli* lacking all SOD had reduced viability and a high mutation rate and was very sensitive to paraquat; these characteristics could be counteracted by a plasmid that overproduced SOD (3). Likewise, a *Drosophila* SOD null mutant had increased sensitivity to O₂, reduced fertility, and shortened lifespan (15).

Finally, there are a few reports of protection provided against paraquat toxicity by organisms engineered to produce higher levels of SOD (for example, see refs. 5, 11, 21). Interestingly, at least in some cases where such has been noted, the levels of H₂O₂ detoxifying enzymes appear to increase in an adaptive response to the higher H₂O₂ levels generated via increased SOD activity (5, 11).

In summary, in this work we find no consistent protection provided to transgenic plants engineered to overproduce the chloroplastic Cu/Zn-SOD. It is clear from the examples listed above that there are many factors that enter into the response of organisms to oxidative stress, and the variation we see within individual experiments and among leaves of different ages from the same plant genotype is probably due to the complexity of the overall antioxidant system.

ACKNOWLEDGMENT

We gratefully acknowledge Dr. Richard Trout for assistance in the statistical analyses.




Figure 2. Visible foliar injury index in response to ozone in control (C, W38.11) tobacco and two independent transgenic lines (S1 and S2) that overproduce the petunia chloroplastic Cu/Zn-SOD. The ratings for three independent experiments are presented. Leaves 5 through 8 on five plants of each genotype were rated as described in "Materials and Methods." The indices presented are means ± SE of these values ($n = 20$). Means within one experiment, designated by the same letter, are not significantly different at the 0.05 level, using an LSD test. Details of the experimental protocol are described in the text.

LITERATURE CITED

1. **Beauchamp C, Fridovich I** (1971) Superoxide dismutase: improved assays and an assay applicable to acrylamide gels. *Anal Biochem* **44**: 276–287
2. **Bradford MM** (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein using the principles of protein dye-binding. *Anal Biochem* **72**: 248–254
3. **Carlioz A, Touati D** (1986) Isolation of superoxide dismutase mutants in *E. coli*: is superoxide dismutase necessary for aerobic life? *EMBO J* **5**: 623–630
4. **Carnahan JE, Jenner EL, Wat EKW** (1978) Prevention of ozone injury to plants by a new protectant chemical. *Phytopathology* **68**: 1225–1229
5. **Ceballos I, Delabar JM, Nicole A, Lynch RE, Hallelwell RA, Kamouss P, Sinet PM** (1988) Expression of transfected human Cu-Zn SOD gene in mouse L cells and NS20Y neuroblastoma cells induces enhancement of glutathione peroxidase activity. *Biochim Biophys Acta* **949**: 5864–5870
6. **Chanway CP, Runeckles VC** (1984) Effect of ethylene diurea (EDU) on ozone tolerance and superoxide dismutase activity in bush bean. *Environ Pollut Ser A Ecol Biol* **35**: 49–56
7. **Chanway CP, Runeckles VC** (1984) The role of superoxide dismutase in the susceptibility of bean leaves to ozone injury. *Can J Bot* **62**: 236–240
8. **Decleire M, De Cat W, de Temmerman L, Baeten H** (1984) Changes of peroxidase, catalase and superoxide dismutase activities in ozone-fumigated spinach leaves. *J Plant Physiol* **116**: 147–152
9. **Heath RL** (1980) Initial events in injury to plants by air pollutants. *Annu Rev Plant Physiol* **31**: 396–431
10. **Jacobs MB** (1960) *The Chemical Analysis of Air Pollutants*. Interscience Publishers Inc, New York, pp 224–226
11. **Kelner MJ, Bagnell R** (1990) Alteration of endogenous glutathione peroxidase, manganese superoxide dismutase, and glutathione transferase activity in cells transfected with a copper-zinc superoxide dismutase expression vector. *J Biol Chem* **265**: 10872–10875
12. **Lee EH, Bennett JH** (1982) Superoxide dismutase: a possible protective enzyme against ozone injury in snap beans (*Phaseolus vulgaris* L). *Plant Physiol* **69**: 1444–1449
13. **Leone I, Brennan E, Daines RH** (1966) Effect of nitrogen nutrition on the response of tobacco to ozone in the atmosphere. *J Air Pollution Control Assn* **16**: 191–196
14. **McKersie BD, Beversdorf WD, Hucl P** (1982) The relationship between ozone insensitivity, lipid-soluble antioxidants, and superoxide dismutase in *Phaseolus vulgaris*. *Can J Bot* **60**: 2686–2691
15. **Phillips JP, Campbell SG, Michaud D, Charbonneau M, Hilliker AJ** (1989) Null mutation of Cu/Zn SOD in *Drosophila* confers hypersensitivity to paraquat and reduced longevity. *Proc Natl Acad Sci USA* **86**: 2761–2765
16. **Scioli JR, Zilinskas BA** (1988) Cloning and characterization of a cDNA encoding the chloroplastic copper-zinc superoxide dismutase from pea. *Proc Natl Acad Sci USA* **85**: 7661–7665
17. **Scott MD, Meshnick SR, Eaton JW** (1987) Superoxide dismutase rich bacteria. Paradoxical increase in oxidant toxicity. *J Biol Chem* **262**: 3640–3645
18. **Seto NO, Hayashi S, Tener GM** (1990) Overexpression of Cu-Zn superoxide dismutase in *Drosophila* does not affect lifespan. *Proc Natl Acad Sci USA* **87**: 4270–4274
19. **Tepperman JM, Dunsmuir P** (1990) Transformed plants with elevated levels of chloroplastic SOD are not more resistant to superoxide toxicity. *Plant Mol Biol* **14**: 501–511
20. **Thomson WW** (1975) Effects of air pollutants on plant ultrastructure. In J Mudd, T Kozlowski, eds, *Responses of Plants to Air Pollution*. Academic Press, New York, pp 179–194
21. **Touati D** (1988) Molecular genetics of superoxide dismutases. *Free Radic Biol Med* **5**: 393–402
22. **United States Environmental Protection Agency** (1986) *Air Quality Criteria for Ozone and Other Photochemical Oxidants*, Vol 3. EPA Office of Research and Development, Research Triangle Park, NC, pp 8–84
23. **Yim MB, Chock PB, Stadtman ER** (1990) Copper, zinc superoxide dismutase catalyzes hydroxyl radical production from hydrogen peroxide. *Proc Natl Acad Sci USA* **87**: 5006–5010