

Title: The prospect of using cyanobacterial bicarbonate transporters to improve leaf photosynthesis in C₃ crop plants.

Authors: *G Dean Price**, *Murray R Badger* and *Susanne von Caemmerer*

Running Title: Chloroplast bicarbonate pumps

Address: Molecular Plant Physiology Cluster, Plant Science Division, Research School of Biology (RSB), Australian National University, Canberra, ACT 0200, Australia.

* Corresponding author:

A/Prof. G. Dean Price

Molecular Plant Physiology Cluster,

Research School of Biology,

Building #46, Sullivan's Creek Road,

Australian National University,

Canberra, ACT 0200, Australia.

Dean.Price@anu.edu.au

Definitions:

Rubisco = ribulose-1,5-bisphosphate carboxylase-oxygenase

CCM = CO₂ concentrating mechanism

DIC = dissolved inorganic carbon (mainly CO₂ and HCO₃⁻)

C_i = leaf intracellular CO₂ concentration (sub-stomatal cavity).

CA = carbonic anhydrase; an enzyme catalyst that inter-converts CO₂ and HCO₃⁻

WUE = water use efficiency; the amount of water required to yield one unit of grain or biomass.

Keywords: Bicarbonate transporters, leaf photosynthesis, modeling, crop improvement.

1. Introduction

The photosynthetic CO₂ fixing enzyme Rubisco arose some 3.5 billion years ago, in an environment when CO₂ was high and O₂ low. Under these conditions, it was CO₂ saturated and presumably performed well (Badger et al., 1998). However, since the advent of oxygenic photosynthesis, the levels of O₂ have risen dramatically and CO₂ has fallen to very low levels. This has gradually created conditions where CO₂ has become limiting for Rubisco and allowed O₂ to act as an alternative inhibitory substrate for the enzyme. To cope with these dramatic environmental changes, two major strategies have evolved to help Rubisco maximize its carboxylation rate at ambient levels of limiting CO₂. Firstly, the enzyme has evolved better kinetic properties where the K_mCO₂ has decreased and the ability to distinguish against O₂ has increased at the expense of catalytic rate (Badger et al., 1998). Alternatively, many photosynthetic organisms ranging from cyanobacteria, to algae to land plants have developed active CO₂ concentrating mechanisms to turbo-charge Rubisco's CO₂ supply at a minor metabolic cost (Badger et al., 1998). Most notably among plants this has led to the development of C₄ photosynthesis (Sage, 2004).

Most of the important grain crops (rice, wheat, barley, canola, soybean), tuber crops and vegetable crops are C₃ species and have applied the first strategy and lack any form of CO₂ concentrating mechanism (CCM) at the leaf or chloroplast level. Much of the inherent inefficiency in C₃ photosynthesis revolves around the need to gain CO₂ through passive diffusion through the leaf pores (stomata), across cell walls and cytoplasm, and eventually through to the chloroplasts. Diffusive resistances to CO₂ passage result in a drawdown of the effective CO₂ concentration in the chloroplast and C₃ plants have adopted strategies to maximize the diffusive conductivity for CO₂ by appressing chloroplasts against the intracellular airspaces and having large chloroplast surface area to leaf area ratios (Evans and von Caemmerer, 1996). Low chloroplast CO₂ concentrations exacerbate the CO₂ limitations and increase the wasteful Rubisco oxygenation reaction of RuBP to produce phosphoglycolate that must be recycled back to RuBP through a complex set of reactions known as the photorespiratory cycle. This is worsened by increased temperature with the affinity for CO₂ dropping and the oxygenase reaction being relatively enhanced (Kubien and Sage, 2008). To achieve acceptable high rates of photosynthetic CO₂ fixation typical C₃ species invest up to 30% of soluble protein and some 25% of leaf nitrogen into Rubisco protein. Evolution of the CCM in C₄ plants effectively circumvented a number of the inefficiencies, creating present day impetus for attempting to introduce C₄-CCMs into important C₃ crops such as rice (Hibberd et al., 2008).

However, while the C₄ CCM is one approach to elevating CO₂ around Rubisco, drawing from our knowledge of single cell CCMs in cyanobacteria (Price et al., 2008) there are also opportunities to elevate CO₂ around Rubisco at the individual leaf chloroplast level. These

prospects are expanded upon below, but in brief we consider two scenarios. The **first**, and simplest approach, is to consider the transplantation of cyanobacterial bicarbonate transporters to the C₃ chloroplasts to provide marginal but significant improvement in photosynthetic performance. The **second**, more elaborate, longer term objective would be to engineer a more functional cyanobacterial CCM in the chloroplast.

2. The cyanobacterial CO₂ concentrating mechanism (CCM)

Cyanobacteria have evolved an extremely efficient CCM (Fig 1; see below) - being able to concentrate CO₂ around Rubisco by a factor of up to 1000-fold. As a result, cyanobacterial CO₂ fixation has been able to retain a Rubisco with relatively high carboxylation rate, although lower selectivity between CO₂ and O₂, compared to the Rubisco in C₃ plants (Badger et al., 1998). Cyanobacterial cells also have high nitrogen-use efficiency as less nitrogen is devoted to Rubisco than in a C₃ plant (Badger et al., 1998). In addition, Rubisco within a cyanobacterium operates at near CO₂ saturation due to the action of the CCM, such that wasteful photorespiration is largely eliminated.

Given that an early cyanobacterial progenitor is considered to have become the original endosymbiont for chloroplast evolution in algae and land plants, the question arises as to why do present-day land plants lack any apparent chloroplast-based CCM? Cyanobacterial progenitors first appeared some 2.7 billion years ago (Buick, 1992), but it is almost certain that cyanobacteria have been subjected to periods of rapid evolutionary change throughout this period. In particular, the marked drop in CO₂ levels, and the rise in O₂ levels, that occurred around 400-350 million years ago (Berner, 1990) represents a likely trigger that forced the evolution of adaptations to cope with photorespiration and low efficiency CO₂ fixation (Fig 2). The poor availability of CO₂ in water where diffusion is 10⁴ times slower than in air and where large unstirred layers can exist, probably provided additional evolutionary pressure. In addition, with a pK_a around 6.4, CO₂ is a rarer species at alkaline pH, whereas HCO₃⁻ is considerably more abundant in many aquatic environments. Evolutionary adaptations to deal with these combined pressures would have included transporters for active uptake of dissolved inorganic carbon species (DIC; CO₂ and HCO₃⁻), the subsequent localised elevation of CO₂ around Rubisco and the partitioning of Rubisco into micro-compartments known as carboxysomes (Badger et al., 2002; Price et al., 2008) – see below. This may have also been the stage when micro-algae developed CCMs. If, as seems likely, cyanobacteria did not evolve fully functional CCMs until 350 Mya, then this is well after the first terrestrial plants are thought to have evolved from eukaryotic algae at around 450 Mya (Kenrick and Crane, 1997) and long after the original endosymbiotic event that gave rise to micro-algae at around 1.5 billion years ago (Dyall et al., 2004). This probably explains why present day crop

plants lack any form of chloroplast-based CCM derived from cyanobacterial or micro-algal ancestors.

The Cyanobacterial CCM functions to actively transport and accumulate DIC into the cell where the accumulated HCO_3^- pool is utilised to generate elevated CO_2 levels around Rubisco (Badger et al., 2002; Price et al., 2008). Rubisco is encapsulated in unique micro-compartments, known as carboxysomes that are typically 90-200 nm in diameter. The functional importance of these proteinaceous, icosahedral bodies that are composed of 20 equilateral triangular side, is that they act as the site of CO_2 elevation within the cell, with the supply rate of CO_2 from accumulated HCO_3^- being catalysed by a carboxysome-located carbonic anhydrase (CA). The carboxysome shell in these cyanobacteria is composed of just 6-8 proteins (Price et al., 2008) and the average unicellular cyanobacterial cells would normally possess 5-15 carboxysomes per cell. The key to the efficiency of any CCM revolves around the ability to minimise the loss of CO_2 from the elevation zone. In model cyanobacteria, this is accomplished by a combination of (a) accumulation of the ionic form of DIC which is less membrane permeable than CO_2 , (b) the complete elimination of CA activity from the general cytosol to help reduce CO_2 leakage out of the cell, (c) the special properties of the carboxysome protein shell acting to retard CO_2 leakage, and (d) the action of the CO_2 pumps in recycling CO_2 leakage from the carboxysome back into HCO_3^- pool (Maeda et al., 2002; Price et al., 2008).

The localisation of CA, which catalyses the reversible hydration and dehydration of CO_2 and HCO_3^- , is a key element of cyanobacterial CCMs. The absence of CA in the cytosol, and the action of the directional CO_2 uptake systems that convert CO_2 to HCO_3^- at the thylakoid membrane, allows the cell to accumulate HCO_3^- and keep it out of rapid chemical equilibrium with CO_2 . This is very effective in minimizing the concentration of the diffusible CO_2 molecule owing to the slow dehydration of HCO_3^- in the absence of CA (Walker et al., 1980). The importance of accumulating HCO_3^- in the cytosol, and maintaining an internal HCO_3^- pool out of chemical equilibrium, was shown by an experiment where human CA was expressed in the cytoplasm of a model cyanobacterium, *Synechococcus elongatus* PCC7942. The ectopic expression caused complete dissipation of the accumulated HCO_3^- pool due to the CA-mediated equilibration between CO_2 and HCO_3^- which in turn lead to increased CO_2 diffusion out of the cell (Price and Badger, 1989). This is very different to the situation in C_3 chloroplasts where CA is highly abundant in the stroma in order to maximise diffusion of CO_2 across the envelope and throughout the chloroplast (Badger and Price, 1994).

Five distinct transport systems for DIC uptake have been identified in cyanobacteria (Fig. 1 and Table 1; see (Price et al., 2008) for more details and related references:-

- (i) **BCT1**, is inducible under DIC limitation, and is a high affinity HCO_3^- transporter (uniporter) belonging to the traffic ATPase family.
- (ii) **SbtA**, an inducible, high affinity Na^+ -dependent HCO_3^- transporter (Price et al., 2004; Shibata et al., 2002) that apparently acts as a $\text{Na}^+/\text{HCO}_3^-$ symporter with relatively low flux rate.
- (iii) **BicA**, a low affinity, high flux, Na^+ -dependent HCO_3^- transporter belonging to the widespread SulP family and related to the human SLC26 family of anion transporters (Price et al., 2004); BicA is a probable $\text{Na}^+/\text{HCO}_3^-$ symporter.
- (iv) **NDH-I₄**, a constitutive CO_2 uptake system based on a specialised NADPH dehydrogenase (NDH-I) complex; this system uses NADPH as an electron donor to drive conversion of CO_2 to HCO_3^- during the uptake step (Price et al., 2002); each complex is composed of ten core subunits that are common to the respiratory NDH-I complex and 3 specialised subunits required for CO_2 uptake. Interestingly, NDH-I type CO_2 uptake systems appear to be located on the thylakoid membranes where they use CO_2 diffusing from outside the cell, or arising from leakage from the carboxysomes, as a substrate for directional conversion to HCO_3^- .
- (v) **NDH-I₃**, a second CO_2 uptake system based on a modified NDH-I complex that is inducible under DIC limitation and is of higher uptake affinity than NDH-I₄; located on the thylakoid membranes in *Synechocystis* PCC6803.

3. Adding a HCO_3^- transporter to the chloroplast envelope of crop plants

With the objective of attaining a modest elevation of CO_2 levels in the C_3 chloroplast, the simplest approach would be to express a cyanobacterial HCO_3^- transporter on the inner envelope of the C_3 chloroplast (Fig. 3a). Single-subunit HCO_3^- transporters such as BicA and SbtA are the most obvious initial candidates. However, within technical restraints the transfer of multi-subunit transporters such as the BCT1 HCO_3^- transporter (4 genes) is also possible. Additionally, the use of HCO_3^- transporters from micro-algae such as *Chlamydomonas* can also be considered as viable candidates (Duanmu et al., 2009). From a technical viewpoint, addition of DIC transporters mentioned above would be dependent on host genome transformation techniques using *Agrobacterium*, which are generally available for a range of important crop species. Chloroplast transformation techniques would not be required for this approach and this is especially important since chloroplast transformation in crop species is not yet available. As can be seen from the associated modeling presented in this report (Fig 4), the approach of installing BicA and/or SbtA transporters into the chloroplast inner envelope could achieve a 5-15% improvement in photosynthetic CO_2 fixation rates at constant sub-stomatal CO_2 levels (see below).

It is clear that a CO_2 diffusion gradient or drawdown exists between the CO_2 level in the sub-stomatal cavity of the leaf (C_i) and the steady state level of CO_2 in chloroplast (C_{chlo}), with the magnitude of this gradient sitting about 40% below C_i at high irradiance (Evans and von

Caemmerer, 1996). It is important to recognize that in the first instance, the primary objective of adding a HCO_3^- pump would be to diminish the size of this CO_2 drawdown at the chloroplast and not to elevate it significantly above the external CO_2 concentration. This minimizes the risk of wasteful CO_2 leakage. Such a situation is very similar to the concept of introducing a C_4 cycle into C_3 cells (Matsuoka et al., 2001), which has been modeled for transplantation into a typical C_3 chloroplast (von Caemmerer, 2003) and found to be theoretically capable of raising the steady state CO_2 level within the chloroplast. More specific modeling data on the theoretical engineering of BicA into a chloroplast is shown in Fig. 4 and discussed below. Notably, for the addition of HCO_3^- pumps to work there is no need to modify CA levels in the chloroplast since CA is needed to promote rapid equilibrium between accumulated HCO_3^- and CO_2 within the stroma.

In terms of establishing active HCO_3^- uptake across the chloroplast envelope the question arises as to whether a Na^+ -dependent HCO_3^- transporter could function in a chloroplast. Estimates indicate at least $250 \mu\text{M}$ HCO_3^- is present in the cytosol of a leaf cell in ambient air (Evans and von Caemmerer, 1996) and this appears to be maintained by cytosolic CA activity. The uptake affinities of SbtA (low flux rate) and BicA (high flux rate) for HCO_3^- in cyanobacteria are 5-15 and 90-170 μM , respectively (Price et al., 2004; Shibata et al., 2002) and would indicate that either transporter would operate well above its intrinsic K_m . An additional concern relates to the question of energisation and the involvement of Na^+ gradients across the chloroplast envelope. Both SbtA and BicA require about 1mM Na^+ for half maximal activity in the form of a standing inward gradient for Na^+ (Price et al., 2004; Shibata et al., 2002). The leaf cytosol possesses 1-3 mM Na^+ (Karley et al., 2000) and proteomic analyses have revealed that the *Arabidopsis* chloroplast envelope possesses several potential Na^+ -coupled transporters and Na^+/H^+ antiporters that are homologous to cyanobacterial forms (Rolland et al., 2003). Thus, there are good prospects that the chloroplast possesses and maintains an inwardly-directed Na^+ gradient. As a potential enhancement, the transfer of a cyanobacterial Na^+/H^+ antiporter (or a version from the C_4 chloroplast) could also be considered if this Na^+ gradient needed to be augmented, perhaps at the expense of any existing H^+ gradient (proton motive force) inferred from the existence of H^+ -coupled transporters in the envelope (Weber et al., 2005). Any possible perturbation of osmotic and pH homeostasis in the chloroplast by elevating steady state HCO_3^- levels by up to 15%, or even by as much as 25-fold relative to air exposed leaves, would be expected to be minimal (Wagner et al., 1990).

Other problems in establishing SbtA or BicA in C_3 chloroplasts would include ensuring the correct targeting to the chloroplast envelop and uncertainty about whether these transporters need to be post-translationally activated. In the case of targeting, we expect that SbtA and BicA can be fused to the cDNAs for known envelope-targeted proteins such that details on targeting are not initially required. We have determined the membrane topology structure of BicA and SbtA as an

initial step in identifying the most likely cytoplasmic regulatory domains in these transporters (Shelden et al., 2010).

It should be noted that there has been one attempt to place a putative cyanobacterial DIC transporter, *IctB*, in *Arabidopsis* and tobacco resulting in a reported improvement in water use efficiency (WUE, the ratio of CO₂ assimilation rate to transpiration rate) in *Arabidopsis* plants grown under dry air conditions; in particular, a drop in the CO₂ compensation point was observed (Lieman-Hurwitz et al., 2003). The basis of this improved WUE is not yet clear since it is now known that *ictB* does not code for a DIC transporter (Price et al., 2008; Shibata et al., 2002) and its role in cyanobacteria is still unclear.

4. Modeling the addition of bicarbonate pumps to the C₃ chloroplast

Our modeling of the consequences of taking the first step of adding one or two cyanobacterial HCO₃⁻ transporters to a C₃ chloroplast is based on previous approaches used to consider the theoretical addition of a CO₂ pump of single-cell-C₄ type of von Caemmerer & Furbank (von Caemmerer, 2003; von Caemmerer and Furbank, 2003); equations and parameters used in the simulations shown in Fig. 4 are detailed in the supplementary file. Much of the discussion of the benefits of introduction of single-cell-C₄ photosynthesis into a C₃ leaf, applies to the introduction of bicarbonate transporters (von Caemmerer 2003). The key point issuing from the modeling is that the addition of either HCO₃⁻ transporter, *BicA* or *SbtA* can lead to an increase in the rate of light saturated CO₂ assimilation at ambient and low intercellular CO₂ partial pressures, C_i. The magnitude of the increase will be very much dependent on the kinetic properties of the transporters and the conductance to CO₂ diffusion of the chloroplast envelope (von Caemmerer 2003). The introduction of a transporter elevates chloroplast CO₂ partial pressures, C_{chlo}, above C_i at low C_i values, resulting in a reduced CO₂ compensation point. The addition of the high affinity *SbtA* transporter is more effective at reducing the compensation point than the *BicA* transporter because of its lower K_m, and introduction of both can be more effective again. At higher C_i levels, transporters serve to reduce the drawdown in CO₂ between intercellular CO₂ and the chloroplast (Fig. 4). At a constant C_i level of 250 μbar, a theoretical transgenic plant, with the assumed activities of both HCO₃⁻ uptake systems, could display an indicative assimilation rate >15% higher than wild type C₃.

One of the disadvantages of C₄ photosynthesis is the requirement of 2 ATP per CO₂ fixed in the C₄ cycle which makes the introduction of a C₄ cycle into current C₃ leaves energetically inefficient (von Caemmerer 2003), although there are examples of a number of single-cell C₄ species that have overcome the anatomical limitations inherent in C₃ leaves (Edwards et al. 2004). It is likely that bicarbonate transport is less expensive. Considering the negative electrogenicity of HCO₃⁻ uptake in cyanobacteria (Ritchie et al., 1996), and the likely Na⁺:proton equivalence, we

estimate a required $\frac{1}{4}$ ATP per HCO_3^- transported by BicA and $\frac{1}{2}$ ATP per HCO_3^- for SbtA. On this basis, ATP requirement per net CO_2 assimilation rate drops with increasing C_i during C_3 photosynthesis as the cost of photorespiration decreases (Fig.4) The introduction of bicarbonate transporters reduces the ATP cost at low C_i below that normally experienced during C_3 photosynthesis and increases marginally above the C_3 requirement at higher C_i (Fig.4).

We have only considered the implication for light saturated photosynthesis here. For healthy crops about half the canopy will be operating under these conditions which should make introduction of bicarbonate transporters a productive strategy. Enhancing leaf photosynthetic rates also has the potential of increasing leaf water use efficiency, WUE, depending on stomatal responses. Bicarbonate transporters provide the largest benefit a low C_i , and the stomata of a plant with HCO_3^- pump enhancement could afford to be less open whilst providing the same rate of assimilation, thereby resulting in less loss of water from the leaf stomates. The SbtA transporter could be capable of improving WUE under dry air conditions more effectively than BicA, and addition of both transporters is likely to be even better. If SbtA or SbtA+BicA can be successfully introduced into C_3 plants, such species might be able to better survive transient episodes of high water deficit under high light and moderate temperature stress.

5. Adding a more elaborate cyanobacterial CCM to the chloroplast

A longer-term objective, involving a greater technical difficulty of the introduction of multiple genes, could be to establish a more elaborate form of the cyanobacterial CCM in the chloroplast (Fig. 3b). This could involve the transfer of one or two functionally active HCO_3^- transporters to the inner envelope membrane, combined with the transfer a CO_2 uptake system to the thylakoid membranes and a Rubisco micro-compartment such as the carboxysomes. The C_3 chloroplast would also need to be, like the cyanobacterial cytosol, converted to a HCO_3^- accumulating organelle where the HCO_3^- pool is held in a state of slow chemical interconversion. To do this, it would be necessary to re-organise chloroplastic C_3 Rubisco into effective carboxysome structures, and devise an effective means of removing the highly abundant chloroplastic CA so that HCO_3^- accumulation can be optimised. Certainly, it has been possible to remove up to 99% of chloroplastic CA activity in tobacco leaves by antisense RNA approaches (Price et al., 1994). Ideally, a complete removal of CA from the stroma would be more desirable, except for retaining the critically important CA in transferred carboxysomes. One of the most significant uncertainties relates to the conductance of the envelope to CO_2 diffusion, with a range of estimates available (Flexas et al., 2008). Aquaporins seem to play a role in CO_2 conductance, thus a useful enhancement might be to reduce aquaporin levels in the envelope by iRNA technology since lower conductance would aid in reducing CO_2 leakage (Flexas et al., 2006).

Our understanding of carboxysome assembly and function has improved greatly in recent years, to the point where engineering the assembly of a carboxysome in the chloroplast is approaching feasibility. This has been aided by advances in determining the crystal structures of some key components of the shell (Yeates et al., 2008) and our own efforts to identify proteins required as key Rubisco- and shell-organizing proteins (Long et al., 2007; Long et al., 2010). The remarkable feature of the small shell proteins is an ability to self-assemble (Yeates et al., 2008). This property, shared with some virus coat proteins, could greatly aid the final goal of assembling functional carboxysomes within the chloroplast. The longer-term objective of engineering a more potent form of the cyanobacterial CCM into the chloroplast may provide greater photosynthetic enhancements than the introduction of bicarbonate transporters alone.

6. Conclusions

Modeling indicates that the addition of cyanobacterial (or micro-algal) HCO_3^- pumps at the chloroplast envelope of a typical C_3 plant could provide a significant boost to the photosynthetic performance of leaf photosynthesis, either as increased assimilation rate or as improved WUE. A present research focus is to target BicA and SbtA transporters to the chloroplast of a model C_3 plant and to extend our understanding of activation and energisation processes for cyanobacterial HCO_3^- transporters. However, it should also be clear that parallel transgenic strategies such as improving the performance of Rubisco or raising capacities for RuBP generation or light interception would provide complementary improvements to crop performance, as discussed in other articles in this issue.

Figure captions:

Figure 1: The cyanobacterial CO₂ concentrating mechanism (CCM) utilizes up to five uptake systems for DIC and the polyhedral micro-compartments known as carboxysomes that contain the cell's complement of Rubisco and act as a localized site for elevation of CO₂ around Rubisco. A key operational feature is that all uptake systems deliver HCO₃⁻ to the general cytoplasm and that this is kept in a state of dynamic disequilibria favoring HCO₃⁻, owing to the recycling of internally generated CO₂ through the CO₂ pumps and the absence of CA in the general cytoplasm. A specific, low level of CA activity is only present in the carboxysomes. In general, BicA, SbtA, BCT1 and NDH-I₃ uptake systems are only induced under DIC limiting conditions (e.g. liquid in near-equilibrium with air CO₂ levels or less). The carboxysomes are typically 90-200 nm diameter (enlarged in this schematic) and a cell may possess 5-15 carboxysomes. By comparison typical unicellular cyanobacteria are up to 3 μm in length and typical C₃ chloroplasts are up to 50 μm in diameter.

Figure 2: A timeline indicating that CO₂ concentrating mechanisms (CCMs) possibly arose in cyanobacteria and micro-algae at around 400-350 mya, and well after the evolution of early land plants.

Figure 3: Schematic representations illustrating the concepts of adding a cyanobacterial HCO₃⁻ transporter to the chloroplast envelope of a notional C₃ leaf chloroplast (a) and the longer-term prospect of constructing a more fully functional cyanobacterial or micro-algal CCM in the C₃ chloroplast (b). The diagrams show CO₂ moving from the intracellular airspace (IAS; sub-stomatal cavity) of a mesophyll leaf cell through the cell wall to the cytoplasm (Cyt) before entering the chloroplast by CO₂ diffusion or via entry through a HCO₃⁻ transporter. The hexagonal structure represents the icosahedral carboxysomes that would contain the full complement of Rubisco in the chloroplast with a specific carbonic anhydrase (CA) partitioned to this compartment and stromal CA removed. Linkages between the carbon reduction in the chloroplast and photorespiration involving peroxisomes (P) and mitochondria (M) are also shown.

Figure 4 : a) Modeled net CO₂ assimilation rate (A) as a function of intercellular CO₂ partial pressure, C_i, with three different options for bicarbonate transport at the chloroplast envelope. 1) a BicA transporter with a maximum activity of 30 μmol m⁻²s⁻¹ and k_{1/2} for HCO₃⁻ = 90 μM (~140 μbar CO₂ at pH=7.4). 2), a SbtA transporter with a maximum activity of 15 μmol m⁻²s⁻¹ and k_{1/2} for HCO₃⁻ = 5 μM (~11 μbar CO₂ at pH=7.4), 3) with both transporters. Only light saturated CO₂ assimilation is considered and maximal Rubisco activity was V_{cmx}=100 μmol m⁻²s⁻¹. The diffusive conductance across the cell wall plasmalemma

interface, and across the chloroplast envelope were, 1 and $0.5 \text{ mol m}^{-2} \text{ s}^{-1} \text{ bar}^{-1}$, respectively. This results in a total conductance $=0.333 \text{ mol m}^{-2} \text{ s}^{-1} \text{ bar}^{-1}$. CO_2 assimilation rates are compared to Rubisco limited C_3 photosynthesis with the same maximal Rubisco activity. Other parameters and model equations used are given in the supplementary document. (The simulation becomes unrealistic at higher C_i as it does not consider RuBP regeneration limitation). **b)** Difference between chloroplast and intercellular CO_2 ($\text{C}_{\text{chlo}} - \text{C}_i$) as a function of C_i for the examples given in a). **c)** ATP consumption per net CO_2 assimilation rate for the examples given in a). Considering the proton transport required we estimate that $\frac{1}{4}$ ATP per HCO_3^- transported by BicA and $\frac{1}{2}$ ATP per HCO_3^- transported by SbtA.

Table 1: a summary of the properties of cyanobacterial DIC transporters

Transport Type	Mechanism	Substrate affinity	Flux rate	Photosynthetic affinity, $K_{0.5}$
BicA	Na ⁺ -dependent HCO ₃ ⁻ uptake	Low-medium	High	90-170 μM HCO ₃ ⁻
SbtA	Na ⁺ -dependent HCO ₃ ⁻ uptake	High	Low	< 5 μM HCO ₃ ⁻
BCT1	Traffic ATPase, HCO ₃ ⁻ uptake	High	Low	10-15 μM HCO ₃ ⁻
NDH-I ₄	NADPH-driven CO ₂ uptake via conversion to HCO ₃ ⁻	Medium	High	10-15 μM CO ₂
NDH-I ₃	NADPH-driven CO ₂ uptake via conversion to HCO ₃ ⁻	High	Low	1-2 μM CO ₂

References

- Badger MR, Andrews TJ, Whitney SM, Ludwig M, Yellowlees DC, Leggat W, Price GD** (1998) The diversity and coevolution of rubisco, plastids, pyrenoids, and chloroplast-based CO₂-concentrating mechanisms in algae. *Can J Bot* 76: 1052-1071
- Badger MR, Hanson D, Price GD** (2002) Evolution and diversity of CO₂ concentrating mechanisms in cyanobacteria. *Funct Plant Biol* 29: 161-173
- Badger MR, Price GD** (1994) The role of carbonic anhydrase in photosynthesis. *Ann Rev Plant Physiol Plant Molec Biol* 45: 369-392
- Berner RA** (1990) Atmospheric carbon dioxide levels over phanerozoic time. *Science* 249: 1382-1386
- Buick R** (1992) The Antiquity of Oxygenic Photosynthesis: Evidence from Stromatolites in Sulphate-Deficient Archaean Lakes. *Science* 255: 74
- Duanmu D, Miller AR, Horken KM, Weeks DP, Spalding MH** (2009) Knockdown of limiting-CO₂-induced gene HLA3 decreases HCO₃⁻ transport and photosynthetic Ci affinity in *Chlamydomonas reinhardtii*. *Proc Natl Acad Sci (USA)* 106: 5990-5995
- Dyall SD, Brown MT, Johnson PJ** (2004) Ancient invasions: From endosymbionts to organelles. *Science* 304: 253-257
- Evans JR, von Caemmerer S** (1996) Carbon Dioxide Diffusion inside Leaves. *Plant Physiol* 110: 339-346
- Flexas J, Ribas-Carbo M, Diaz-Espej A, Galmes J, Medrano H** (2008) Mesophyll conductance to CO₂: current knowledge and future prospects. *Plant Cell Environ* 31: 602-621
- Flexas J, Ribas-Carbo M, Hanson DT, Bota J, Otto B, Cifre J, McDowell N, Medrano H, Kaldenhoff R** (2006) Tobacco aquaporin NtAQP1 is involved in mesophyll conductance to CO₂ in vivo. *Plant J* 48: 427-439
- Hibberd JM, Sheehy JE, Langdale JA** (2008) Using C-4 photosynthesis to increase the yield of rice - rationale and feasibility. *Curr Op Plant Biol* 11: 228-231
- Karley AJ, Leigh RA, Sanders D** (2000) Where do all the ions go? The cellular basis of differential ion accumulation in leaf cells. *Trends Plant Sci* 5: 465-470
- Kenrick P, Crane PR** (1997) The origin and early evolution of plants on land. *Nature* 389: 33-39
- Kubien DS, Sage RF** (2008) The temperature response of photosynthesis in tobacco with reduced amounts of Rubisco. *Plant Cell Environ* 31: 407-418
- Lieman-Hurwitz J, Rachmilevitch S, Mittler R, Marcus Y, Kaplan A** (2003) Enhanced photosynthesis and growth of transgenic plants that express *ictB*, a gene involved in HCO₃⁻ accumulation in cyanobacteria. *Plant Biotech J* 1: 43-50
- Long BM, Badger MR, Whitney SM, Price GD** (2007) Analysis of carboxysomes from *Synechococcus* PCC7942 reveals multiple Rubisco complexes with carboxysomal proteins CcmM and CcaA. *J Biol Chem* 282: 29323-29335

Long BM, Tucker L, Badger MR, Price GD (2010) Functional Cyanobacterial beta-Carboxysomes Have an Absolute Requirement for Both Long and Short Forms of the CcmM Protein. *Plant Physiol* 153: 285-293

Maeda S, Badger MR, Price GD (2002) Novel gene products associated with NdhD3/D4-containing NDH-1 complexes are involved in photosynthetic CO₂ hydration in the cyanobacterium, *Synechococcus* sp. PCC7942. *Molec Microbiol* 43: 425-435

Matsuoka M, Furbank RT, Fukayama H, Miyao M (2001) Molecular engineering of C-4 photosynthesis [Review]. *Ann Rev Plant Physiol Plant Molec Biol* 52: 297-314

Price GD, Badger MR (1989) Expression of human carbonic anhydrase in the cyanobacterium *Synechococcus* PCC7942 creates a high CO₂-requiring phenotype. Evidence for a central role for carboxysomes in the CO₂ concentrating mechanism. *Plant Physiol* 91: 505-513

Price GD, Badger MR, Woodger FJ, Long BM (2008) Advances in understanding the cyanobacterial CO₂-concentrating-mechanism (CCM): functional components, Ci transporters, diversity, genetic regulation and prospects for engineering into plants. *J Exp Bot* 59: 1441-1461

Price GD, Maeda S, Omata T, Badger MR (2002) Modes of active inorganic carbon uptake in the cyanobacterium, *Synechococcus* sp PCC7942. *Funct Plant Biol* 29: 131-149

Price GD, von Caemmerer S, Evans JR, Yu JW, Lloyd J, Oja V, Kell P, Harrison K, Gallagher A, Badger MR (1994) Specific reduction of chloroplast carbonic anhydrase activity by antisense RNA in transgenic tobacco plants has a minor effect on photosynthetic CO₂ assimilation. *Planta* 193: 331-340

Price GD, Woodger FJ, Badger MR, Howitt SM, Tucker L (2004) Identification of a SulP-type bicarbonate transporter in marine cyanobacteria. *Proc Natl Acad Sci (USA)* 101: 18228-18233

Ritchie RJ, Nadolny C, Larkum AWD (1996) Driving forces for bicarbonate transport in the cyanobacterium *Synechococcus* R2 (PCC 7942). *Plant Physiol* 112: 1573-1584

Rolland N, Ferro M, Seigneurin-Berny D, Garin J, Douce R, Joyard J (2003) Proteomics of chloroplast envelope membranes. *Photosyn Res* 78: 205-230

Sage RF (2004) The evolution of C-4 photosynthesis. *New Phytol* 161: 341-370

Shelden MC, Howitt SM, Price GD (2010) Membrane topology of the cyanobacterial bicarbonate transporter, BicA, a member of the SulP (SLC26A) family. *Mol Membr Biol* 27: 12-23

Shibata M, Katoh H, Sonoda M, Ohkawa H, Shimoyama M, Fukuzawa H, Kaplan A, Ogawa T (2002) Genes essential to sodium-dependent bicarbonate transport in cyanobacteria - Function and phylogenetic analysis. *J Biol Chem* 277: 18658-18664

von Caemmerer S (2003) C₄ photosynthesis in a single C₃ cell is theoretically inefficient but may ameliorate internal CO₂ diffusion limitations of C₃ leaves. *Plant Cell Environ* 26: 1191-1197


von Caemmerer S, Furbank RT (2003) The C₄ pathway: an efficient CO₂ pump. *Photosyn Res* 77: 191-207

Wagner U, Kolbowski J, Oja V, Laisk A, Heber U (1990) pH Homeostasis of the Chloroplast Stroma Can Protect Photosynthesis of Leaves During the Influx of Potentially Acidic Gases. *Biochim Biophys Acta* 1016: 115-120

Walker NA, Smith FA, Cathers IR (1980) Bicarbonate Assimilation by Freshwater Charophytes and Higher-Plants. 1. Membrane-Transport of Bicarbonate Ions Is Not Proven. *J Membr Biol* 57: 51-58

Weber APM, Schwacke R, Flugge UI (2005) Solute transporters of the plastid envelope membrane. *Ann Rev Plant Biol* 56: 133-164

Yeates TO, Kerfeld CA, Heinhorst S, Cannon GC, Shively JM (2008) Protein-based organelles in bacteria: carboxysomes and related microcompartments. *Nature Rev Microbiol* 6: 681-691



Freshwater β -Cyanobacteria

