

Root Systems Biology: *integrative modeling across scales, from gene regulatory networks to the rhizosphere*

Kristine Hill¹, Silvana Porco¹, Guillaume Lobet², Susan Zappala¹, Sacha Mooney¹
Xavier Draye³ and Malcolm J Bennett^{1*}

¹The Centre for Plant Integrative Biology, School of Biosciences, University of Nottingham, Sutton Bonington Campus, Loughborough, LE12 5RD, UK.

²PhytoSYSTEMS, Université de Liège, Boulevard du Rectorat 27, 4000 Liège, Belgium.

³ Earth and Life Institute, Université catholique de Louvain, Croix du Sud 2 – L7.05.11, 1348, Louvain-la-Neuve, Belgium.

*Corresponding author

malcolm.bennett@nottingham.ac.uk

FON +44 (0) 115 951 3255

FAX +44 (0) 115 951 6334

Word and figure count

Abstract – 146

Main Text – 9926

Figures – 3

Tables - 1

Abstract

Genetic and genomic approaches in model organisms have advanced our understanding of root biology over the last decade. Recently, however, systems biology and modeling have emerged as important approaches, as our understanding of root regulatory pathways has become more complex and interpreting pathway outputs has become less intuitive. To relate root genotype to phenotype, we must move beyond examination of interactions at the genetic network scale and employ multi-scale modeling approaches to predict emergent properties at the tissue, organ, organism and rhizosphere scales. Understanding the underlying biological mechanisms and the complex interplay between systems at these different scales requires an integrative approach. Here, we describe examples of such approaches and discuss the merits of developing models to span multiple scales, from network to population levels, and to address dynamic interactions between plants and their environment.

136 words

1 Introduction

Root architecture critically influences nutrient and water uptake efficiency and plays a central role in plant productivity (Lynch, 1995); selecting new crop varieties with improved root traits may produce a second '*Green Revolution*' (Lynch, 2007). Over the past several decades, reductionist approaches have pinpointed the individual genes or mechanisms that control root system architecture (reviewed in Benfey et al., 2010). However, as our knowledge of biological systems has increased, researchers have realized that the underlying components (e.g. gene products, cells, tissues and organs) function in highly complex, dynamic networks. The existence of emergent traits, robustness, and hierarchical organization of biological systems make them challenging to conventional reductionist experimental approaches (Bruggeman and Westerhoff, 2007). Understanding these properties requires studying the system, rather than its individual components. Also, in contrast to the simple linear networks conveyed in textbooks, biological networks usually contain multiple branches, feedback and/or feed-forward loops, and other complex regulatory motifs. For example, most signal transduction pathways include negative feedback and/or positive feed-forward loops to switch off or amplify a response pathway. Hence, logic alone often fails to predict the output from such pathways. Systems approaches often employ mathematical or computational models to simulate the behaviors of these non-linear networks and predict emergent behaviors (reviewed in Middleton et al., 2012). Robust biological systems also have compensatory mechanisms that come into play when a key gene is removed. For example, 65% of knock-out lines for *Arabidopsis* transcription factors in the root stele gene regulatory network (GRN) showed molecular phenotypes, but only 16% showed morphological phenotypes (Brady et al. 2011). Thus, in the robust stele GRN, the network architecture or genetic redundancy can buffer or canalize changes in gene expression. Hierarchy is inherent to all biological systems. At scales relevant to agriculture, hierarchy starts at the field scale. Field-scale populations comprise individual plants, which are composed of organs, tissues, cells, organelles and molecules, respectively. Each hierarchical level interacts with higher and lower scales. Hence, understanding the underlying biological mechanisms of the complex, critical interplay between these scales requires an integrative systems approach.

Root researchers currently use several distinct systems-based approaches. 'Top-down' approaches automatically analyze large-scale datasets to uncover relationships between levels of transcripts and/or proteins (reviewed in Bassel et al., 2012). This approach starts with experimental 'omics' data, followed by data analysis and data integration to determine correlations, and ends with the formulation of hypotheses concerning co- and inter-regulation of groups of molecules. These hypotheses predict new correlations that can be iteratively tested (Bruggeman and Westerhoff, 2007).

Another approach, termed 'bottom-up,' constructs detailed models of well-characterized networks that can be simulated mathematically and/or computationally. This approach starts with a manageable part of a system, such as a small GRN, and employs mechanism-based models (reviewed in Middleton et al., 2012). Bottom-up studies take an integrative approach to look for emergent properties of a network with available quantitative information such as kinetic data, transcription rates and protein stability. As information about GRN increases, so does the opportunity to implement mechanistic models for bottom-up studies.

'Middle-out' approaches construct models starting at the level for which we have the most information and combining predictive bottom-up and inference-based top-down approaches, thus building on the current body of knowledge (Noble, 2008). The middle-out approach can start at any scale. However, cells provide a natural level of organization for an organism, cell-based models can endow individual cells with considerable internal machinery (representing the subcellular networks) and the interactions of numerous individual cells can lead to the (emergent) tissue-scale properties. This makes cell-based models an attractive approach, as they naturally incorporate heterogeneity via cellular properties.

'Multi-scale' approaches employ models that consider behaviors on two or more spatial scales, ranging from the subcellular up to whole-organism and beyond. Multi-scale models must also incorporate temporal scales ranging over many orders of magnitude. Spatially, systems can be measured at the molecular scale (10^{-10} m) through to the whole plant scale (0.1-100 m range), or up to the field level (>1000 m).

Likewise, temporal scales can range from nanoseconds (10^{-9} s) for enzymatic reaction to months and years (10^7 s) for a plant lifecycle. Given this range of temporal and spatial scales, multi-scale models are often challenging to construct and relatively few examples involve plant roots (Band et al., 2012a).

Although our review focuses on roots, we do not ignore links to the whole plant and field systems. We ultimately study roots because they take up water and nutrients, which influences crop yield and environmental adaptation. While the first point – roots are fundamental to the whole plant – is obvious, the second point – the effect of the whole plant context on the root – is often overlooked.

This review attempts to cover the plethora of activities that make up root systems research employing the approaches detailed above. In recent years, ‘root systems biology’ has become synonymous with the ‘top-down’ approach pioneered by Benfey, Birnbaum, Coruzzi and co-workers, involving transcriptomics and bioinformatic analysis (Poultney et al., 2007; Benfey et al., 2010; Katari et al., 2010). Here, we first describe the impressive progress made using this approach to build root GRN and elucidate novel biological mechanisms. We also discuss how mathematical and computational modeling employing ‘bottom-up,’ ‘middle-out’ and ‘multi-scale’ approaches helps us understand the mechanistic behavior of root systems from the GRN scale to whole root system and rhizosphere scales, thereby enabling us to bridge the genotype to phenotype gap by integrating environmental information.

2 *Top-down* root systems biology approaches

Until recently, root-related gene and protein networks have been characterized using low throughput molecular genetic and cell biological approaches (e.g. double mutants, FRET; Benfey et al., 2010). Although these approaches have enabled the discovery and characterization of many small GRN controlling root developmental programs, they do not work for large-scale network identification as they build GRN in a gene-by-gene manner. In contrast, ‘top-down’ systems biology approaches are ideally suited to construct large-scale root networks of genes or proteins (reviewed in Bassel et al., 2012).

New 'omic' profiling technologies such as transcriptomics have enabled explosive growth of 'top-down' approaches in recent years and the large-scale datasets required to conduct top-down approaches have increased almost exponentially since the sequencing of the *Arabidopsis* genome (*Arabidopsis* genome Initiative, 2000; see reviews by Mardis, 2008; Egan et al., 2012; Hamilton and Buell, 2012). Expression analysis of specific cell types and developmental stages in *Arabidopsis* roots can be accomplished using techniques such as laser capture micro-dissection (Kerk et al., 2003; Jiao et al., 2009). The most widely-used technique uses fluorescence-activated sorting of marked cells, usually from transgenic lines with tissue-specific expression of green fluorescent protein (GFP). This technique uses total RNA prepared from sorted protoplasts of these transgenic lines (Birnbaum et al., 2005). Another method involves expressing a tagged ribosomal protein behind a tissue-specific promoter (Zanetti et al., 2005). Immunoprecipitation of ribosomes from such transgenic plants allows researchers to profile only the actively translated transcripts.

The majority of current 'top-down' examples in roots examine mRNA transcript abundance (transcriptomics), but transcript levels do not necessarily predict protein abundance. Directly examining proteins, a protein expression map of the *Arabidopsis* root provided the identity and cell type-specific localization of nearly 2,000 proteins, and co-localization data provided support for numerous protein-protein network interactions (Petricka et al., 2012). Other 'omics' profiling techniques recently employed in roots include metabolomics, which measures the pool sizes of metabolites (molecules of $M_r \leq 1,000$ D) that collectively define the metabolome (Fiehn et al., 2000). Moussaleff et al, (2013) used a cell-sorting approach to perform non-targeted metabolomics assays on core cell types in the *Arabidopsis* root. This study provided a metabolic map of the key root cell types, identifying fifty metabolites.

The challenge root researchers face is how to use these rich 'omics' datasets to elucidate meaningful regulatory relationships between mRNAs, proteins and other biological molecules. Two distinct approaches have been employed to construct large-scale root regulatory networks (summarized in Figure 1, using two specific

examples). The **direct** approach involves a systematic experimental screen of pairwise interactions among a carefully selected subset of potentially interacting root genes and/or proteins. The second, **indirect** approach exploits the availability of root 'omics' (predominantly transcriptomic) datasets to infer GRN through statistical analysis and identification of regulatory relationships between tens to many thousands of genes. In the next section we review both approaches using recent examples.

2.1 Building root networks using interaction screens

Brady et al. (2011) recently described an interaction-based screen to unravel the GRN controlling the identity of the stele, the innermost *Arabidopsis* root tissues. The authors initially used a high-resolution spatio-temporal root gene expression map (Brady et al., 2007) to select 167 genes encoding stele-enriched transcription factors (TFs). They carried out a yeast 1-hybrid screen using these 167 TF genes as preys and the promoters of 65 out of the 167 genes as baits. They also included in their screen the promoters of 28 microRNAs (miRNAs) that target some of the selected TFs. Next, they completed their network with a yeast 2-hybrid screen to identify protein-protein interactions between the 167 TFs. The two screens generated a network of 66 protein-DNA interactions and 25 protein-protein interactions. 59% of the protein-DNA interaction could be validated *in planta* by chromatin immunoprecipitation (ChIP) and genetic analysis. To determine whether the protein-DNA interactions detected by yeast 1-hybrid and ChIP-PCR resulted in positive or negative effects on transcription, they next analysed changes in transcript accumulation of the predicted targets in lines knocked-down or over-expressing each putative regulator. This approach allowed them to quantify the strength of the interactions and identify the strongest regulators for each target. This systematic screening approach successfully recovered the core of the GRN controlling stele tissue identity and quantitatively measured the strength of the interactions in this GRN. This time-consuming method yields only a partial GRN, but it provides an excellent experimentally validated framework to undertake more detailed analysis of network topology through further lab-based and *in silico* modeling studies.

One of the key challenges to building GRN lies in defining and confirming the *cis*-acting regulatory elements in the promoters of transcription factor targets. Recent genome-aided investigations of the developmental pathway have begun to provide insights into how useful this can be (Won et al., 2009). This approach depends on knowledge of the consensus binding motifs for the transcription factors, knowledge largely lacking in *Arabidopsis*. High throughput genomic approaches such as ChIP-seq and yeast 1-hybrid have begun to fill in this knowledge gap, but the large number of known and putative transcription factors suggests that *in silico* scans with subsequent validation will remain the standard approach.

2.2 Building root networks using statistical inference algorithms

Network inference approaches typically look for correlative associations (e.g. sets of genes expressed under a particular condition) in genome-wide 'omics' datasets, commonly using transcriptomic data. Initially the approach identifies the components correlated with specific variables (e.g. time, developmental stage, genetic differences, and growth conditions). Conditional and temporal/developmental transcriptomic data can be used to infer GRN. Conditional expression data compares variables such as wild-type versus root hair defective mutants (Jones et al., 2006), treated versus mock treated (Paponov et al., 2008), or combinations of genetic and environmental variables (Okushima et al., 2005; Bruex et al., 2012). Time-series expression data can be collected following developmental or experimental induction of the system. Temporal and developmental transcriptomic datasets relevant to roots include those associated with lateral root formation (De Smet et al., 2008), root hair nodulation (Libault et al., 2010), or responses to hormone or nutrient treatments (Krouk et al., 2010; Bielach et al., 2012). For example, Ruffel et al. (2011) studied how systemic nitrate signaling affected genome-wide programming in root development (Figure 1). This involved physically isolating root systems of the same plant using a 'split-root' and challenging them with different nitrate conditions. This system was used to probe systemic nitrate responses by comparative transcriptomic analysis of *Arabidopsis* mutants impaired in nitrate reduction and hormone synthesis, and on decapitated plants. The authors performed three-way analysis of variance (ANOVA) to analyze the data as a whole, including the nitrogen and split-root effects.

This analysis identified genes whose nitrate responses were affected by the split-root conditions and found least two genetically independent systemic signaling mechanisms important for communicating the nitrate supply and demand of a plant (Ruffel et al., 2011).

Clustering provides a useful first analysis of transcriptomic datasets (Eisen et al., 1998; Bansal et al., 2007). Hierarchical clustering uses a pairwise measure of distance between gene expression profiles to compute distance trees (dendrograms) that delimit clusters of co-expressed genes; genes showing similar expression profiles are likely to be co-regulated. Clustering can help identify regulatory modules, but it does not reveal the regulatory relationships of the genes inside a cluster. If two genes cluster together, they could be targets of another gene or one gene could be directly or indirectly regulate the other.

In contrast to clustering approaches, correlation network reconstruction algorithms aim to infer the regulatory relationships between correlated genes. The simplest way of building a correlation network consists of first calculating all pairwise Pearson's correlation between expression profiles and then determining the edges of the network through a thresholding procedure. However, these simple correlation networks analyses have the same drawbacks as clustering analyses.

Several algorithms based on various types of mathematical formalisms have been developed over the last decade to infer GRN State-space representation models, one of the earliest and widely used methods for modeling gene networks (Noor et al., 2013), attempt to capture the dynamic evolution of the gene network. Krouk et al. (2010) employed a state-space modeling approach termed machine learning to analyze time series transcriptome data and generate testable hypotheses to probe potential mechanisms underlying the root nitrate signalling pathway. Out of 550 nitrogen-regulated genes, the authors extracted 67 that correspond to the predicted transcription factors plus 9 N-regulated target genes belonging to the primary nitrogen assimilation pathway. The machine learning approach proposed a regulatory network derived from the high-resolution transcriptome dataset that was sampled at rapid timepoints after roots were treated with nitrate. Intriguingly, of the 76 studied genes, 60 have 500 significant connections, suggesting that nitrate

regulation is highly complex. Indeed, the authors speculated that this high level of connectivity “may explain why, to date, (reductionist) experimental analyses have uncovered only few molecular actors specifically involved in the control of NO_3^- -induced gene expression” (Krouk et al. 2010). Nevertheless, by determining the topology of the GRN, the authors pinpointed a key node, the SBP-box-like 9 (SPL9) transcription factor. The model identified SPL9 as the third-most influential transcription factor regulating nitrogen assimilation genes, since it is induced minutes after nitrate treatment, and controls at least 6 genes, including two highly important genes. Furthermore, SPL9 had the greatest magnitude and number of ‘in’ connections controlling it. Hence, the authors concluded that: “SPL9 constitutes a potential crucial bottleneck in the flux of information mediated by the proposed nitrogen regulatory network,” a prediction that was later validated experimentally through studies demonstrating that *SPL9* overexpression modifies the characteristics of nitrate signal propagation.

Probabilistic graphical modeling techniques such as Bayesian, Gaussian, and Qualitative Probabilistic Networks have also emerged as useful tools for reverse engineering GRN (for a comprehensive review, see Noor et al., 2013). Bayesian approaches were initially developed to infer GRN from steady-state expression data (Friedman et al., 2000). They were later extended to solve inference problems on time-series expression data (Dynamic Bayesian Network, Perrin et al., 2003; Yu et al., 2004). For example, to infer the GRN important for root hair differentiation, Bruex et al. (2012) examined the transcriptomes of 17 root epidermal mutants and 2 plant hormone treatments. They used a Bayesian approach to infer the regulatory relationships between 208 core genes and used expression information from a developmental time series dataset to position genes temporally within the network (Bruex et al., 2012).

2.3 Tools and resources available for roots system biology studies

A wealth of available tools and databases can be used for data collection, data mining, analysis and meta-analysis. The high spatial resolution of transcriptome datasets has proved particularly valuable for root studies (Brady et al., 2011). For

example, 85% of the 3,656 differentially expressed genes identified by a cell-sorting analysis were not detected in whole-root experiments (Gifford et al. 2008). This illustrates the sensitivity of these approaches; also, transcriptional responses to environmental cues are likely to be tissue-specific and masked in whole-root assays.

As the capacity to generate high-throughput large-scale datasets increases, so does the need for tools designed to collect and examine this wealth of new information. Despite the focus on above-ground tissues and whole-plant expression, many tools and resources are already available for root biologists to adopt for their studies. The publicly available *Arabidopsis* eFP Browser visualizes relative and absolute gene expression data across approximately 22,000 genes from *Arabidopsis*, as represented on the ATH1 GeneChip from Affymetrix (Winter et al., 2007). This platform includes root-expressed genes profiled using 19 marker lines and 13 developmental stages, corresponding to cell types and tissues at progressive developmental stages, and transcript profiles from quiescent center cells of the root meristem (Brady et al. 2007). The updated *Arabidopsis* eFP Browser incorporates gene expression data from a variety of sources, including roots under abiotic stresses such as high salinity, iron deprivation, nitrogen availability, low pH, and sulfur and phosphate deficiency (Dinneny et al., 2008; Gifford et al., 2008; Iyer-Pascuzzi et al., 2011; Lin et al., 2011).

The majority of root transcriptome datasets contain microarray data, but many future datasets will contain next-generation sequencing data, from RNA analysis through cDNA sequencing at a massive scale (RNA-seq) (Ozsolak and Milos, 2011). Indeed, RNA-seq overcomes many of the limitations of microarray technologies. For example, RNA-seq can detect alternative isoforms of RNA transcripts and discover miRNAs (Fahlgren et al. 2007), or siRNAs (Kasschau et al. 2007; reviewed in Ghildiyal and Zamore, 2009) (Ozsolak and Milos, 2011). Next-generation sequencing-based investigation of *Arabidopsis* root system architecture in response to nitrate availability discovered a nitrate responsive miRNA and its target (Vidal et al., 2010). The authors found that nitrate induces the expression of both miRNA393 and its target, the auxin receptor *AFB3*, with a time delay, creating a negative feedback loop of regulation; this module regulates primary and lateral roots by modifying auxin responses via *AFB3* (Vidal et al., 2010). Vidal et al. (2013) have recently extended their studies to dissect regulatory networks activated by nitrate in

roots and acting downstream of AFB3. Employing the Sungear tool (see details in Table 1) they found that the NAC4 transcription factor controls a nitrate-responsive network. Intriguingly, the *nac4* mutant exhibits altered lateral root growth but normal primary root growth in response to nitrate. The authors conclude that AFB3 activates two independent pathways to control root system architecture.

The Arabidopsis Interactions Viewer searchable database includes experimental and predictive information from protein-protein interactions (Geisler-Lee et al., 2007). This freely available and curated resource (Table 1) queries a database of 70944 predicted and 28556 confirmed *Arabidopsis* interacting proteins against a protein(s) of interest. The experimentally confirmed and predicted interactions come from BIND, the Biomolecular Interaction Network Database (Willis and Hogue, 2006), from high-density *Arabidopsis* protein microarrays (Popescu et al., 2007; Popescu et al., 2009), from the *Arabidopsis* Interactome Mapping Consortium (2011), the Membrane protein INteractome Database MIND (Lalonde et al., 2010), the *Arabidopsis* G-signaling Interactome Database (Klopffleisch et al., 2011), and over 1190 other literature sources. The subcellular localization data in *Arabidopsis Interactions Viewer* comes from an older version of SUBA (which does not include predicted localization from SUBA3). Other freely available tools include user-friendly platforms designed to facilitate hypothesis formulation, by depositing information on high-throughput profiling information relating to plant tissue metabolome (Bais et al., 2010) and mineral composition (termed the ionome; Baxter et al., 2007).

Resources for mining high-throughput profiling datasets increasingly include user-friendly platforms, but the ever-increasing information contributed by low-throughput studies remains less well-annotated. A wealth of information relevant to root biology remains buried within various repositories, including the scientific literature. Text mining can retrieve these data, but as the body of information grows, the feasibility of manual annotation decreases. However, advanced text mining methodology has recently been implemented (Van Landeghem et al., 2013), including work to collate information about specific tissues or processes in a readily accessible format. For example, the more than 1,300 articles that describe processes related to root hair cells would require extensive time and effort to extract relevant information by a traditional literature review (Kwasniewski et al., 2013).

iROOTHAIR addresses this problem and provides a freely-available resource that collects, presents and shares the available information in a systematic, curated fashion (Kwasniewski et al., 2013). Features of iRootHair include four searchable sections: 'Genes', 'Processes of root hair formation', 'Root hair mutants' and 'References'. The database also integrates bioinformatic tools and can be used to identify particular genes of interest or find broader information about root hair genomics.

Bottom-up root systems biology approaches

In contrast to the phenomenological-type models used in 'top-down' studies, the hypothesis-driven 'bottom-up' approach uses predictive models that can be tested experimentally (recently reviewed by Middleton et al., 2012). This requires quantitative data on key components of the steady-state system to build mechanism-based models to predict emergent behaviors of the system. Mechanism-based models help us conceptualize biological processes and uncover gaps in our understanding. These models can be used as simulations, which assist in experiential design by enabling researchers to make well-founded and testable predictions. Models, by nature, oversimplify the system we wish to study, and should ideally be improved in an iterative loop, in association with experimental validation. The more quantitative data available for key components of a particular system, the greater the opportunity to construct meaningful models.

New mathematical and computational models probe the biological significance of identified interactions within increasingly complex GRN. *BioModels Database* serves as a reliable repository of computational models of biological processes (Li et al., 2010). The *BioModels Database* website (<http://www.ebi.ac.uk/biomodels-main/>) hosts a collection of models described in peer-reviewed scientific literature, as well as models generated automatically from pathway resources (Path2Models). These freely available models can be used or adapted to address relevant biological questions. As information on root regulatory networks increases, so does the opportunity to implement mechanistic models for bottom-up studies (reviewed by Middleton et al., 2012). Below, we discuss some examples of how these types of models have successfully addressed important biological questions related to root systems.

The molecular and biochemical reactions that underpin biological systems are often non-linear, with feed-forward and feed-backward loops that contribute to the robustness of the system (Mosconi et al., 2008). For example, the phytohormone auxin regulates cell division, elongation and differentiation in roots (Péret et al., 2009; Stahl and Simon, 2010; Sablowski, 2011). Auxin functions in part by mediating the degradation of Aux/IAA family proteins, which repress auxin response

transcription factors (ARFs). ARFs regulate various downstream genes that define the auxin responsive cellular processes described above, including Aux/IAs. Thus the pathway comprises multiple negative feedback loops. Middleton et al. (2010) developed a mathematical model of the auxin response network by employing a series of coupled ordinary differential equations. The model captures all key auxin response network components, including auxin, its receptor protein TIR1, ARFs and interacting AUX/IAA repressor proteins, together with their known regulatory relationships. The quantitative data from molecular studies on protein turnover and transcription rates enabled Middleton et al., (2010) to build a model using realistic parameters. Simulations using this model revealed the importance of the turnover rates of Aux/IAA protein and mRNA for the dynamic behavior of the model.

Model predictions that are inconsistent with the experimental data can point to the existence of unknown interactions or components missing from the network model. For example, the model described above showed discrepancies between its behavior and the experimental data (Middleton et al., 2010). In experiments, the abundance of some mRNA species can change by up to 30-fold (Abel and Theologis, 1996). However, achieving this level of fold-change in the original model developed by Middleton et al. (2010) requires an extremely large (and possibly unrealistic) increase in auxin levels. This points to the existence of additional regulatory mechanisms that might amplify the cellular response (as measured by *Aux/IAA* mRNA levels) to changes in auxin. A number of ARF-encoding genes respond to auxin (Okushima et al., 2005), which could represent a feed-forward loop that allows signal amplification.

The ARF family member MONOPTEROS (MP) up-regulates both its own expression and the expression of the gene encoding Aux/IAA family member BODENLOS (BDL) (Lau et al. 2011). Thus, the network consists of an ARF-based positive feedback loop coupled to the Aux/IAA negative feedback. The presence of a positive feedback loop in a model can lead to bistability, in which the system can rest in either of two stable states. The two stable states allows for switch-like behavior that functions as a form of signal amplification. For example, increasing levels of exogenous auxin can switch the system from a low-MP/BDL concentration steady-state to a high one (Lau et al., 2011). Bistable systems also exhibit hysteresis, where the behavior of a

system depends not only on its current condition, but also on its past environment. As auxin levels increase, *MP* expression increases and the system approaches a high-*MP*/*BDL* steady state. However, as auxin levels decrease, *MP* and *BDL* levels decrease much more slowly (i.e. the *MP* positive feedback loop is self-sustaining) and remain up-regulated for several hours (Lau et al., 2011). Moreover the system can generate switch-like responses to changes in auxin concentration, so that an *MP*-dependent developmental program can only initiate over a certain threshold. Once the switch has been activated (and the system is in a *MP* up-regulated state), the network's behavior will be robust to large changes in the level of auxin. Hence, modeling has revealed how auxin can induce all-or-nothing developmental outputs.

Another increasingly important aspect of root developmental biology is the study of cross-regulation between two or more hormones. For example, the interplay between auxin and cytokinin controls the size of the root meristem in an antagonistic manner (Ioio et al., 2008). Local accumulation of auxin promotes root cell division whereas high cytokinin concentrations promote differentiation. To study the interplay between auxin and cytokinin signaling, Muraro et al. (2011) developed an ordinary differential equation-based mathematical model of known interactions. In addition to the auxin-mediated up-regulation of *Aux/IAA* genes (notably *SHORT HYPOCOTYL 2* (*SHY2*)), auxin also up-regulates the transcription of genes encoding ARR-A inhibitors of the cytokinin response. ARR-A genes are also up-regulated by ARR-Bs in response to cytokinin. In contrast, cytokinin signaling activates an ARR-B transcription factor, which in turn induces transcription of *SHY2*. *In silico* perturbations to the model predict experimental changes in cell division observed in both the gain-of-function and the loss-of-function *shy2* mutants, suggesting that the network had accurately captured the regulatory mechanisms controlling meristem size.

3 Middle-out root systems biology approaches

The middle-out approach starts at the physical scale at which there is the most information, and takes bottom-up and top-down approaches. Models constructed at the cell scale are particularly well suited to the middle-out approach

because the cell comprises subcellular network systems (revealed using a top-down approach), and cell-cell interactions lead to the emergence of tissue-scale properties (revealed using a bottom-up approach). However, the middle-out approach can start at any scale with sufficient available information. In the following section we review examples where root researchers have developed models focused either at sub-cellular, cell, tissues, organ, whole plant or plant-environment scales.

3.1 Sub-cellular scale systems models

At the subcellular scale, root models have been described that simulate gene regulatory, signaling, or metabolic networks. These typically employ Boolean, ordinary differential equations, or stochastic approaches (when low copy numbers are important; reviewed by Band et al., 2012a).

Dupeaux et al. (2011) described a stochastic model developed to study abscisic acid (ABA) signal perception within a cell. The key hormone ABA regulates root growth and responses to biotic and abiotic stresses. Within the cell, ABA binds to a family of intracellular receptors, termed ABA binding to pyrabactin resistance (PYR)/PYR1-like (PYL)/Regulatory Component of Abscisic acid Receptor (RCAR) proteins. ABA binding promotes the formation of complexes involving the PYR/PYL/RCAR receptors and several types of protein phosphatase type 2C (PP2C) proteins that activate ABA responses. PYR/PYL/RCAR family members bind ABA as a monomer or dimer. Dupeaux et al. (2011) developed a computational model capturing this relatively simple network of interactions to probe the response to ABA when both monomeric and dimeric PYR/PYL/RCAR receptors compete for ABA and P2C molecules. Intriguingly, the study revealed that monomeric receptors have a competitive advantage for binding, particularly at lower ABA concentrations. Hence, tissue receptor composition and oligomerization likely impact ABA responsiveness.

Cruz-Ramírez et al. (2012) recently used a mathematical model of a small GRN to demonstrate how two nested feed-forward loops precisely control asymmetric cell divisions within the root stem cell niche (Figure 2). The authors employed modeling

to help understand how spatial restriction of these divisions requires physical binding of the stem cell regulator SCARECROW (SCR) by the RETINOBLASTOMA-RELATED (RBR) protein. They generated coupled ordinary differential equations to describe the wiring of the network. Initial models tested the importance of the CycD6 feedback loop on phosphorylation of RBR and its interaction with SCR, and subsequent feedback on *SCR* expression. Because asymmetric cell divisions occur at low or high SCR-RBR levels, the authors analyzed equilibrium levels of nuclear Short Root (SHR) and free SCR (unbound to RBR). SHR and auxin distribution and their effect on the precise positioning of CycD6 in the cortex endodermis determine radial and longitudinal positional information. Coupling of cell-cycle progression to protein degradation resets the circuit, resulting in a “flip-flop” (i.e. a circuit with two stable states) that constrains asymmetric cell division to the stem cell region. RBR binds SCR, and together with the RBR regulator CYCD6;1, defines the position of asymmetric cell divisions. The resulting network creates a robust, bi-stable switch combining the auxin gradient along the longitudinal axis with the SHR distribution pattern along the radial axis (Cruz-Ramírez et al., 2012).

3.2 Cell scale systems models

At the cell level, models capture interactions between neighboring root cells. For example, Savage et al. (2008) modeled patterning of the *Arabidopsis* root epidermis, which forms alternating files of trichoblasts (bearing root hairs) and atrichoblasts (lacking root hairs). Epidermal cells overlying two cortical cell files form trichoblasts. The mobile activator proteins GALABRA3/ENHANCER OF GALABRA3 (GL3/EGL3), regulate this patterning and form an activator complex with TRANSPARENT TESTA GALABRA1 (TTG1), WEREWOLF (WER), and the mobile inhibitor protein CAPRICE (CPC). Savage et al. (2008) formulated a stochastic Boolean model, embedded into a circle of cells, with the levels of CPC and GL3/EGL3 depending on the rate of transcription in the neighboring cells. They modeled potential network structures, comparing WER self-activation and CPC inhibition via competition for GL3/EGL3 with constitutive WER transcription and CPC directly inhibiting WER. Whilst both models replicated the patterning in wild-type roots, the latter model best explained the mutant phenotypes, in particular *cpc*.

Benítez et al. (2008) explored the same problem using a three-state deterministic model within a two-dimensional array of cells; they also found that WER auto-regulation was not required for pattern formation in wild-type roots, but it proved necessary to explain some mutant phenotypes in the two-dimensional geometry. The two-dimensional model was also used to explore the effects of cell shape, with cell elongation acting to stabilize patterns. Whilst experiments of Savage et al. (2008) provided evidence that WER auto-regulation is not important for the early patterning of epidermal cells, local self-activation does occur at later stages (Kwak and Schiefelbein, 2008; Kang et al., 2009), although its functional role remains unclear.

3.3 Tissue to organ scale systems models

Tissue and organ scale models describe growth, transport, and deformation on the macroscale, usually employing continuum (mainly partial differential) equations that do not explicitly subdivide a tissue into discrete cells. Péret et al. (2012) described a tissue-scale modeling approach to investigate how aquaporins regulate lateral root emergence. In *Arabidopsis*, lateral root primordia originate from pericycle cells located deep within the parental root and emerge through endodermal, cortical and epidermal tissues. Recent studies have highlighted the importance of the localized auxin signal originating from the tip of the lateral root primordia to induce specific physiological responses in overlying tissues (Swarup et al., 2008; Péret et al., 2012). Biomechanical properties targeted by auxin include changes in aquaporin spatial expression patterns to modify water fluxes. Based on these data, Péret et al. (2012) included a mathematical model describing how aquaporin-dependent tissue hydraulics could affect the timing of lateral root development. In the model, water fluxes are coupled to primordium expansion, with the lateral root primordia and overlying tissue represented as two fluid compartments. The dividing primordium cells have a prescribed increasing osmotic potential. The model predicts the resulting water fluxes and pressure dynamics and shows how these drive primordium expansion. The model predicts that aquaporin repression by auxin promotes lateral root emergence, and explains the experimentally observed delay in lateral root primordia emergence for both aquaporin loss-of-function and overexpression lines.

3.4 Plant scale systems models

Plant-scale models aim to integrate the behavior of individual plant organs, or even sub-organ characteristics, at the whole plant level, into functional structural plant models (Godin and Sinoquet, 2005). At this level, major dynamic processes come into play, including resource capture (carbon, water, nutrients), long distance transport, resource allocation, long distance signaling, and concurrent growth of many organs. This type of model implements most processes, such as the production and consumption of carbohydrate assimilates (Drouet and Pagès, 2007)

or the development of the plant (growth, branching), at the organ or sub-organ level. These processes are integrated at the plant level by connecting the different parts (up to several thousand) according to plant topology. These models often adopt a tree-like structure, well suited to representing plant networks.

The analysis of carbon transport and allocation within the root system has been described using the MassFlowDyn model (Bidel et al., 2000), which explicitly simulates phloem transport of sugars following simple source/sinks rules, in relation to the growth and development of complex root system. The authors demonstrated the importance of the initial diameter of individual root meristems as a determinant of sink strength. Formation of a hierarchy of root meristems (primary > lateral) leads to herringbone type systems (akin to the arrangement of bones in a fish), while similar meristem sizes lead to fibrous root systems (composed of many roots of approximately equivalent lengths).

3.5 Plant-environment scale systems models

A second category of functional structural plant models explicitly takes into account the bidirectional flow of material or energy and the biological, physical and chemical interactions between the plant and the soil compartment. Compared to the previous models, which generally exclude environmental effects or consider them uniform, these models explicitly include the environment, with a focus on the processes of interest. As for plant scale models, the different processes-- in the plant, in the environment or at the plant/environment interface-- are implemented at the local scale and integrated at the whole plant scale. The plant structure still follows a tree-like structure, but the environment, especially the soil, may require a three-dimensional graph, as all neighboring soil voxels are inter-connected. The integration of the plant tree and environment graph requires a complex modeling framework in which the positioning of the different structures can be matched in time and space. Indeed, since most of the processes are computed locally, the exact position of every plant and environment element is needed.

Soil-root models developed in the last two decades address mostly water and nutrient dynamics. SimRoot (Lynch et al., 1997; Postma and Lynch, 2011a) and

ROOTMAP (Diggle, 1988; Dunbabin et al., 2002) link root growth and root system architecture with nutrient acquisition (reviewed by Dunbabin et al., 2013). SimRoot, for example, was used to demonstrate the influence of root type-gravitropism on phosphorus acquisition in bean (Ge et al., 2000). Other models have been developed to analyse the dynamics of water flow in the soil root-system, taking into account the long distance movement of water in the soil (Doussan et al., 1998; Javaux et al., 2008; Schröder et al., 2008; Lobet and Draye, 2013) (Figure 2C). The three-dimensional model R-SWMS, for example, simulates the uptake of water by individual root segments as a function of the root hydraulic architecture and soil properties (Doussan et al., 2006; Draye et al., 2010) and will help our understanding of the contribution of long distance regulatory signals (e.g. ABA) to whole plant hydraulics and water use.

The recent development of methods for imaging live roots in real soil at the microscopic scale (Mooney et al, 2012) enables a step change in the modeling of nutrient uptake. The ability to uncover the three-dimensional interactions between the root surface and soil porosity has allowed the development of a model of phosphate uptake by root hairs based directly on the real geometry of hairs and associated soil pores (Keyes et al, 2013). The new model questioned the currently accepted relative contribution of the root surface versus root hairs during the uptake of phosphorus.

The interface between the root and its environment also has many biological interactions between the root system and its surrounding communities. The German plant physiologist Lorenz Hiltner (1904) coined the term “rhizosphere” to describe the region inhabited by a unique population of microorganisms, which he postulated were influenced by the chemicals released from plant roots (Hartmann et al., 2008). Root exudation represents a significant carbon cost to the plant (Lynch and Whipps, 1990). In addition, the presence of exudates and microbes profoundly affects physical properties of the soil-root interface such as its hydraulic conductivity (Carminati and Vetterlein, 2013). Therefore, understanding plant behavior in the field requires a fundamental understanding of the rhizosphere. The functional structural plant models framework has all the required components to address the spatio-temporal dynamics of the rhizosphere, yet such work remains to be initiated.

3.6 Crop scale systems models

At the crop scale, the description of plants can be further simplified, mainly to reduce computational cost as the total number of plants (and subsequently of individual plant elements) increases. Compared to the plant-environment scale, modeling at the crop scale still requires the space and time localization of the plants, but to a lesser extent; for example, the position of every root is not required anymore. Such models mainly aim to predict the construction of final crop yield under different environmental conditions, in different agronomic management practices, or in different genetic backgrounds.

Despite the fact that plants are conceptualized at a higher level than in lower scale models, some crop models includes physiological and genetic modules influencing crop development. For example, SUNFLO (Casadebaig et al., 2011) models sunflower adaptation to environmental stimuli (water and nitrogen stress) and aims to assess genotypes for their success in these changing environments, as evaluated by their yield. The SUNFLO model successfully predicted genotype when tested against validation data from over 50 field experiments. This could be due to incorporation into the model of genotype via gene relationship to dynamic processes such as nutrient uptake and oil content, rather than the model's inherent predictive power. These results also highlight the need to incorporate information about rooting phenotypes to enhance the model's applicability in water stress situations.

In development for more than 15 years, the crop-level model Agricultural Production Systems sIMulator (APSIM) aims to integrate genetic information, management practices and environmental conditions to predict yields of economically important crop plants (Keating et al., 2003). APSIM has been used to test the effect of genetic variation, either at the root (Hammer et al., 2010; Manschadi et al., 2010) or the shoot level (Chenu et al., 2008). Using APSIM, Chenu et al. (2009) modeled the effect of organ-level quantitative trait loci (QTLs) on maize yield. Using the model, they simulated the effect of QTLs of two selected developmental processes (leaf and silk elongation) by generating hundreds of hypothetical recombinant inbred lines and testing them across several patterns of drought seasons. The simulations predicted the QTLx E interactions observed under field

conditions. In particular, yield increase depended on higher leaf elongation rate in well-watered conditions, but depended on decreased anthesis to silking interval in all environments. Additionally, the pleiotropic effects of the different QTLs limited possible trait combinations. Using APSIM, Manschadi et al. (2010) evaluated the effect on final yield of specific root traits. By reducing the insertion angle of primary roots in wheat, the authors demonstrated the capacity of the crop as a whole to extract more water and increase yield. Hence, integration of local processes (root emergence, leaf growth rate) at the crop scale can be useful to help understand the crop system as a whole.

Crop models initially aimed to simulate yield construction over the season based on resource capture and environmental conditions. They do not obviously lend themselves to organ-based representation of the plant, because the elementary unit is the crop itself. Rather, parameters such as mass, leaf area index and root depth represent aspects of plant architecture. These variables are essential to compute light interception or soil resource capture. Although crop models are much less mechanistic than the previous examples discussed, reasons to include them into the root systems biology approach include: (1) Crop models represent one further step in up-scaling, and, in essence, that step resembles the step from cell to organ, or from organ to plant. Each step requires the identification of the most relevant source of variation between elementary units. (2) These models generate unique paths in time that integrate environmental, management and genetic factors that influence the future of the crop. They have been used to simulate the emergence of genotype by environment (GxE) interactions, which remain one of the major challenges to moving from the lab to the field. They, therefore, provide very useful insights to orient research at lower scales. (3) They offer the only simulation framework that allows testing (systematically) of many (virtual) genotypes in many seasons, which has become a major step in plant breeding, and provide a means to address the agronomic potential of gene targets identified in basic research.

4 *Multi-scale* root systems biology approaches

Multi-scale models consider behaviors on two or more scales, ranging from the subcellular up to whole-organism scales and beyond (see Figure 3). They must also

incorporate different temporal scales, ranging from minutes for transcription to weeks for developmental adaptation. Relatively few examples of multi-scale models have been described to date for roots (or other plant organs). We selected the following examples to illustrate the potential for developing multi-scale models to probe the mechanisms underlying complex, non-linear biological processes in roots. One key to predicting the relationship between genotype and phenotype is linking the spatio-temporal concentrations of regulators, such as auxin, with cell growth and division. We currently lack a mechanistic understanding of the effect of auxin on growth, but several existing examples employ multi-scale approaches, incorporating phenomenological models to describe this missing step.

Grieneisen et al. (2007) proposed that a gradient of auxin controls the specification of the root's developmental zones, supposing that cell division occurs at high hormone concentrations, and cell elongation at lower concentrations. Due to the predicted auxin distribution, the model simulates cell division and growth dynamics in the meristem and elongation zone, capturing the gradual expansion of the meristem over the first 8 days post germination and the reduction in meristem size after root excision. Chavarria-Krauser et al. (2005) modeled the growth dynamics at the root tip, supposing that the ratio between auxin and cytokinin governs the production and degradation of a remodeling enzyme, which in turn regulates cell growth and division. For root developmental responses, Lucas et al. (2008), in predicting lateral root initiation, considered a pool of auxin at the root tip that is consumed by root gravitropic bending and/or emergence. Using such a phenomenological model between auxin and root development, the authors accurately predicted experimentally observed perturbations in lateral root initiation for the *aux1* root gravitropic and *lax3* lateral root emergence mutants.

The hormone gibberellin controls root growth by, for example, controlling cell growth within the elongation zone (Ubeda-Tomás et al., 2008). Band et al. (2012b) recently developed a multi-scale model of gibberellin dynamics in the *Arabidopsis* root elongation zone, prescribing cell growth (using experimental measurements) and simulating gibberellin dilution, diffusion of gibberellin between cellular compartments and the response network through which gibberellin degrades the growth-repressing DELLA proteins (describing the dynamics with a system of

ordinary differential equations). The model (see Figure 3) revealed that, as cells pass through the elongation zone, dilution (rather than degradation) creates a decline in gibberellin concentration, leading to spatial gradients in the levels of downstream DELLA mRNAs and proteins. This model prediction was confirmed experimentally with knock out mutants lacking every root-expressed GA2ox degradation enzyme. The study also considered the dynamics in plants treated with paclobutrazol (an inhibitor of gibberellin biosynthesis) and plants with mutations in the gibberellin biosynthesis and signaling pathways; these cases revealed that the growth rates appear to reflect the fold-change in DELLA as cells traverse the elongation zone. Furthermore, the model provided new insights into the normal phenotype exhibited in the *ga1-3/gai-t6/rga-24* triple mutant. The model demonstrated that the effect of the *ga1-3* mutation in reducing gibberellin biosynthesis (leading to higher functional DELLA levels) counteracts the effect of the *gai-t6/rga-24* mutation in reducing the translation of functional DELLA; if these two processes are suitably balanced, the levels of functional DELLA are similar to wild-type, explaining why the triple mutant exhibits normal cell elongation. In summary, by assimilating a range of data and knowledge, the model revealed the dominant effect of gibberellin dilution (rather than degradation) on the emergent DELLA distribution, providing new insights into gibberellin's growth regulation.

Jeuffroy et al. (2012) also used multi-scale modeling to identify genotypes with desirable characteristics for breeding programs. The pea crop model AFISOL for yield characteristics (incorporating water and nitrogen flow) was linked to SISOL modeling soil parameters (soil moisture content, soil biota activity effects, impacts of machinery and natural soil weathering) and OTELO for farm practices (fertilizer, pesticides and soil amendments). These models all include dynamic parameters and this work demonstrates the capability to link several models of different scale, revealing the applicability of crop models in the systems biology approach across multiple plant physical and temporal scales. Additionally, the model assessment included real and virtual genotypes, which shows the value for isolating desirable characteristics that could then be harnessed through more molecular techniques.

5 Future challenges for root systems biology

Major advances have recently been made employing 'top-down' systems approaches to identify the key molecular players (genes, RNAs, proteins, etc.) and to elucidate several GRN that control root growth and development (Benfey et al., 2010; Ruffel et al., 2010; Bassel et al., 2012). However, regulatory networks not only operate at the molecular-scale, but must also integrate information from cell to organ to rhizosphere scales, to probe the mechanisms controlling root development. Mathematical and computational models provide invaluable tools to bridge these distinct spatial and temporal scales and generate new mechanistic insights about the regulation of root growth and development (Band et al, 2012a). Nevertheless, we are only at the beginning of this new research area and several challenging issues remain to be addressed for this field to move forward.

Integrating models

In general, the models highlighted in this review have been designed independently of one another, raising important issues relating to interoperability. To proceed further, models must start to be assembled into unified frameworks. A number of conceptual and technological reasons make model integration challenging. First, models may operate at different spatial resolutions or dimensions. For example, coupling a two-dimensional mechanical model of root tissues at a subcellular resolution with a two-dimensional model of auxin transport designed at cellular resolution would require homogenizing the spatial resolution such that the mechanical model updates cell geometries over time. Second, models may also operate at different temporal resolutions. For example, mechanical processes are often assumed to be much faster than biochemical processes. Hence, cell size and shape can be considered constant in computing chemical simulations. Third, combining different types of mathematical models (e.g. Boolean, stochastic, ordinary differential equations, and partial differential equations) can often prove challenging. Nevertheless, models integrating mechanistic and stochastic models have been successfully designed at the organism level for aerial tissues (Costes et al., 2008).

Creating combinations of models often represents a large amount of theoretical and experimental effort. Hence, it is important that these models are freely available to

the community as shared data sets, open-source model formats such as SBML (Hucka et al., 2003) or CellML (Cuellar et al., 2003), or common modeling software platforms, such as VV (Prusinkiewicz, 2004), OpenAlea (Pradal et al., 2008) or MorphoGraphX (Kierzkowski et al., 2012). The move toward model integration and flexibility in the crop modeling community should be translatable to the root system biology ethos. For example, RECORD (Renovation and COORDination of agroecosystem modeling) represents a concerted effort by French researchers at INRA to develop an “integrated modeling platform” that could eventually include Geographic Information System data, open source statistical resources and data handling resources (Bergez et al., 2012). Additionally, planned as a repository for crop models, the RECORD platform includes many principles of root system biology including multiple time scales (seconds to months), spatial variation and environmental dynamics. RECORD aimed to use as many existing models as possible. However, to foster the incorporation of sub-crop scale models as advocated in root systems biology, RECORD would need to look beyond crop-soil interactions to the root system itself.

Towards digital plants and populations

Developing a mechanistic model of a whole plant represents a logical next step. Indeed, given the exchange of water, nutrients and signals between root and shoot organs, developing a virtual root or shoot model independent of each other could be considered naïve in the longer term. To date, models of diverse root system sub-processes have been developed at different scales. Compared to initial approaches in systems biology, most of these models make explicit use of spatial information. Such spatial information represents different aspects of realistic root structures and can take the form of a continuous medium, a branching structure of connected elements (e.g. root meristems), a multicellular population or a set of interacting subcellular compartments. By progressively integrating more functional aspects into these realistic representations, researchers have created new models (Sievänen et al., 2000; Godin and Sinoquet, 2005). Functional structural plant models provide a promising platform with which to create a digital plant model.

Compared to many previous models developed on aerial parts (e.g. Vos et al., 2010) and on root systems (Bidel et al., 2000; Pagès et al., 2004), recent functional structural plan models also integrate gene regulation and signaling as a new dimension in the analysis of development (Prusinkiewicz et al., 2007; Han et al., 2010). Through the combined modeling of genetic networks, physiological processes and spatial interaction between components, development of a new generation of functional structural plant models opens the way to building digital versions of real plants (Coen et al., 2004; Cui et al., 2010) and testing biological hypotheses *in silico* (Stoma et al., 2008; Perrine-Walker et al., 2010).

Ultimately, breeders measure plant performance at a population scale, rather than an individual scale. Hence, mechanistic multi-scale models need to be developed that bridge the remaining physical scale between the plant and the field to relate genotype to phenotype and enable engineering of crop traits. Relatively few examples exist where modeling results have been used to direct the selection of new crop varieties. One promising example is the optimization of bean root systems for phosphorus (P) and water acquisition. Using the model SimRoot, Lynch and co-workers investigated the impact of root gravitropism (Ge et al., 2000), root hair production (Ma et al., 2001) and aerenchyma production (Postma and Lynch, 2011, 2011) on plant performance under various phosphorus and/or water conditions. These results defined root system ideotypes (Lynch and Brown, 2001; Lynch, 2011; Lynch, 2013) and were validated by several experimental trials (e.g. Bates and Lynch, 2001; Rubio et al., 2003). This identified QTLs for basal root gravitropism (Liao et al., 2004) and root hairs (Yan et al., 2004) and led to the selection of new bean varieties, now used by breeding programs in South America and Africa. This example illustrates how modeling at the plant level, can drive research and untangle processes at the molecular level, and inform crop breeding.

Integrating environmental information

Phenotype represents the output of the interaction between genotype and environment. In this review, almost all of the models described study intrinsic root regulatory processes. This provides a solid foundation for describing basic developmental mechanisms, but to account for root plasticity, models must also

integrate environmental information. Developing mixed genetic-ecophysiological models that bridge the gap between genetic and environmental regulation represents an important goal (Roose and Schnepf, 2008; Draye et al., 2010). These environmental factors include: soil physical properties; water, nutrient and macro/micro element availability and distribution; mycorrhization and nodulation; and competition and interaction with other root systems. Obtaining such spatial information remains very challenging. Researchers traditionally assess soil structure by imaging and then physically removing successive layers of soil, and examine root structure (of soil-grown roots) by root washing. As both are destructive, construction of combined root and soil data sets requires integration of measurements obtained from different samples. High resolution synchrotron and X-ray micro CT imaging has the potential to provide rich data sets from single samples, with concurrent measurements of root and soil (Mooney et al., 2012; Keyes et al., 2013).

Root systems biology also urgently requires new methods to assay rhizosphere parameters in addition to root and soil. For example, having tools to dynamically monitor quantitative changes in root biology (hormones, water status, nutrients, etc) and root environment (pH, nutrient content) would address a major challenge. The development of novel biosensors based on new understanding of the pathways responsible for the perception of small signalling molecules and advances in imaging technologies and mathematical modelling, will enable a truly quantitative analysis of biological processes (Band et al, 2012b; reviewed by Wells et al, 2013). Combined with new dynamic sensors for environmental parameters like optodes (e.g. Elberling et al. 2011), new biosensors promise to provide a deeper understanding of how plant systems interact with their environment. Moreover, these sensors could result in innovative technologies to monitor biotic or abiotic stresses in the field (Chaerle et al., 2009). Biosensors could help optimize the use of inputs such as water, fertilisers or pesticides and therefore achieve more environmentally-friendly and sustainable agricultural practices. Such tools and the information generated will also greatly aid the development of more realistic root-rhizosphere models and, ultimately, help optimize crop root architectures for soil types and nutrient regimens.

Acknowledgments:

We would like to thank Rodrigo Gutierrez and Miriam Gifford for sharing information prior to publication. We thank the Biotechnology and Biological Sciences Research Council (BBSRC) and Engineering and Physical Sciences Research Council (EPSRC) funding for the Centre for Plant Integrative Biology (CPIB) to SM and MJB. In addition, we gratefully acknowledge the support of the Royal Society and Wolfson Foundation (MJB); ERC 'FUTUREROOTS' (MJB and SM) and BBSRC Professorial Research Fellowship funding (MJB); Communauté française de Belgique (ARC 0510-329 grant to X.D.) and the Belgian Science Policy (Interuniversity Attraction Poles Program P7/29 grant to S.P., G.L., X.D. and M.J.B.) for financial support.

Figure legends

Figure 1. Examples of top-down approaches to infer gene regulatory networks.

(far left column): Krouk et al., 2010 combined time-series transcriptomic data with published hormone-treatment datasets (Nemhauser et al 2006) to obtain a temporal model for nitrate-driven gene expression networks. They then used a machine learning approach (state-space modeling) to infer a regulatory network governing nitrate response over a time.

(middle, left column): Ruffel et al., 2011 combined genetic and genomic approaches within a 'split-root' system to study how systemic nitrate signaling affected genome-wide programming in root development.

(right column): Brady et al., 2011 employed a modeling approach to predict the regulatory potential of each of the relationships they identified within the stele GRN. Stele-expressed transcription factors were selected from the GRN dataset published previously (Brady et al., 2007). Protein-protein and protein DNA maps were generated from yeast 1-hybrid and 2-hybrid screens of stele-expressed TFs and miRNAs. The key interactions between TFs and their target genes were measured by qRT-PCR of target gene expression in wild type and mutant lines. Regulatory interaction strength is represented by edge and arrow width with thicker lines or arrowheads representing steeper slopes. P-value strength is represented by edge opacity with darker edges representing more significant interactions. Slopes were determined by plotting quantitative PCR (qPCR) expression values of the TF and its target and fitting a line using weighted least squares regression. This predictive framework enables the identification of the most influential upstream activators or repressors to be manipulated to regulate expression of a target.

Figure 2. Examples of hypothesis driven bottom-up and middle-out approaches.

Models were constructed for processes at different spatial and temporal scales. (A) At the sub-cellular level, gene regulatory networks can be modeled to include the input of signals such as auxin and output in the form of cell-scale events, such as mitosis. (B) Stem cell division and the bi-stable switch that regulates it are modeled using longitudinal gradients of auxin and radial gradients of the transcription factor SHORTROOT (SHR) (Cruz-Ramirez et al., 2012). (C) Root system scale models,

such as RootTyp (Pagés et al., 2004) can be linked to hydraulic models to explicitly simulate water flow in the soil-plant domain (Javaux et al., 2008).

Figure 3: Root biological models can be multi-scale.

Band et al. (2012b) demonstrate the power of multi-scale modeling for understanding gibberellin dynamics during root elongation. The elongation zone is modeled as a collection of cells (organ scale) (A) and used to motivate modeling of cell expansion and dilution in the vacuole (cell scale) (B). This naturally leads to the link between cell partitioning of gibberellin and sub-cellular processes, revealing the influence of dilution of gibberellin concentrations on DELLA expression and growth.

References

- Abel S, Theologis A** (1996) Early genes and auxin action. *Plant Physiol* **111**: 9-17
- Bais P, Moon SM, He K, Leitao R, Dreher K, Walk T, Sucaet Y, Barkan L, Wohlgemuth G, Roth MR, Wurtele ES, Dixon P, Fiehn O, Lange BM, Shulaev V, Sumner LW, Welti R, Nikolau BJ, Rhee SY, Dickerson JA** (2010) PlantMetabolomics.org: A Web Portal for Plant Metabolomics Experiments. *Plant Physiol* **152**: 1807-1816
- Band LR, Fozard JA, Godin C, Jensen OE, Pridmore T, Bennett MJ, King JR** (2012a) Multiscale Systems Analysis of Root Growth and Development: Modeling Beyond the Network and Cellular Scales. *Plant Cell* **24**: 3892-3906
- Band LR, Wells DM, Larrieu A, Sun J, Middleton AM, French AP, Brunoud G, Sato EM, Wilson MH, Peret B, Oliva M, Swarup R, Sairanen I, Parry G, Ljung K, Beeckman T, Garibaldi JM, Estelle M, Owen MR, Vissenberg K, Hodgman TC, Pridmore TP, King JR, Vernoux T, Bennett MJ** (2012b) Root gravitropism is regulated by a transient lateral auxin gradient controlled by a tipping-point mechanism. *Proc Natl Acad Sci* **109**: 4668-4673
- Bansal M, Belcastro V, Ambesi-Impiombato A, Di Bernardo D** (2007) How to infer gene networks from expression profiles. *Mol Syst Biol* **3**: 122
- Bassel GW, Gaudinier A, Brady SM, Hennig L, Rhee SY, De Smet I** (2012) Systems analysis of plant functional, transcriptional, physical interaction, and metabolic networks. *Plant Cell* **24**: 3859-3875
- Bates T, Lynch J** (2001) Root hairs confer a competitive advantage under low phosphorus availability. *Plant and Soil* **236**: 243-250
- Baxter I, Ouzzani M, Orcun S, Kennedy B, Jandhyala SS, Salt DE** (2007) Purdue Ionomics Information Management System. An Integrated Functional Genomics Platform. *Plant Physiol* **143**: 600-611
- Benfey PN, Bennett M, Schiefelbein J** (2010) Getting to the root of plant biology: impact of the Arabidopsis genome sequence on root research. *Plant J* **61**: 992-1000
- Benítez M, Espinosa-Soto C, Padilla-Longoria P, Alvarez-Buylla ER** (2008) Interlinked nonlinear subnetworks underlie the formation of robust cellular patterns in Arabidopsis epidermis: a dynamic spatial model. *BMC Syst Biol* **2**: 98
- Bergez J-E, Chabrier P, Gary C, Jeuffroy M, Makowski D, Quesnel G, Ramat E, Raynal H, Rouse N, Wallach D** (2012) An open platform to build, evaluate and simulate integrated models of farming and agro-ecosystems. *Environmental Modelling & Software* **39**: 39-49
- Bidel LP, Pagès L, Rivièrè LM, Pelloux G, Lorendeau JY** (2000) MassFlowDyn I: A Carbon Transport and Partitioning Model for Root System Architecture. *Ann of Botany* **85**: 869-886
- Bielach A, Podlesakova K, Marhavy P, Duclercq J, Cuesta C, Muller B, Grunewald W, Tarkowski P, Benkova E** (2012) Spatiotemporal regulation of lateral root organogenesis in Arabidopsis by cytokinin. *Plant Cell* **24**: 3967-3981
- Birnbaum K, Shasha DE, Wang JY, Jung JW, Lambert GM, Galbraith DW, Benfey PN** (2003) A gene expression map of the Arabidopsis root. *Science* **302**: 1956-1960

- Birnbaum K, Jung JW, Wang JY, Lambert GM, Hirst JA, Galbraith DW, Benfey PN** (2005) Cell type-specific expression profiling in plants via cell sorting of protoplasts from fluorescent reporter lines. *Nat Methods* **2**: 615-619
- Brady SM, Orlando DA, Lee J-Y, Wang JY, Koch J, Dinneny JR, Mace D, Ohler U, Benfey PN** (2007) A high-resolution root spatiotemporal map reveals dominant expression patterns. *Science* **318**: 801-806
- Brady SM, Zhang L, Megraw M, Martinez NJ, Jiang E, Yi CS, Liu W, Zeng A, Taylor-Teeple M, Kim D** (2011) A stele-enriched gene regulatory network in the *Arabidopsis* root. *Mol Syst Biol* **7**: 456
- Bruex A, Kainkaryam RM, Wieckowski Y, Kang YH, Bernhardt C, Xia Y, Zheng X, Wang JY, Lee MM, Benfey P** (2012) A gene regulatory network for root epidermis cell differentiation in *Arabidopsis*. *PLoS genetics* **8**: e1002446
- Bruggeman FJ, Westerhoff HV** (2007) The nature of systems biology. *Trends in Microbiol* **15**: 45-50
- Carminati A, Vetterlein D** (2013) Plasticity of rhizosphere hydraulic properties as a key for efficient utilization of scarce resources. *Ann Bot* **112**: 277-290
- Casadebaig P, Guillioni L, Lecoeur J, Christophe A, Champolivier L, Debaeke P** (2011) SUNFLO, a model to simulate genotype-specific performance of the sunflower crop in contrasting environments. *Agricultural and forest meteorology* **151**: 163-178
- Chaerle L, Lenk S, Leinonen I, Jones HG, Van Der Straeten D, Buschmann C.** (2009) Multi-sensor plant imaging: towards the development of a stress-catalogue. *Biotechnology Journal* **4**, 1152–1167
- Chavarría-Krauser A, Jäger W, Schurr U** (2005) Primary root growth: a biophysical model of auxin-related control. *Funct Plant Biol* **32**: 849–862.
- Chenu K, Chapman SC, Hammer GL, Mclean G, Salah HBH, Tardieu F** (2008) Short-term responses of leaf growth rate to water deficit scale up to whole-plant and crop levels: an integrated modelling approach in maize. *Plant, Cell & Environment* **31**: 378-391
- Chenu K, Chapman SC, Tardieu Fo, McLean G, Welcker C, Hammer GL** (2009) Simulating the yield impacts of organ-level quantitative trait loci associated with drought response in maize: a "gene-to-phenotype" modeling approach. *Genetics* **183**: 1507-1523
- Coen E, Rolland-Lagan A-G, Matthews M, Bangham JA, Prusinkiewicz P** (2004) The genetics of geometry. *Proc Natl Acad Sci* **101**: 4728-4735
- Costes E, Smith C, Renton M, Guédon Y, Prusinkiewicz P, Godin C** (2008) MAppleT: simulation of apple tree development using mixed stochastic and biomechanical models. *Functional Plant Biology* **35**: 936-950
- Cruz-Ramírez A, Díaz-Triviño S, Blilou I, Grieneisen VnA, Sozzani R, Zamioudis C, Miskolczi PI, Nieuwland J, Benjamins R, Dhonukshe P** (2012) A bistable circuit involving SCARECROW-RETINOBLASTOMA integrates cues to inform asymmetric stem cell division. *Cell* **150**: 1002-1015
- Cuellar AA, Lloyd CM, Nielsen PF, Bullivant DP, Nickerson DP, Hunter PJ** (2003) An Overview of CellML 1.1, a Biological Model Description Language. *SIMULATION* **79**: 740-747
- Cui M-L, Copsey L, Green AA, Bangham JA, Coen E** (2010) Quantitative Control of Organ Shape by Combinatorial Gene Activity. *PLoS Biol* **8**: e1000538
- De Smet I, Vassileva V, De Rybel B, Levesque MP, Grunewald W, Van Damme D, Van Noorden G, Naudts M, Van Isterdael G, De Clercq R, Wang JY, Meuli N, Vanneste S, Friml J, Hilson P, Jurgens G, Ingram GC, Inze D,**

- Benfey PN, Beeckman T** (2008) Receptor-like kinase ACR4 restricts formative cell divisions in the Arabidopsis root. *Science* **322**: 594-597
- Diggle A** (1988) ROOTMAP—a model in three-dimensional coordinates of the growth and structure of fibrous root systems. *Plant and Soil* **105**: 169-178
- Dinneny JR, Long TA, Wang JY, Jung JW, Mace D, Pointer S, Barron C, Brady SM, Schiefelbein J, Benfey PN** (2008) Cell identity mediates the response of Arabidopsis roots to abiotic stress. *Science* **320**: 942-945
- Doussan C, Pagès L, Vercambre G** (1998) Modelling of the hydraulic architecture of root systems: an integrated approach to water absorption-model description. *Ann Bot* **81**: 213-223
- Doussan C, Pierret A, Garrigues E, Pagès L** (2006) Water uptake by plant roots: II - Modelling of water transfer in the soil root-system with explicit account of flow within the root system - Comparison with experiments. *Plant and Soil* **283**: 99-117
- Draye X, Kim Y, Lobet G, Javaux M** (2010) Model-assisted integration of physiological and environmental constraints affecting the dynamic and spatial patterns of root water uptake from soils. *J Exp Bot* **61**: 2145-2155
- Drouet J-L, Pagès L** (2007) GRAAL-CN: A model of GRowth, Architecture and ALlocation for Carbon and Nitrogen dynamics within whole plants formalised at the organ level. *Ecological Modelling* **206**: 231-249
- Dunbabin V, Postma J, Schnepf A, Pagès L, Javaux M, Wu L, Leitner D, Chen Y, Rengel Z, Diggle A** (2013) Modelling root–soil interactions using three–dimensional models of root growth, architecture and function. *Plant and Soil*: 1-32
- Dunbabin VM, Diggle AJ, Rengel Z, van Hugten R** (2002) Modelling the interactions between water and nutrient uptake and root growth. *Plant and Soil* **239**: 19-38
- Dupeux F, Santiago J, Betz K, Twycross J, Park S-Y, Rodriguez L, Gonzalez-Guzman M, Jensen MRb, Krasnogor N, Blackledge M** (2011) A thermodynamic switch modulates abscisic acid receptor sensitivity. *EMBO J* **30**: 4171-4184
- Elberling B, Aaskaer L, Jørgensen CJ, Joensen HP, Kuhl M, Glud RN, Lauritsen F** (2011) Linking O₂, CO₂ and CH₄ concentrations in a wetland soil: Implications for CO₂ and CH₄ fluxes. *Environ Science Tech* **45**:3393-3399
- Egan AN, Schlueter J, Spooner DM** (2012) Applications of next-generation sequencing in plant biology. *Am J. Bot* **99**: 175-185
- Eisen MB, Spellman PT, Brown PO, Botstein D** (1998) Cluster analysis and display of genome-wide expression patterns. *Proc Natl Acad Sci* **95**: 14863-14868
- Fahlgren N, Howell MD, Kasschau KD, Chapman EJ, Sullivan CM, Cumbie JS, Givan SA, Law TF, Grant SR, Dangl JL, Carrington JC** (2007) High-throughput sequencing of Arabidopsis microRNAs: evidence for frequent birth and death of miRNA genes. *PLoS One* **2**: e219
- Fiehn O, Kopka J, Dormann P, Altmann T, Trethewey RN, Willmitzer L** (2000) Metabolite profiling for plant functional genomics. *Nat Biotechnol* **18**: 1157-1161
- Friedman N, Linial M, Nachman I, Pe'er D** (2000) Using Bayesian networks to analyze expression data. *Journal of computational biology* **7**: 601-620

- Ge Z, Rubio G, Lynch JP** (2000) The importance of root gravitropism for inter-root competition and phosphorus acquisition efficiency: results from a geometric simulation model. *Plant Soil* **218**: 159-171
- Geisler-Lee J, O'Toole N, Ammar R, Provart NJ, Millar AH, Geisler M** (2007) A predicted interactome for Arabidopsis. *Plant Physiol* **145**: 317-329
- Ghildiyal M, Zamore PD** (2009) Small silencing RNAs: an expanding universe. *Nat Rev Genet* **10**: 94-108
- Gifford ML, Dean A, Gutierrez RA, Coruzzi GM, Birnbaum KD** (2008) Cell-specific nitrogen responses mediate developmental plasticity. *Proc Natl Acad Sci* **105**: 803-808
- Godin C, Sinoquet H** (2005) Functional and structural plant modelling. *New Phytologist* **166**: 705-708
- Grieneisen VA, Xu J, Maree AF, Hogeweg P, Scheres B** (2007) Auxin transport is sufficient to generate a maximum and gradient guiding root growth. *Nature* **449**: 1008-1013
- Hamilton JP, Buell RC** (2012) Advances in plant genome sequencing. *Plant J* **70**: 177-190
- Hammer GL, van Oosterom E, McLean G, Chapman SC, Broad I, Harland P, Muchow RC** (2010) Adapting APSIM to model the physiology and genetics of complex adaptive traits in field crops. *J Exp Bot* **61**: 2185-2202
- Han L, Hanan J, Gresshoff PM** (2010) Computational complementation: a modelling approach to study signalling mechanisms during legume autoregulation of nodulation. *PLoS Comput Biol* **6**: e1000685
- Hartmann A, Rothballer M, Schmid M** (2008) Lorenz Hiltner, a pioneer in rhizosphere microbial ecology and soil bacteriology research. *Plant and Soil* **312**: 7-14
- Hucka M, Finney A, Sauro HM, Bolouri H, Doyle JC, Kitano H, Arkin AP, Bornstein BJ, Bray D, Cornish-Bowden A, Cuellar AA, Dronov S, Gilles ED, Ginkel M, Gor V, Goryanin II, Hedley WJ, Hodgman TC, Hofmeyr JH, Hunter PJ, Juty NS, Kasberger JL, Kremling A, Kummer U, Le Novere N, Loew LM, Lucio D, Mendes P, Minch E, Mjolsness ED, Nakayama Y, Nelson MR, Nielsen PF, Sakurada T, Schaff JC, Shapiro BE, Shimizu TS, Spence HD, Stelling J, Takahashi K, Tomita M, Wagner J, Wang J** (2003) The systems biology markup language (SBML): a medium for representation and exchange of biochemical network models. *Bioinformatics* **19**: 524-531
- Ioio RD, Nakamura K, Moubayidin L, Perilli S, Taniguchi M, Morita MT, Aoyama T, Costantino P, Sabatini S** (2008) A genetic framework for the control of cell division and differentiation in the root meristem. *Science* **322**: 1380-1384
- Iyer-Pascuzzi AS, Jackson T, Cui H, Petricka JJ, Busch W, Tsukagoshi H, Benfey PN** (2011) Cell Identity Regulators Link Development and Stress Responses in the Arabidopsis Root. *Dev Cell* **21**: 770-782
- Javaux M, Schröder T, Vanderborght J, Vereecken H** (2008) Use of a three-dimensional detailed modeling approach for predicting root water uptake. *Vadose Zone Journal* **7**: 1079-1088
- Jeuffroy M-H, Vocanson A, Roger-Estrade J, Meynard J-M** (2012) The use of models at field and farm levels for the ex ante assessment of new pea genotypes. *Eur J Agronomy* **42**: 68-78
- Jiao Y, Tausta SL, Gandotra N, Sun N, Liu T, Clay NK, Ceserani T, Chen M, Ma L, Holford M, Zhang H-Y, Zhao H, Deng X-W, Nelson T** (2009) A

- transcriptome atlas of rice cell types uncovers cellular, functional and developmental hierarchies. *Nat Genet* 41: 258-263
- Jones MA, Raymond MJ, Smirnov N** (2006) Analysis of the root-hair morphogenesis transcriptome reveals the molecular identity of six genes with roles in root-hair development in *Arabidopsis*. *Plant J* 45: 83-100
- Kang YH, Kirik V, Hulskamp M, Nam KH, Hagely K, Lee MM, Schiefelbein J** (2009) The MYB23 gene provides a positive feedback loop for cell fate specification in the *Arabidopsis* root epidermis. *Plant Cell* 21: 1080-1094
- Kasschau KD, Fahlgren N, Chapman EJ, Sullivan CM, Cumbie JS, Givan SA, Carrington JC** (2007) Genome-wide profiling and analysis of *Arabidopsis* siRNAs. *PLoS Biol* 5: e57
- Katari MS, Nowicki SD, Aceituno FF, Nero D, Kelfer J, Thompson LP, Cabello JM, Davidson RS, Goldberg AP, Shasha DE, Coruzzi GM, Gutiérrez RA** (2010) VirtualPlant: A Software Platform to Support Systems Biology Research. *Plant Physiol* 152: 500-515
- Keating BA, Carberry P, Hammer G, Probert ME, Robertson M, Holzworth D, Huth N, Hargreaves J, Meinke H, Hochman Z** (2003) An overview of APSIM, a model designed for farming systems simulation. *Eur J Agronomy* 18: 267-288
- Kerk NM, Ceserani T, Tausta SL, Sussex IM, Nelson TM** (2003) Laser capture microdissection of cells from plant tissues. *Plant Physiol* 132: 27-35
- Keyes SD, Boardman RP, Marchant A, Roose T, Sinclair I** (2013) A robust approach for determination of the macro-porous volume fraction of soils with X-ray computed tomography and an image processing protocol. *Eur J Soil Sci* 64: 298-307
- Kierzkowski D, Nakayama N, Routier-Kierzkowska A-L, Weber A, Bayer E, Schorderet M, Reinhardt D, Kuhlemeier C, Smith RS** (2012) Elastic Domains Regulate Growth and Organogenesis in the Plant Shoot Apical Meristem. *Science* 335: 1096-1099
- Kloppfleisch K, Phan N, Augustin K, Bayne RS, Booker KS, Botella JR, Carpita NC, Carr T, Chen JG, Cooke TR, Frick-Cheng A, Friedman EJ, Fulk B, Hahn MG, Jiang K, Jorda L, Kruppe L, Liu C, Lorek J, McCann MC, Molina A, Moriyama EN, Mukhtar MS, Mudgil Y, Pattathil S, Schwarz J, Seta S, Tan M, Temp U, Trusov Y, Urano D, Welter B, Yang J, Panstruga R, Uhrig JF, Jones AM** (2011) *Arabidopsis* G-protein interactome reveals connections to cell wall carbohydrates and morphogenesis. *Mol Syst Biol* 7: 532
- Krouk G, Lacombe B, Bielach A, Perrine-Walker F, Malinska K, Mounier E, Hoyerova K, Tillard P, Leon S, Ljung K** (2010) Nitrate-regulated auxin transport by NRT1.1 defines a mechanism for nutrient sensing in plants. *Dev Cell* 18: 927-937
- Kwak S-H, Schiefelbein J** (2008) A Feedback Mechanism Controlling SCRAMBLED Receptor Accumulation and Cell-Type Pattern in *Arabidopsis*. *Current Biol* 18: 1949-1954
- Kwasniewski M, Nowakowska U, Szumera J, Chwialkowska K, Szarejko I** (2013) iRootHair: a comprehensive root hair genomics database. *Plant Physiol* 161: 28-35
- Lalonde S, Sero A, Pratelli R, Pilot G, Chen J, Sardi MI, Parsa SA, Kim DY, Acharya BR, Stein EV, Hu HC, Villiers F, Takeda K, Yang Y, Han YS, Schwacke R, Chiang W, Kato N, Loque D, Assmann SM, Kwak JM,**

- Schroeder JI, Rhee SY, Frommer WB** (2010) A membrane protein/signaling protein interaction network for Arabidopsis version AMPv2. *Front Physiol* **1**: 24
- Lau S, De Smet I, Kolb M, Meinhardt H, Jurgens G** (2011) Auxin triggers a genetic switch. *Nat Cell Biol* **13**: 611-615
- Li C, Donizelli M, Rodriguez N, Dharuri H, Endler L, Chelliah V, Li L, He E, Henry A, Stefan MI, Snoep JL, Hucka M, Le Novère N, Laibe C** (2010) BioModels Database: An enhanced, curated and annotated resource for published quantitative kinetic models. *BMC syst biol* **4**: 92
- Liao H, Yan X, Rubio G, Beebe SE, Blair MW, Lynch JP** (2004) Genetic mapping of basal root gravitropism and phosphorus acquisition efficiency in common bean. *Functional Plant Biology* **31**: 959-970
- Libault M, Farmer A, Brechenmacher L, Drnevich J, Langley RJ, Bilgin DD, Radwan O, Neece DJ, Clough SJ, May GD, Stacey G** (2010) Complete transcriptome of the soybean root hair cell, a single-cell model, and its alteration in response to *Bradyrhizobium japonicum* infection. *Plant Physiol* **152**: 541-552
- Lin W-D, Liao Y-Y, Yang TJ, Pan C-Y, Buckhout TJ, Schmidt W** (2011) Coexpression-based clustering of Arabidopsis root genes predicts functional modules in early phosphate deficiency signaling. *Plant Physiol* **155**: 1383-1402
- Lobet G, Draye X** (2013) Novel scanning procedure enabling the vectorization of entire rhizotron-grown root systems. *Plant Methods* **9**: 1
- Lucas M, Godin C, Jay-Allemand C, Laplaze L** (2008) Auxin fluxes in the root apex co-regulate gravitropism and lateral root initiation. *J Exp Bot* **59**: 55-66
- Lynch J** (1995) Root Architecture and Plant Productivity. *Plant Physiol* **109**: 7-13
- Lynch J, Brown K** (2001) Topsoil foraging – an architectural adaptation of plants to low phosphorus availability. *Plant and Soil* **237**: 225-237
- Lynch JM, Whipps JM** (1990) Substrate flow in the rhizosphere. *Plant and Soil* **129**: 1-10
- Lynch JP** (2007) Roots of the Second Green Revolution. *Aus J Bot* **55**: 493-512
- Lynch JP** (2011) Root phenes for enhanced soil exploration and phosphorus acquisition: tools for future crops. *Plant Physiol* **156**: 1041-1049
- Lynch JP** (2013) Steep, cheap and deep: an ideotype to optimize water and N acquisition by maize root systems. *Ann of Botany* **112**: 347-57
- Lynch JP, Nielsen KL, Davis RD, Jablankow AG** (1997) SimRoot: modelling and visualization of root systems. *Plant and Soil* **188**: 139-151
- Ma Z, Bielenberg DG, Brown KM, Lynch JP** (2001) Regulation of root hair density by phosphorus availability in *Arabidopsis thaliana*. *Plant, Cell & Environment* **24**: 459-467
- Manschadi A, Christopher J, Hammer G, Devoil P** (2010) Experimental and modelling studies of drought-adaptive root architectural traits in wheat (*Triticum aestivum* L.). *Plant Biosystems* **144**: 458-462
- Mardis ER** (2008) The impact of next-generation sequencing technology on genetics. *Trends in genetics* **24**: 133
- Middleton A, King J, Bennett M, Owen M** (2010) Mathematical modelling of the Aux/IAA negative feedback loop. *Bulletin of mathematical biology* **72**: 1383-1407

- Middleton AM, Farcot E, Owen MR, Vernoux T** (2012) Modeling regulatory networks to understand plant development: small is beautiful. *Plant Cell* **24**: 3876-3891
- Mooney SJ, Pridmore TP, Helliwell J, Bennett MJ** (2012) Developing X-ray Computed Tomography to non-invasively image 3-D root systems architecture in soil. *Plant and Soil* **352**: 1-22
- Mosconi F, Julou T, Desprat N, Sinha DK, Allemand J-F, Croquette V, Bensimon D** (2008) Some nonlinear challenges in biology. *Nonlinearity* **21**: T131
- Moussaieff A, Rogachev I, Brodsky L, Malitsky S, Toal TW, Belcher H, Yativ M, Brady SM, Benfey PN, Aharoni A** (2013) High-resolution metabolic mapping of cell types in plant roots. *Proc Natl Acad Sci* **110**: 1232-41
- Muraro D, Byrne H, King J, Voß U, Kieber J, Bennett M** (2011) The influence of cytokinin-auxin cross-regulation on cell-fate determination in *Arabidopsis thaliana* root development. *J Theoretical Biol* **283**: 152-167
- Nawy T, Lee J-Y, Colinas J, Wang JY, Thongrod SC, Malamy JE, Birnbaum K, Benfey PN** (2005) Transcriptional profile of the *Arabidopsis* root quiescent center. *Plant Cell* **17**: 1908-1925
- Nemhauser JL, Hong F, Chory J** (2006) Different plant hormones regulate similar processes through largely nonoverlapping transcriptional responses. *Cell* **126**: 467-475
- Noble D** (2008) Claude Bernard, the first systems biologist, and the future of physiology. *Exp Physiol* **93**: 16-26
- Noor A, Serpedin E, Nounou M, Nounou H, Mohamed N, Chouchane L** (2013) An overview of the statistical methods used for inferring gene regulatory networks and protein-protein interaction networks. *Adv Bioinformatics* **2013**: 953814
- Okushima Y, Overvoorde PJ, Arima K, Alonso JM, Chan A, Chang C, Ecker JR, Hughes B, Lui A, Nguyen D, Onodera C, Quach H, Smith A, Yu G, Theologis A** (2005) Functional Genomic Analysis of the AUXIN RESPONSE FACTOR Gene Family Members in *Arabidopsis thaliana*: Unique and Overlapping Functions of ARF7 and ARF19. *Plant Cell* **17**: 444-463
- Ozsolak F, Milos PM** (2011) RNA sequencing: advances, challenges and opportunities. *Nat Rev Genet* **12**: 87-98
- Pagès L, Vercambre G, Drouet J-L, Lecompte F, Collet C, Le Bot J** (2004) Root Typ: a generic model to depict and analyse the root system architecture. *Plant and Soil* **258**: 103-119
- Paponov IA, Paponov M, Teale W, Menges M, Chakrabortee S, Murray JA, Palme K** (2008) Comprehensive transcriptome analysis of auxin responses in *Arabidopsis*. *Mol Plant* **1**: 321-337
- Parizot B, De Rybel B, Beeckman T** (2010) VisualRTC: A New View on Lateral Root Initiation by Combining Specific Transcriptome Data Sets. *Plant Physiol* **153**: 34-40
- Péret B, Larrieu A, Bennett MJ** (2009) Lateral root emergence: a difficult birth. *J Exp Bot* **60**: 3637-3643
- Péret B, Li G, Zhao J, Band LR, Vos U, Postaire O, Luu D-T, Da Ines O, Casimiro I, Lucas MI** (2012) Auxin regulates aquaporin function to facilitate lateral root emergence. *Nat Cell Biol* **14**: 991-998

- Perrin B-E, Ralaivola L, Mazurie A, Bottani S, Mallet J, d'Alché-Buc F** (2003) Gene networks inference using dynamic Bayesian networks. *Bioinformatics* **19**: ii138-ii148
- Perrine-Walker F, Doumas P, Lucas M, Vaissayre V, Beauchemin NJ, Band LR, Chopard J, Crabos A, Conejero G, Peret B, King JR, Verdeil JL, Hocher V, Franche C, Bennett MJ, Tisa LS, Laplaze L** (2010) Auxin carriers localization drives auxin accumulation in plant cells infected by *Frankia* in *Casuarina glauca* actinorhizal nodules. *Plant Physiol* **154**: 1372-1380
- Petricka JJ, Schauer MA, Megraw M, Breakfield NW, Thompson JW, Georgiev S, Soderblom EJ, Ohler U, Moseley MA, Grossniklaus U, Benfey PN** (2012) The protein expression landscape of the *Arabidopsis* root. *Proc Natl Acad Sci* **109**: 6811-6818
- Popescu SC, Popescu GV, Bachan S, Zhang Z, Gerstein M, Snyder M, Dinesh-Kumar SP** (2009) MAPK target networks in *Arabidopsis thaliana* revealed using functional protein microarrays. *Genes Dev* **23**: 80-92
- Popescu SC, Popescu GV, Bachan S, Zhang Z, Seay M, Gerstein M, Snyder M, Dinesh-Kumar SP** (2007) Differential binding of calmodulin-related proteins to their targets revealed through high-density *Arabidopsis* protein microarrays. *Proc Natl Acad Sci* **104**: 4730-4735
- Postma JA, Lynch JP** (2011a) Root cortical aerenchyma enhances the growth of maize on soils with suboptimal availability of nitrogen, phosphorus, and potassium. *Plant Physiol* **156**: 1190-1201
- Postma JA, Lynch JP** (2011) Theoretical evidence for the functional benefit of root cortical aerenchyma in soils with low phosphorus availability. *Ann Bot* **107**: 829-841
- Poultney CS, Gutiérrez RA, Katari MS, Gifford ML, Paley WB, Coruzzi GM, Shasha DE** (2007) Sungear: interactive visualization and functional analysis of genomic datasets. *Bioinformatics* **23**: 259-261
- Pradal C, Dufour-Kowalski S, Boudon F, Fournier C, Godin C** (2008) OpenAlea: a visual programming and component-based software platform for plant modelling. *Functional Plant Biology* **35**: 751-760
- Prusinkiewicz P** (2004) Art and science for life: Designing and growing virtual plants with L-systems. *In* C Davidson, T Fernandez, eds, *Nursery Crops: Development, Evaluation, Production and Use: Proceedings of the XXVI International Horticultural Congress, Vol 360. Acta Horticulturae*, pp 15-28
- Prusinkiewicz P, Erasmus Y, Lane B, Harder LD, Coen E** (2007) Evolution and Development of Inflorescence Architectures. *Science* **316**: 1452-1456
- Roose T, Schnepf A** (2008) Mathematical models of plant-soil interaction. *Philos Trans R Soc Lond B Biol Sci* **366**: 4597-4611
- Rubio G, Zhu J, Lynch JP** (2003) A critical test of the two prevailing theories of plant response to nutrient availability. *Am J Bot* **90**: 143-152
- Ruffel S, Krouk G, Coruzzi GM** (2010) A systems view of responses to nutritional cues in *Arabidopsis*: toward a paradigm shift for predictive network modeling. *Plant Physiol* **152**: 445-452
- Savage NS, Walker T, Wieckowski Y, Schiefelbein J, Dolan L, Monk NAM** (2008) A Mutual Support Mechanism through Intercellular Movement of CAPRICE and GLABRA3 Can Pattern the *Arabidopsis* Root Epidermis. *PLoS Biol* **6**: e235

- Ruffel S, Krouk G, Ristova D, Shasha D, Birnbaum KD, Coruzzi GM** (2011) Nitrogen economics of root foraging: Transitive closure of the nitrate-cytokinin relay and distinct systemic signaling for N supply vs. demand. *Proc Natl Acad Sci* **108**: 18524-18529
- Sablowski R** (2011) Plant stem cell niches: from signalling to execution. *Current Opinion in Plant Biology* **14**: 4-9
- Schröder T, Javaux M, Vanderborght J, Körfgen B, Vereecken H** (2008) Effect of local soil hydraulic conductivity drop using a three-dimensional root water uptake model. *Vadose Zone Journal* **7**: 1089-1098
- Sievänen R, Nikinmaa E, Nygren P, Ozier-Lafontaine H, Perttunen J, Hakula H** (2000) Components of functional-structural tree models. *Ann. For. Sci.* **57**: 399-412
- Stahl Y, Simon R** (2010) Plant primary meristems: shared functions and regulatory mechanisms. *Current Opinion in Plant Biology* **13**: 53-58
- Stoma S, Lucas M, Chopard J, Schaedel M, Traas J, Godin C** (2008) Flux-Based Transport Enhancement as a Plausible Unifying Mechanism for Auxin Transport in Meristem Development. *PLoS Comput Biol* **4**: e1000207
- Swarup K, Benková E, Swarup R, Casimiro I, Péret B, Yang Y, Parry G, Nielsen E, De Smet I, Vanneste S, Levesque MP, Carrier D, James N, Calvo V, Ljung K, Kramer E, Roberts R, Graham N, Marillonnet S, Patel K, Jones JD, Taylor CG, Schachtman DP, May S, Sandberg G, Benfey P, Friml J, Kerr I, Beeckman T, Laplace L, Bennett MJ** (2008) The auxin influx carrier LAX3 promotes lateral root emergence. *Nat Cell Biol* **10**: 946-954
- Tanz SK, Castleden I, Hooper CM, Vacher M, Small I, Millar HA** (2013) SUBA3: a database for integrating experimentation and prediction to define the SUBcellular location of proteins in Arabidopsis. *Nuc Acids Res* **41**: 1185-1191
- Ubeda-Tomás S, Swarup R, Coates J, Swarup K, Laplace L, Beemster GT, Hedden P, Bhalerao R, Bennett MJ** (2008) Root growth in Arabidopsis requires gibberellin/DELLA signalling in the endodermis. *Nat Cell Biol* **10**: 625-628
- Van Landeghem S, De Bodt S, Drebert ZJ, Inzé D, Van de Peer Y** (2013) The Potential of Text Mining in Data Integration and Network Biology for Plant Research: A Case Study on Arabidopsis. *Plant Cell* **25**: 794-807
- Vidal EA, Moyano TC, Riveras E, Contreras-Lopez O, Gutierrez RA** (2013) Systems approaches map regulatory networks downstream of the auxin receptor AFB3 in the nitrate response of Arabidopsis thaliana roots. *Proc Natl Acad Sci* **110**: 12840-12845
- Vidal EA, Tamayo KP, Gutierrez RA** (2010) Gene networks for nitrogen sensing, signaling, and response in Arabidopsis thaliana. *Wiley Interdiscip Rev Syst Biol Med* **2**: 683-693
- Vos J, Evers JB, Buck-Sorlin GH, Andrieu B, Chelle M, de Visser PH** (2010) Functional-structural plant modelling: a new versatile tool in crop science. *J Exp Bot* **61**: 2101-2115
- Wells D, Laplace L, Bennett MJ, Vernoux T** (2013) Biosensors for phytohormone quantification: challenges, solutions and opportunities. *Trends in Plant Science*.18:244-249
- Willis RC, Hogue CW** (2006) Searching, viewing, and visualizing data in the Biomolecular Interaction Network Database (BIND). *Curr Protoc Bioinformatics* **Chapter 8**: Unit 8-9

- Winter D, Vinegar B, Nahal H, Ammar R, Wilson GV, Provart NJ** (2007) An electronic fluorescent pictograph browser for exploring and analyzing large-scale biological data sets. *Plos One* **2**: e718
- Won S-K, Lee Y-J, Lee H-Y, Heo Y-K, Cho M, Cho H-T** (2009) Cis-element-and transcriptome-based screening of root hair-specific genes and their functional characterization in Arabidopsis. *Plant physiol* **150**: 1459-1473
- Yan X, Liao H, Beebe S, Blair M, Lynch J** (2004) QTL mapping of root hair and acid exudation traits and their relationship to phosphorus uptake in common bean. *Plant and Soil* **265**: 17-29
- Yu J, Smith VA, Wang PP, Hartemink AJ, Jarvis ED** (2004) Advances to Bayesian network inference for generating causal networks from observational biological data. *Bioinformatics* **20**: 3594-3603
- Zanetti ME, Chang F, Gong F, Galbraith DW, Bailey-Serres J** (2005) Immunopurification of polyribosomal complexes of Arabidopsis for global analysis of gene expression. *Plant Physiol* **138**: 624-635

Table 1 Databases, data mining and analysis tools for root systems biology.

Freely available databases and mining tools for root systems biology. Many of these databases also may be useful for study of other plant organs and systems.

Resource	Description	Reference
VirtualPlant 1.3 http://www.virtualplant.org	Integrates genomic data and provides visualization and analysis tools to aid generation of biological hypotheses. Also includes predicted and validated protein-DNA interactions	Katari et al., 2010
Sungear http://virtualplant.bio.nyu.edu/cgi-bin/sungear/index.cgi	Enables rapid, visually interactive exploration of large sets of genomic data. Allows browsing of gene sets by experiment, gene annotation, and ontological term. Makes otherwise complicated queries quick and visually intuitive.	Part of the VirtualPlant suite of tools
Arabidopsis eFP Browser http://bar.utoronto.ca/efp/cgi-bin/efpWeb.cgi	Enables visualization of relative and absolute gene expression data from ~22,000 <i>Arabidopsis</i> genes in different tissues and under a variety of conditions	Winter et al., 2007
GENIUS (Gene Network Inference Using Signatures) http://networks.bio.puc.cl/genius/	Uses large-scale gene expression data to identify signatures that characterize and discriminate the process of interest, and then uses these signatures to predict a functional gene network.	
VisuaLRTC	Lateral root transcriptome compendium	Parizot et al., 2010
iROOTHAIR http://www.iroothair.org/	A comprehensive database of root hair genomics information	Kwasniewski et al., 2013
SUBA3	Combines manual literature curation of	Tanz et al., 2013

http://suba.plantenergy.uwa.edu.au	<p>large-scale subcellular proteomics, fluorescent protein visualization and protein–protein interaction datasets from <i>Arabidopsis</i> with subcellular targeting calls from 22 prediction programs.</p>	
<p>Arabidopsis interaction viewer</p> <p>http://bar.utoronto.ca/interactions/cgi-bin/arabidopsis_interactions_viewer.cgi</p>	<p>Queries a database of 70944 predicted and 28556 confirmed <i>Arabidopsis</i> interacting proteins against the protein(s) of interest. Includes subcellular localization from SUBA</p>	<p>Geisler-Lee et al., 2007</p>
<p>PlantMetabolomics.org (PM)</p> <p>http://plantmetabolomics.vrac.iastate.edu/ver2/</p>	<p>An NSF-funded multi-institutional project that is developing metabolomics as a functional genomics tool for elucidating the functions of <i>Arabidopsis</i> genes. The Consortium has established metabolomic platforms that detect approximately 1,800 metabolites.</p>	<p>Bais et al., 2010</p>
<p>Ionomics Hub (iHub)</p> <p>http://www.ionomicshub.org/home/PiiMS</p>	<p>Contains curated ionomic data on many thousands of plant samples (from <i>Arabidopsis</i>, <i>Brassica napus</i>, rice, maize, and soybean).</p>	

Figure 1. Examples of top-down approaches to infer gene regulatory networks.




Figure 1. Examples of top-down approaches to infer gene regulatory networks.

(far left column): Krouk et al., 2010 combined time-series transcriptomic data with published hormone-treatment datasets (Nemhauser et al 2006) to obtain a temporal model for nitrate-driven gene expression networks. They then used a machine learning approach (state-space modeling) to infer a regulatory network governing nitrate response over a time.

(middle, left column): Ruffel et al., 2011 combined both genetic and genomic approaches within a 'split-root' system to study how systemic nitrate signaling effected genome-wide programming in root development.

(right column): Brady et al., 2011 employed a modeling approach to predict the regulatory potential of each of the relationships they identified within the stele GRN. Stele expressed transcription factors were selected from the GRN dataset published previously (Brady et al., 2007). Protein-protein and protein DNA maps were generated from Y1H and Y2H screens of stele expressed TFs and miRNAs. The key interactions between TFs and their target genes were subsequently measured by qRT-PCR analysis target gene expression in wild type and mutant lines. Regulatory interaction strength is represented by edge and arrow width with thicker lines or arrowheads representing steeper slopes. P-value strength is represented by edge opacity with darker edges representing more significant interactions. Slopes were determined by plotting quantitative PCR (qPCR) expression values of the transcription factor (TF) and its target and fitting a line using weighted least squares regression (WLSR). This predictive framework enables the identification of the most influential upstream activators or repressors to be manipulated in order to regulate expression of a target.

Figure 2. Examples of hypothesis driven bottom-up and middle-out approaches.

Examples of hypothesis driven 'bottom-up' and 'middle-out' approaches




Figure 2. Examples of hypothesis driven bottom-up and middle-out approaches.

Models were constructed for processes at different spatial and temporal scales. (A) At the sub-cellular level, gene regulatory networks can be modeled to include the input of signals such as auxin and output in the form of cell-scale events, such as mitosis. (B) Stem cell division and the bistable switch that regulates it are modeled using longitudinal gradients of auxin and radial gradients of the transcription factor SHORTROOT (SHR) (Cruz-Ramirez et al., 2012). (C) Root system scale models, such as RootTyp (Pagés et al., 2004) can be linked to hydraulic models to explicitly simulate water flow in the soil-plant domain (Javaux et al., 2008).

Figure 3: Root biological models can be multi-scale.




Figure 3: Root biological models can be multi-scale.

Band et al. (2012b) demonstrate the power of multi-scale modeling for understanding gibberellin dynamics during root elongation. The elongation zone is modeled as a collection of cells (organ scale) (A) and used to motivate modelling of cell expansion and dilution in the vacuole (cell scale) (B). This naturally leads to the link between cell partitioning of gibberellin and sub-cellular processes, revealing the influence of dilution of gibberellin concentrations on DELLA expression and growth.