

1 **Blue light regulation of stomatal opening and the plasma membrane H⁺-ATPase**

2

3 **Shin-ichiro Inoue^{a,*} and Toshinori Kinoshita^{a,b,*}**

4

5 ^aDivision of Biological Science, Graduate School of Science, Nagoya University,
6 Chikusa, Nagoya 464-8602, Japan; ^bInstitute of Transformative Bio-Molecules
7 (WPI-ITbM), Nagoya University, Chikusa, Nagoya 464-8602, Japan

8 *Address correspondence to shin@bio.nagoya-u.ac.jp and
9 kinoshita@bio.nagoya-u.ac.jp

10

11

12 This work was supported by KAKENHI from the Japan Society for the Promotion of
13 Science (nos. 25840105 and 15K07101 to S.I. and JP15H05956 and 15H04386 to T.K.).

14

15

16

17

18

19

20

21

22

23 Light-induced stomatal responses were first reported by Francis Darwin (1898).
24 Stomata open in response to light, including blue and red light (Shimazaki et al., 2007).
25 Red light induces stomatal opening via photosynthesis in the mesophyll and guard cell
26 chloroplasts (Mott et al., 2008; Suetsugu et al., 2014). In contrast, blue light as a signal
27 induces stomatal opening. Phototropins expressed in guard cells act as major blue light
28 receptors for stomatal opening (Kinoshita et al., 2001; Kinoshita et al., 2003; Inoue et
29 al., 2008). Blue light-induced stomatal opening is mediated through activation of a
30 plasma membrane (PM) H⁺ pump, later identified as the PM H⁺-ATPase, in guard cells
31 (Assmann et al., 1985; Shimazaki et al., 1986; Kinoshita and Shimazaki, 1999). The
32 blue light-activated pump provides driving force for stomatal opening concomitant with
33 ion accumulation and cell volume increase in guard cells (Schroeder et al., 1987;
34 Kinoshita and Hayashi, 2011). Note that stomatal opening in response to weak blue
35 light as a signal requires background red light, indicating that red light has a synergistic
36 effect on the blue light response in guard cells (Shimazaki et al., 2007).

37 Recent investigations of guard cells with respect to blue light-induced stomatal
38 opening have greatly advanced our understanding. In this review, we focus on the recent
39 progress of the blue light signaling pathway in guard cells and its regulation of the PM
40 H⁺-ATPase activity.

41

42 **Blue light-signaling for stomatal opening**

43 Stomata effectively open in response to blue light, especially under strong red light
44 (Shimazaki et al., 2007; Marten et al., 2010). Our understanding of the signaling model
45 for stomatal opening was mainly constructed from studies of blue light-induced
46 stomatal opening (Fig. 1). A single guard cell possesses all signaling components, from
47 blue light-perception to cell-volume increase, for stomatal opening. When guard cells
48 are irradiated by blue light, blue light-photoreceptor protein kinases, phototropins, are
49 activated through autophosphorylation and initiate signaling for stomatal opening
50 (Kinoshita et al., 2001; Christie, 2007). Blue light induces autophosphorylation of two
51 Ser residues in the kinase activation-loop of phototropin molecules, and
52 phosphorylation is required for downstream signaling, probably through substrate
53 recognition (Inoue et al., 2008; Inoue et al., 2010; Inoue et al., 2011). The activated
54 phototropins directly phosphorylate another protein kinase BLUE LIGHT SIGNALING
55 1 (BLUS1), and phosphorylated BLUS1 indirectly transmits the signal to type 1 protein

Figure 1




Figure 1. Blue light signaling pathway in stomatal guard cells.

Arrows and a T-bar represent positive and negative regulation, respectively. The P in the white circles indicates a phosphorylation of each protein. phot, phototropins; BLUS1, BLUE LIGHT SIGNALING1; BHP, BLUE LIGHT-DEPENDENT H⁺-ATPASE PHOSPHORYLATION; PP1, Protein phosphatase 1; PRSL1, PP1 regulatory subunit2-like protein1; 14-3-3, 14-3-3 protein; K_{in} channel, inward rectifying K⁺ channel; TAG, triacylglycerols; PAR, photosynthetically active radiation; Chl., Chloroplasts; NHX1/2, Na⁺/H⁺ EXCHANGER 1/2; ALMT9, aluminium-activated malate transporter 9; CLCc, chloride channel c. The time scale of each peak of the key signaling events for blue light-induced stomatal opening (approximately 2 h) is shown as follows: phototropin activation (within 1 min), H⁺ pumping (approximately 2.5 min), Hyperpolarization (several min), K⁺ accumulation (between 30 and 60 min). TAG breakdown, starch degradation, and vacuolar remodeling are observed within 1 to 2 h after the start of light illumination.

56 phosphatase (PP1) and its regulatory subunit PRSL1 (Takemiya et al., 2006; Takemiya
 57 et al., 2013a; Takemiya et al., 2013b; Takemiya et al., 2016). Note that BLUS1
 58 expression is specific to guard cells and is not involved in the other

59 phototropin-mediated responses, such as phototropism, chloroplasts movements, and
60 leaf flattening (Takemiya et al., 2013a), suggesting that BLUS1 defines signaling
61 specificity of stomatal opening among the phototropin-mediated responses. The signal
62 generated by BLUS1 finally activates the PM H⁺-ATPase, mainly isoform AHA1, in
63 guard cells through phosphorylation of a penultimate threonine in the C-terminus with
64 subsequent binding of a 14-3-3 protein (Shimazaki et al., 2007; Hayashi et al., 2011;
65 Yamauchi et al., 2016).

66 Very recently, a Raf-like protein kinase, BLUE LIGHT-DEPENDENT H⁺-ATPASE
67 PHOSPHORYLATION (BHP), was reported as the novel signaling component in blue
68 light-dependent stomatal opening (Hayashi et al., 2017). BHP was identified through a
69 screening of protein kinase inhibitors that suppress blue light-dependent PM H⁺-ATPase
70 phosphorylation in guard cells and similarities to the mammalian targets of the
71 inhibitors. BHP does not bind to the PM H⁺-ATPase but to BLUS1 and forms an early
72 signaling complex with phototropins to mediate phosphorylation of a penultimate
73 threonine of the PM H⁺-ATPase (Hayashi et al., 2017). Guard cells in *bhp* mutant
74 exhibited normal phosphorylation of the PM H⁺-ATPase in response to the PM
75 H⁺-ATPase activator fusicoccin, suggesting that BHP is not likely to directly
76 phosphorylate the penultimate threonine of PM H⁺-ATPase. There may be an
77 unidentified protein kinase that directly phosphorylates the PM H⁺-ATPase in stomatal
78 opening. In addition, whether BLUS1 phosphorylates and activates BHP is unknown at
79 the present time. Further analyses are needed to clarify the early signaling for stomatal
80 opening from phototropins to the PM H⁺-ATPase activation and to identify the
81 endogenous substrates of BLUS1, BHP, and PP1 in guard cells.

82 The blue light-activated PM H⁺-ATPase drives H⁺ transport across the PM and
83 hyperpolarizes of the membrane (Shimazaki et al., 2007; Marten et al., 2010). This
84 membrane hyperpolarization activates inward-rectifying K⁺ (K⁺_{in}) channels and induces
85 an influx of K⁺ (Lebaudy et al., 2008; Kim et al., 2010), resulting in the accumulation of
86 K⁺, and the counter anions Cl⁻, nitrate, and malate²⁻. K⁺ and Cl⁻ are immediately
87 transported into the vacuole via the tonoplast-localized K⁺/H⁺ exchangers NHX1 and
88 NHX2 and vacuolar chloride channels ALMT9 and CLCc, respectively (Jossier et al.,
89 2010; Chen et al., 2012; De Angeli et al., 2013; Andrés et al., 2014), and K⁺
90 accumulation into the vacuole involves dynamic remodeling of vacuolar structure for
91 stomatal opening (Andrés et al., 2014). Accumulation of these ions decreases the water

92 potential of guard cells, which leads to water uptake into the vacuole and turgor increase,
93 leading to stomatal opening (Inoue et al., 2010; Marten et al., 2010). The details of ion
94 transports in guard cells are reviewed in this Focus Issue (Eisenacha and De Angeli,
95 2017; Jezek and Blatt, 2017). Recently, Santelia and colleagues demonstrated that starch
96 in guard cell chloroplasts is degraded by phototropin-mediated signaling downstream of
97 PM H⁺-ATPase activity and the degradation contributes to stomatal opening (Horrer et
98 al., 2016), probably through malate synthesis (Shimazaki et al. 2007). It remains an
99 interesting and important question on how the activity of the PM H⁺-ATPase is linked to
100 the starch degradation pathway in chloroplasts (Santelia and Lunn, 2017).

101 In addition to the pursuit of signaling components as described above, regulation of
102 the expression and localization of key signaling factors that determines stomatal
103 aperture indirectly have also been investigated recently. For example, the bHLH family
104 transcription factors of ABA-RESPONSIVE KINASE SUBSTRATEs (AKSs) and
105 GARP transcription factors of GOLDEN 2-LIKE 1 and 2 (GLK1 and 2) bind to the
106 promoter of the K⁺_{in} channel *KATI* gene and increase the K⁺_{in} channel expression in
107 guard cells (Takahashi et al., 2013; Nagatoshi et al., 2016). GLKs also positively
108 regulate *BLUSI* expression (Nagatoshi et al., 2016). These transcriptional regulations
109 contribute to enhance stomatal opening. Moreover, the signaling components for
110 photoperiodic flowering including cryptochromes, GIGANTEA, CONSTANS, EARLY
111 FLOWERING 3, FLOWERING LOCUS T (FT), TWIN SISTER OF FT, and
112 SUPPRESSOR OF OVEREXPRESSION OF CO 1, are expressed in guard cells and
113 also affect light-induced stomatal opening probably via transcriptional regulations in
114 guard cells, but details of the regulatory targets are unknown (Kinoshita et al., 2011;
115 Ando et al., 2013; Kimura et al., 2015). In addition, the Munc13-like protein PATROL1
116 may be involved in the enhancement of light-induced stomatal opening by promoting
117 the recruitment of AHA1 to the PM of guard cells (Hashimoto-Sugimoto et al., 2013).
118 Both aminophospholipid ATPase (ALA10) flippase and phospholipase A₂β (PL A₂β) are
119 involved in light-induced stomatal opening through lysophospholipid generation-(Seo et
120 al., 2008; Poulsen et al., 2015). ALA10 promotes phospholipid uptake into guard cells
121 and PL A₂β generates lysophospholipids using phospholipids as a substrate. One of the
122 products, lysophosphatidylcholine functions as a specific activator of the PM
123 H⁺-ATPase (Palmgren, 2001), because both ALA10 and PL A₂β may have a positive
124 effect on the stomatal opening through enhance the PM H⁺-ATPase activity.

125 Furthermore, photosynthetic processes in guard cell chloroplasts provide fuel (ATP
126 and/or reducing equivalents) for blue light-dependent H⁺ pumping of the PM
127 H⁺-ATPase and contribute to stomatal opening (Suetsugu et al., 2014). Correspondingly,
128 guard cells lacking chloroplasts in the *crumpled leaf* mutant displayed attenuation of
129 both guard cell ATP levels and stomatal opening in response to light (Wang et al., 2014).
130 Stored triacylglycerols in guard cells are broken down in response to light and the
131 catabolic process is also thought to supply ATP for PM H⁺-ATPase activity in stomatal
132 opening (McLachlan et al., 2016).

133

134 **Crosstalk between blue light- and ABA-signaling in light-induced stomatal opening**

135 The plant hormone ABA synthesized in response to drought stress conditions drastically
136 reduces stomatal aperture to prevent water loss in the presence of light (Bauer et al.,
137 2013; Waadt et al., 2014; Kim et al., 2010; Osakabe et al., 2014). ABA induces stomatal
138 closure in already open stomata, called “stomatal closure” and simultaneously inhibits
139 light-induced stomatal opening, called “inhibition of stomatal opening”, and both
140 physiological regulatory mechanisms are required to close stomata efficiently under
141 sunlight. It is known that ABA-signaling in guard cells effectively suppresses blue
142 light-signaling in inhibition of stomatal opening with various ways. First, ABA
143 accelerates the release of ions from guard cells by activating S- and R-type anion
144 channels, outward-rectifying K⁺ channels, and K⁺ uptake transporters (KUPs) in
145 stomatal closure (Negi et al., 2008; Vahisalu et al., 2008; Kim et al., 2010; Osakabe et
146 al., 2013). ABA suppresses these processes via ABA-receptor components
147 PYR/PYL/RCAR-PP2Cs-SnRK2s in guard cells. (Fujii et al., 2009; Ma et al., 2009;
148 Park et al., 2009; Umezawa et al., 2009; Vlad et al., 2009; Jezek and Blatt, 2017) with
149 subsequent second messengers H₂O₂, H₂S, NO, phosphatidic acid (PA), and cytosolic
150 Ca²⁺ (Inoue et al., 2010; Scuffi et al., 2014). It has been reported that H₂S also functions
151 in parallel to ABA signaling events (Papanatsiou et al., 2015). Simultaneously, in the
152 inhibition of stomatal opening, ABA suppresses blue light-signaling of the activation of
153 PM H⁺-ATPase via ABA-receptor components and H₂O₂, NO, PA, and Ca²⁺ (Shimazaki
154 et al., 2007; Zhang et al., 2007; Inoue et al., 2010; Kim et al., 2010; Takemiya and
155 Shimazaki, 2010; Hayashi et al., 2011; Hayashi and Kinoshita, 2011). PA directly
156 inhibits PP1 catalytic activity and blocks blue light-signaling between phototropins and
157 the PM H⁺-ATPase (Takemiya and Shimazaki, 2010). In addition, ABA inhibits K⁺_{in}

158 channel activity and many ABA-signaling components affect this inhibition (Kim et al.,
159 2010; Jezek and Blatt, 2017). OPEN STOMATA 1 (OST1), an ABA-activated protein
160 kinase that operates downstream of the ABA-receptor components, suppresses K^+_{in}
161 channel KAT1 activity through direct phosphorylation (Sato et al., 2009). The second
162 messengers, such as Ca^{2+} , and G-proteins $G\alpha$ and $G\beta$ are also involved in ABA-induced
163 K^+_{in} channel inhibition (Fan et al., 2008; Kim et al., 2010). Moreover, the S-type anion
164 channels SLAC1 and SLAH3, which drive stomatal closure, are strongly upregulated by
165 drought stress in guard cells and inhibit KAT1 activity by direct binding (Zhang et al.,
166 2016). Second, ABA decreases the expression of K^+_{in} channel genes via inactivation of
167 AKS transcription factors through phosphorylation by OST1 in guard cells (Takahashi
168 et al., 2013; Takahashi et al., 2016). Third, ABA promotes internalization of KAT1 from
169 the PM into endomembrane compartments by endocytosis, thereby reducing the amount
170 of K^+_{in} channels functioning at the PM (Sutter et al., 2007).

171 Conversely, when plants are grown under well-watered conditions, blue light
172 suppresses signaling of ABA-induced stomatal closure to promote stomatal opening.
173 Blue light-receptor cryptochromes reduce ABA content in the plant body, and this
174 process is thought to affect ABA-signaling in guard cells (Boccalandro et al., 2012).
175 More directly, blue light also stimulates stomatal opening by suppressing anion release
176 from guard cells. This process involves light-dependent inhibition of S-type anion
177 channels in a phototropin-dependent manner (Marten et al., 2007). It was also reported
178 that light-produced phosphatidylinositol 4,5-bisphosphate inhibits anion channel
179 activity in guard cells (Lee et al., 2007). Consistent with these results, stomata in the
180 *slac1-1* mutant are slightly open under dark-adapted conditions and open larger than
181 those in wild type in response to light (Wang et al., 2012).

182

183 **Involvement of PM H^+ -ATPase in stomatal opening**

184 PM H^+ -ATPases, a family of P-type ATPases, consist of a functional polypeptide with
185 10 transmembrane domains and three cytosolic domains, including the N-terminal
186 domain, catalytic domain, and C-terminal autoinhibitory domain (Fig. 2). In
187 *Arabidopsis*, eleven genes encode PM H^+ -ATPases (*AHA1*–*AHA11*), and a
188 double-knockout mutant of *AHA1* and *AHA2*, which are predominantly expressed in all
189 cell types, displays an embryonic lethal phenotype (Haruta et al., 2010). The membrane
190 potential and proton gradient created by PM H^+ -ATPases energize multiple ion channels

Figure 2




Figure 2. Schematic structure of PM H⁺-ATPases.

PM H⁺-ATPases possess 10 transmembrane domains (TM1-10) and three cytosolic domains, including the N-terminal domain, catalytic domain, and C-terminal autoinhibitory domain containing the R-I and R-II regions (Palmgren, 2001). There are several phosphorylation sites in the C-terminal domain (Thr881, Ser899, Ser931, and Thr947). Thr881, Ser899, and Ser931 are phosphorylated by PSY1R, FERONIA, and PKS5, respectively. 14-3-3 protein binds to the phosphorylated penultimate threonine (Thr947). Phosphorylations of Thr881 and Thr947 lead to activation of PM H⁺-ATPase (Green). Phosphorylations of Ser899 and Ser931 lead to inactivation of PM H⁺-ATPase (Red). The numbering of the amino acid residues corresponds to *Arabidopsis* H⁺-ATPase2 (AHA2).

191 and various H⁺-coupled transporters in the PM for diverse physiological responses
192 including stomatal movement, phloem loading and unloading, xylem loading and
193 unloading, seed germination, solute uptake in roots, leaf movement, tip growth, and cell

194 expansion (Haruta et al., 2015; Wang et al., 2014b; Takahashi and Kinoshita, 2016). In
195 addition, proton transport through PM H⁺-ATPases controls cytosolic pH homeostasis
196 and apoplastic pH (Falhof et al., 2016). It is worthy of note that modeling analyses of
197 ion transport in the stomatal guard cells reveal importance of PM H⁺-ATPase activity
198 not only in proton gradient formation and pH control, but also in driving K⁺ Ca²⁺, anion
199 transport and metabolism in guard cells (Chen et al. 2012; Hills et al. 2012;
200 Minguet-Parramona et al. 2016), demonstrating that PM H⁺-ATPase plays a pivotal role
201 in guard cells physiology.

202 Since the 1970s, it has been known that light induces proton extrusion from stomatal
203 guard cells, K⁺ uptake, and swelling of guard cells. Physiological and
204 electro-physiological analyses revealed the properties of blue light-activated proton
205 pump using *Vicia* guard cell protoplasts (Assmann et al. 1985; Shimazaki et al., 1986).
206 Later, conclusive evidence was obtained by biochemical analysis. PM H⁺-ATPase is
207 activated and phosphorylated in response to blue light on a penultimate threonine
208 residue in the C-terminus and 14-3-3 protein binds to the phosphorylated C-terminus
209 region (Kinoshita and Shimazaki, 1999; Kinoshita and Shimazaki, 2002).

210 In addition to biochemical evidence, genetic evidence has also been obtained.
211 Dominant mutants of *AHA1*, a major H⁺-ATPase isoform in *Arabidopsis*, *ost2-1D* and
212 *ost2-2D*, displayed constitutively open stomatal phenotypes, since dominant mutations
213 cause constitutive activity of H⁺-ATPase (Merlot et al., 2007). More recently, it was
214 reported that loss of function mutants of *AHA1* showed reduced blue light-induced
215 stomatal opening (Yamauchi et al., 2016) and a closed stomatal phenotype (Osakabe et
216 al., 2016). Furthermore, a loss of function mutant of *OSA7*, a major PM H⁺-ATPase
217 isoform in rice, also showed reduced blue light-induced increase of stomatal
218 conductance (Toda et al., 2016). Overexpression of *AHA2* using a strong guard cell
219 promoter enhanced light-induced stomatal opening, leading to increased photosynthetic
220 activity and plant biomass (Yang et al., 2008; Wang et al., 2014a). These biochemical
221 and genetic findings clearly demonstrate that the PM H⁺-ATPases act as H⁺ pumps in
222 the PM and are vital for stomatal opening.

223

224 **Regulation of PM H⁺-ATPase by reversible phosphorylation**

225 It has been demonstrated that PM H⁺-ATPase activity is regulated by phosphorylation of
226 several sites (Haruta et al. 2015; Falhof et al. 2016). The first reported and most studied

227 phosphorylation site is the penultimate threonine in the C-terminus (Palmgren, 2001;
228 Wang et al., 2014b). Blue light activates PM H⁺-ATPase via the guard cell-specific
229 signaling pathway, leading to phosphorylation of the penultimate threonine in guard
230 cells (Shimazaki et al. 2007). In addition, recent investigations revealed that
231 phosphorylation level of the penultimate threonine in PM H⁺-ATPase were modulated in
232 response to physiological and environmental signals, such as light, salt, sucrose, auxin,
233 gibberellin, and ABA in several tissues and cell types besides guard cells (Niittylä et al.,
234 2007; Chen et al., 2010; Okumura et al., 2012; Takahashi et al., 2012; Hayashi et al.,
235 2014; Inoue et al., 2016; Okumura et al., 2016). These results indicate that there are
236 several unique signaling pathways in each tissue and cell type, but a final regulatory
237 mechanism, that is, phosphorylation of the penultimate threonine at the C-terminus of
238 the H⁺-ATPase, is common in these responses.

239 The protein kinase responsible for the phosphorylation of the penultimate threonine
240 in PM H⁺-ATPase has not been identified, despite strenuous efforts, although *in vitro*
241 protein kinase activity for the penultimate threonine of PM H⁺-ATPase was found in the
242 plasma membrane isolated from spinach leaves, the microsomes from guard cell
243 protoplasts of *Vicia*, and the plasma membrane from etiolated seedlings of *Arabidopsis*
244 (Kinoshita and Hayashi 2011). On the other hand, it was suggested that
245 dephosphorylation of the phosphorylated penultimate threonine of PM H⁺-ATPases is
246 mediated by the membrane-localized Mg²⁺/Mn²⁺-dependent protein phosphatase 2C
247 (PP2C)-like activity in *Vicia* guard cells and *Arabidopsis* etiolated seedlings (Hayashi et
248 al., 2010). Eventually, D-clade PP2Cs were shown to be involved in the
249 dephosphorylation of PM H⁺-ATPase in etiolated seedlings (Schweighofer et al. 2004;
250 Spartz et al., 2014; Ren and Gray, 2015). In addition, *SMALL AUXIN-UP RNAs*
251 (*SAURs*), a large multigene family of early-auxin-responsive genes, inhibit D-clade
252 PP2C activity through physical interaction (Spartz et al., 2014; Sun et al. 2016). It is
253 noteworthy that SAUR19-overexpressing plants displayed enhanced water loss in
254 detached leaves, wilted faster than wild type upon cessation of watering, and exhibited
255 delayed stomatal closure (Spartz et al., 2014; Spartz et al., 2017), and that many *SAUR*
256 genes are repressed by ABA, which certainly reduces stomatal aperture (Nemhauser et
257 al. 2006; Kodaira et al. 2011). Taken together these biochemical and genetic data
258 strongly suggest that clade D of PP2C and SAURs are involved in the regulation of PM
259 H⁺-ATPase in stomatal guard cells. However, it is still unknown that how SAURs are

260 regulated in response to blue light in guard cells.

261 In addition to the penultimate threonine, three residues were demonstrated to
262 regulate PM H⁺-ATPase activity using non-guard cells (Haruta et al., 2015;
263 Rudashevskaya et al., 2012) (Fig. 2). A receptor kinase, FERONIA, phosphorylates a
264 serine residue (Ser899 in AHA2) in the C-terminal autoinhibitory domain of PM
265 H⁺-ATPase and this phosphorylation suppresses proton efflux by PM H⁺-ATPase
266 (Haruta et al., 2014). Moreover, the phosphorylation of a threonine residue (Thr881 in
267 AHA2) in the C-terminal autoinhibitory domain of PM H⁺-ATPase is induced by a
268 receptor kinase, PSY1R, and application of the ligand peptide for PSY1R, PSY1,
269 increased proton efflux, suggesting that phosphorylation of Thr881 activates PM
270 H⁺-ATPase activity (Fuglsang et al., 2014). In addition, the serine residue (Ser931 in
271 AHA2) in the C-terminal autoinhibitory domain of the PM H⁺-ATPase is
272 phosphorylated by PKS5, a Ser/Thr protein kinase. Phosphorylation of Ser931 inhibits
273 interaction of the PM H⁺-ATPase with 14-3-3 protein and decreases PM H⁺-ATPase
274 activity (Fuglsang et al., 2007). The physical interaction of chaperone J3 with PKS5
275 induces the activation of PM H⁺-ATPase by repressing PKS5 activity (Yang et al., 2010).
276 In addition, it has been reported that type 2A protein phosphatase scaffolding subunit A
277 interacts with the C-terminus region of PM H⁺-ATPase (Fuglsang et al., 2006), and that
278 a negative regulator of plant immunity RIN4 activates PM H⁺-ATPase activity and the
279 *rin4* mutant shows reduced stomatal aperture (Liu et al., 2009). Thus, post-translational
280 modifications, such as phosphorylation of multiple sites in the C-terminal domain
281 and/or protein-protein interactions, regulate PM H⁺-ATPase activity. Blue light induces
282 phosphorylation on multiple serine and threonine residues in the C-terminus of the PM
283 H⁺-ATPase in *Vicia* guard cells (Kinoshita and Shimazaki, 1999). However, it is still
284 unclear whether other phosphorylation sites, such as Thr881, Ser899, and Ser931, are
285 involved in the regulation of PM H⁺-ATPase in guard cells in response to blue light.
286 Further investigations will be needed to clarify this.

287

288 **Conclusion and perspective**

289 In this review, we described recent advances in the blue light signaling pathway in
290 stomatal guard cells and the regulatory mechanisms of PM H⁺-ATPase. Stomata open in
291 response to blue light to facilitate gas exchange between the plant and the atmosphere.
292 This response is key to terrestrial plant life, as gas exchange is necessary not only for

293 photosynthesis but also for water uptake from the roots. So far, major signaling
294 components involved in the blue light signaling pathway in stomatal guard cells have
295 been identified, such as phototropin, BLUS1, BHP, PP1, and PM H⁺-ATPase; however,
296 the signaling mechanism is not fully understood. For example, how do blue light signals
297 induce phosphorylation of the penultimate threonine of PM H⁺-ATPase, which is a key
298 enzyme for stomatal opening? Whether blue light signal induces the phosphorylation of
299 PM H⁺-ATPase or suppresses dephosphorylation of PM H⁺-ATPase, or both, is still
300 unknown. Elucidation of this mechanism and identification of the kinase and
301 phosphatase for PM H⁺-ATPase in guard cells will provide novel insights into both the
302 blue light signaling pathway through phototropins and the regulation of H⁺-ATPase in
303 plant cells. Given the importance of stomatal regulation, future investigations will not
304 only improve our understanding of the molecular mechanisms of signaling pathways in
305 plants, but also provide important clues for agricultural strategies to improve
306 photosynthetic or water use efficiency, leading to an increase in the biomass and harvest
307 of crops.

308
309

310 **Figure legends**

311

312 Figure 1. Blue light signaling pathway in stomatal guard cells.

313 Arrows and a T-bar represent positive and negative regulation, respectively. The P in the
314 white circles indicates a phosphorylation of each protein. phot, phototropins; BLUS1,
315 BLUE LIGHT SIGNALING1; BHP, BLUE LIGHT-DEPENDENT H⁺-ATPASE
316 PHOSPHORYLATION; PP1, Protein phosphatase 1; PRSL1, PP1 regulatory
317 subunit2-like protein1; 14-3-3, 14-3-3 protein; K⁺_{in} channel, inward rectifying K⁺
318 channel; TAG, triacylglycerols; PAR, photosynthetically active radiation; Chl.,
319 Chloroplasts; NHX1/2, Na⁺/H⁺ EXCHANGER 1/2; ALMT9, aluminium-activated
320 malate transporter 9; CLCc, chloride channel c. The time scale of each peak of the key
321 signaling events for blue light-induced stomatal opening (approximately 2 h) is shown
322 as follows: phototropin activation (within 1 min), H⁺ pumping (approximately 2.5 min),
323 Hyperpolarization (several min), K⁺ accumulation (between 30 and 60 min). TAG
324 breakdown, starch degradation, and vacuolar remodeling are observed within 1 to 2 h
325 after the start of light illumination.

326

327 Figure 2. Schematic structure of PM H⁺-ATPases.

328 PM H⁺-ATPases possess 10 transmembrane domains (TM1-10) and three cytosolic
329 domains, including the N-terminal domain, catalytic domain, and C-terminal
330 autoinhibitory domain containing the R-I and R-II regions (Palmgren, 2001). There are
331 several phosphorylation sites in the C-terminal domain (Thr881, Ser899, Thr924,
332 Ser931, and Thr947). Thr881, Ser899, and Ser931 are phosphorylated by PSY1R,
333 FERONIA, and PKS5, respectively. 14-3-3 protein binds to the phosphorylated
334 penultimate threonine (Thr947). The numbering of the amino acid residues corresponds
335 to *Arabidopsis* H⁺-ATPase2 (AHA2).

336

337

338

ADVANCES

- Phototropin kinases activate the plasma membrane (PM) H⁺-ATPase in guard cells, which provides driving force for the accumulation of K⁺ through K⁺_{in} channels in the PM.
- ABA suppresses blue light-induced activation of PM H⁺-ATPase and K⁺_{in} channel via ABA receptor components PYR/PYL/RCAR-PP2Cs-SnRK2s in guard cells.
- The novel protein kinase BLUS1 acts as a phototropin substrate and transduces blue light signal to the PM H⁺-ATPase via the type 1 protein phosphatase and its regulatory subunit PRSL1.
- A Raf-like kinase BHP interacts with BLUS1 and mediates blue light signaling between BLUS1 and PM H⁺-ATPase.
- Degradation of starch in guard cell chloroplasts is required for phototropin-mediated signaling downstream of PM H⁺-ATPase activity and contributes to stomatal opening through malate synthesis.
- PP2C-Ds directly dephosphorylate the penultimate threonine of PM H⁺-ATPase in etiolated seedlings and are potentially involved in dephosphorylation of PM H⁺-ATPase in guard cells.

OUTSTANDING QUESTIONS

- How does the blue light signal induce phosphorylation of the penultimate threonine in PM H⁺-ATPase in guard cells? How do phototropin-BLUS1, BHP, PP1, and H⁺-ATPase connect in this signaling pathway?
- What are substrates of BLUS1, BHP, and PP1 in guard cells?
- What protein kinase is involved in direct phosphorylation of the penultimate threonine of PM H⁺-ATPase in guard cells and other cell types?
- Are PP2C-Ds involved in direct dephosphorylation of PM H⁺-ATPase in guard cells?
- How does PM H⁺-ATPase activation induce degradation of starch in guard cell chloroplasts?
- What is a next game-changing technique, following patch-clamp, gas-exchange, and thermal imaging, in stomatal research? Are live-cell imaging, chemical biology, and comprehensive proteomics in guard cells powerful approaches for future studies?

Parsed Citations

Ando E, Ohnishi M, Wang Y, Matsushita T, Watanabe A, Hayashi Y, Fujii M, Ma JF, Inoue S, Kinoshita T (2013) TWN SISTER OF FT, GIGANTEA, and CONSTANS have a positive but indirect effect on blue light-induced stomatal opening in *Arabidopsis thaliana*. *Plant Physiol* 162: 1529-1538.

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Andrés Z, Pérez-Hormaeche J, Leidi EO, Schlücking K, Steinhorst L, McLachlan DH, Schumacher K, Hetherington AM, Kudla J, Cubero B, Pardo JM (2014) Control of vacuolar dynamics and regulation of stomatal aperture by tonoplast potassium uptake. *Proc Natl Acad Sci USA* 111: 1806-1814

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Assmann SM, Simoncini L, Schroeder JI (1985) Blue-light activates electrogenic ion pumping in guard cell protoplasts of *Vicia faba*. *Nature* 318: 285-287

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Chen Y, Hoehenwarter W, Weckwerth W (2010) Comparative analysis of phytohormone-responsive phosphoproteins in *Arabidopsis thaliana* using TiO₂-phosphopeptide enrichment and mass accuracy precursor alignment. *Plant J* 63: 1-17

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Bauer H, Ache P, Lautner S, Fromm J, Hartung W, Al-Rasheid KAS, Sonnewald S, Sonnewald U, Kneitz S, Lachmann N, Mendel RR, Bittner F, Hetherington AM, Hedrich R (2013) The stomatal response to reduced relative humidity requires guard cell-autonomous ABA synthesis. *Curr Biol* 23: 53-57

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Boccalandro HE, Giordano CV, Ploschuk EL, Piccoli PN, Bottini R, Casal JJ (2012) Phototropins but not cryptochromes mediate the blue light-specific promotion of stomatal conductance, while both enhance photosynthesis and transpiration under full sunlight. *Plant Physiol* 158: 1475-1484

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Chen ZH, Hills A, Bätz U, Arntmann A, Lew VL, Blatt MR (2012) Systems dynamic modeling of the stomatal guard cell predicts emergent behaviors in transport, signaling, and volume control. *Plant Physiol* 159: 1235-1251

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Christie JM (2007) Phototropin blue-light receptors. *Annu Rev Plant Biol* 58: 21-45

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Darwin F (1989) Observations on stomata. *Philosophical Transactions of the Royal Society of London, Series B* 190: 531-621.

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

De Angeli A, Zhang J, Meyer S, Martinoia E (2013) AtALMT9 is a malate-activated vacuolar chloride channel required for stomatal opening in *Arabidopsis*. *Nat Commun.* 4: 1804

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Eisenacha C, De Angeli A (2017) Ion transport at the vacuole during stomatal movements. *Plant Physiol.* in press.

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Falhof J, Pedersen JT, Fuglsang AT, Palmgren M (2016) Plasma membrane H⁺-ATPase regulation in the center of plant physiology. *Mol Plant* 9: 323-337

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Fan LM, Zhang W, Chen JG, Taylor JP, Jones AM, Assmann SM (2008) Abscisic acid regulation of guard-cell K⁺ and anion channels in G[?]- and RGS-deficient *Arabidopsis* lines. *Proc Natl Acad Sci U S A* 105: 8476-8481

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Fujii H, Chinnusamy V, Rodrigues A, Rubio S, Antoni R, Park SY, Cutler SR, Sheen J, Rodriguez PL, Zhu JK (2009) In vitro reconstitution of an abscisic acid signalling pathway. Nature 462: 660-664

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Fuglsang AT, Tulinius G, Cui N, Palmgren MG (2006) Protein phosphatase 2A scaffolding subunit A interacts with plasma membrane H⁺-ATPase C-terminus in the same region as 14-3-3 protein. Physiol Plant 128: 334-340

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Fuglsang AT, Guo Y, Cuin TA, Qiu Q, Song C, Kristiansen KA, Bych K, Schulz A, Shabala S, Schumaker KS, Palmgren MG, Zhu JK (2007) Arabidopsis protein kinase PKS5 inhibits the plasma membrane H⁺-ATPase by preventing interaction with 14-3-3 protein. Plant Cell 19: 1617-1634

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Fuglsang AT, Kristensen A, Cuin TA, Schulze WX, Persson J, Thuesen KH, Ytting CK, Oehlenschlaeger CB, Mahmood K, Sondergaard TE, Shabala S, Palmgren MG (2014) Receptor kinase-mediated control of primary active proton pumping at the plasma membrane. Plant J 80: 951-964

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Haruta M, Burch HL, Nelson RB, Barrett-Wilt G, Kline KG, Mohsin SB, Young JC, Otegui MS, Sussman MR (2010) Molecular characterization of mutant Arabidopsis plants with reduced plasma membrane proton pump activity. J Biol Chem 285: 17918-17929

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Haruta M, Sabat G, Stecker K, Minkoff BB, Sussman MR (2014) A peptide hormone and its receptor protein kinase regulate plant cell expansion. Science 343: 408-411

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Haruta M, Gray WM, Sussman MR (2015) Regulation of the plasma membrane proton pump (H⁺-ATPase) by phosphorylation. Curr Opin Plant Biol 28: 68-75

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Hashimoto-Sugimoto M, Higaki T, Yaeno T, Nagami A, Irie M, Fujimi M, Miyamoto M, Akita K, Negi J, Shirasu K, Hasezawa S, Iba K (2013) A Munc13-like protein in Arabidopsis mediates H⁺-ATPase translocation that is essential for stomatal responses. Nat Commun 4: 2215

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Hayashi M, Inoue S, Takahashi K, Kinoshita T (2011) Immunohistochemical detection of blue light-induced phosphorylation of the plasma membrane H⁺-ATPase in stomatal guard cells. Plant Cell Physiol 52: 1238-1248

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Hayashi M, Kinoshita T (2011) Crosstalk between blue-light- and ABA-signaling pathways in stomatal guard cells. Plant Signal Behav 6: 1662-1664

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Hayashi M, Inoue S, Ueno Y, Kinoshita T (2017) A Raf-like protein kinase BHP mediates blue light-dependent stomatal opening. Sci Rep 7: 45586

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Hayashi Y, Nakamura S, Takemiya A, Takahashi Y, Shimazaki S, Kinoshita T (2010) Biochemical characterization of in vitro phosphorylation and dephosphorylation of the plasma membrane H⁺-ATPase. Plant Cell Physiol 51: 1186-1196

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Hayashi Y, Takahashi K, Inoue S, Kinoshita T (2014) Abscisic acid suppresses hypocotyl elongation by dephosphorylating plasma membrane H⁺-ATPase in Arabidopsis thaliana. Plant Cell Physiol 55: 845-853

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)
Google Scholar: [Author Only Title Only Author and Title](#)

Hills A, Chen ZH, Amtmann A, Blatt MR, Lew VL (2012) OnGuard, a computational platform for quantitative kinetic modeling of guard cell physiology. Plant Physiol 159: 1026-1042

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only Title Only Author and Title](#)

Horrer D, Flütsch S, Pazmino D, Matthews JS, Thalmann M, Nigro A, Leonhardt N, Lawson T, Santelia D (2016) Blue light induces a distinct starch degradation pathway in guard cells for stomatal opening. Curr Biol 26: 362-370

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only Title Only Author and Title](#)

Inoue S, Kinoshita T, Matsumoto M, Nakayama K, Doi M, Shimazaki K (2008) Blue light-induced autophosphorylation of phototropin is a primary step for signaling. Proc Natl Acad Sci USA 105: 5626-5631

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only Title Only Author and Title](#)

Inoue S, Takemiya A, Shimazaki K (2010) Phototropin signaling and stomatal opening as a model case. Curr Opin Plant Biol 13: 587-593

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only Title Only Author and Title](#)

Inoue S, Matsushita T, Tomokiyo Y, Matsumoto M, Nakayama KI, Kinoshita T, Shimazaki K (2011) Functional analyses of the activation loop of phototropin2 in Arabidopsis. Plant Physiol 156: 117-128

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only Title Only Author and Title](#)

Inoue S, Takahashi K, Okumura-Noda H, Kinoshita T (2016) Auxin influx carrier AUX1 confers acid resistance for Arabidopsis root elongation through the regulation of plasma membrane H⁺-ATPase. Plant Cell Physiol 57: 2194-2201.

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only Title Only Author and Title](#)

Jezeq M, Blatt MR (2017) The membrane transport system of the guard cell and its integration for stomatal dynamics. Plant Physiol. in press.

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only Title Only Author and Title](#)

Jossier M, Kroniewicz L, Dalmás F, Le Thiec D, Ephritikhine G, Thomine S, Barbier-Brygoo H, Vavasseur A, Filleur S, Leonhardt N (2010) The Arabidopsis vacuolar anion transporter, AtCLCc, is involved in the regulation of stomatal movements and contributes to salt tolerance. Plant J 64: 563-576.

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only Title Only Author and Title](#)

Kim TH, Böhmer M, Hu H, Nishimura N, Schroeder JI (2010) Guard cell signal transduction network: advances in understanding abscisic acid, CO₂, and Ca²⁺ signaling. Annu Rev Plant Biol 61: 561-591

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only Title Only Author and Title](#)

Kimura Y, Aoki S, Ando E, Kitatsuji A, Watanabe A, Ohnishi M, Takahashi K, Inoue S, Nakamichi N, Tamada Y, Kinoshita T (2015) A flowering integrator, SOC1, affects stomatal opening in Arabidopsis thaliana. Plant Cell Physiol 56: 640-649

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only Title Only Author and Title](#)

Kinoshita T, Shimazaki K (1999) Blue light activates the plasma membrane H⁺-ATPase by phosphorylation of the C-terminus in stomatal guard cells. EMBO J 18: 5548-5558

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only Title Only Author and Title](#)

Kinoshita T, Doi M, Suetsugu N, Kagawa T, Wada M, Shimazaki K (2001) phot1 and phot2 mediate blue light regulation of stomatal opening. Nature 414: 656-660

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only Title Only Author and Title](#)

Kinoshita T, Shimazaki K (2002) Biochemical evidence for the requirement of 14-3-3 protein binding in activation of the guard-cell plasma membrane H⁺-ATPase by blue light. Plant Cell Physiol 43: 1359-1365

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only Title Only Author and Title](#)

Kinoshita T, Emi T, Tominaga M, Sakamoto K, Shigenaga A, Doi M, Shimazaki K (2003) Blue light- and phosphorylation-dependent binding of a 14-3-3 protein to phototropins in stomatal guard cells of broad bean. *Plant Physiol* 133: 1453-1463

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Kinoshita T, Hayashi Y (2011) New insights into the regulation of stomatal opening by blue light and plasma membrane H⁺-ATPase. *Int Rev Cell Mol Biol* 289: 89-115

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Kinoshita T, Ono N, Hayashi Y, Morimoto S, Nakamura S, Soda M, Kato Y, Ohnishi M, Nakano T, Inoue S, Shimazaki K (2011) FLOWERING LOCUS T regulates stomatal opening. *Curr Biol* 21: 1232-1238

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Kodaira KS, Qin F, Tran LS, Maruyama K, Kidokoro S, Fujita Y, Shinozaki K, Yamaguchi-Shinozaki K (2011) Arabidopsis Cys2/His2 zinc-finger proteins AZF1 and AZF2 negatively regulate abscisic acid-repressive and auxin-inducible genes under abiotic stress conditions. *Plant Physiol* 157: 742-756.

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Lebaudy A, Hosy E, Simonneau T, Sentenac H, Thibaud JB, Dreyer I (2008) Heteromeric K⁺ channels in plants. *Plant J* 54: 1076-1082.

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Lee Y, Kim YW, Jeon BW, Park KY, Suh SJ, Seo J, Kwak JM, Martinoia E, Hwang I, Lee Y (2007) Phosphatidylinositol 4,5-bisphosphate is important for stomatal opening. *Plant J* 52: 803-816

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Liu J, Elmore JM, Fuglsang AT, Palmgren MG, Staskawicz BJ, Coaker G (2009) RIN4 functions with plasma membrane H⁺-ATPases to regulate stomatal apertures during pathogen attack. *PLoS Biol* 7: e10000139

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Ma Y, Szostkiewicz I, Korte A, Moes D, Yang Y, Christmann A, Grill E (2009) Regulators of PP2C phosphatase activity function as abscisic acid sensors. *Science* 324: 1064-1068

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

McLachlan DH, Lan J, Geilfus CM, Dodd AN, Larson T, Baker A, Hörak H, Kollist H, He Z, Graham I, Mickelbart MV, Hetherington AM (2016) The breakdown of stored triacylglycerols is required during light-induced stomatal opening. *Curr Biol* 26: 707-172

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Marten H, Hedrich R, Roelfsema MR (2007) Blue light inhibits guard cell plasma membrane anion channels in a phototropin-dependent manner. *Plant J* 50: 29-39

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Marten I, Deeken R, Hedrich R, Roelfsema MR (2010) Light-induced modification of plant plasma membrane ion transport. *Plant Biol* 12 s1: 64-79

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Merlot S, Leonhardt N, Fenzi F, Valon C, Costa M, Piette L, Vavasseur A, Genty B, Boivin K, Muller A, Giraudat J, Leung J (2007) Constitutive activation of a plasma membrane H⁺-ATPase prevents abscisic acid-mediated stomatal closure. *EMBO J* 26: 3216-3226

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Minguet-Parramona C, Wang Y, Hills A, Vialet-Chabrand S, Griffiths H, Rogers S, Lawson T, Lew V, Blatt MR (2016) An optimal frequency in Ca²⁺ oscillations for stomatal closure is an emergent property of ion transport in guard cells. *Plant Physiol* 170: 33-42

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Mott KA, Sibbersen ED, Shope JC (2008) The role of the mesophyll in stomatal responses to light and CO₂. Plant Cell Environ 31: 1299-1306.

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Murata Y, Mori IC, Munemasa S (2015) Diverse stomatal signaling and the signal integration mechanism. Annu Rev Plant Biol 66: 369-392.

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Nagatoshi Y, Mitsuda N, Hayashi M, Inoue S, Okuma E, Kubo A, Murata Y, Seo M, Saji H, Kinoshita T, Ohme-Takagi M (2016) GOLDEN 2-LIKE transcription factors for chloroplast development affect ozone tolerance through the regulation of stomatal movement. Proc Natl Acad Sci U S A 113: 4218-4223

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Negi J, Matsuda O, Nagasawa T, Oba Y, Takahashi H, K-Y M, Uchimiya H, Hashimoto M, Iba K (2008) CO₂ regulator SLAC1 and its homologues are essential for anion homeostasis in plant cells. Nature 452: 483-486

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Nemhauser JL, Hong F, Chory J (2006) Different plant hormones regulate similar processes through largely nonoverlapping transcriptional responses. Cell 126: 467-475.

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Niittylä T, Fuglsang AT, Palmgren MG, Frommer WB, Schulze WX (2007) Temporal analysis of sucrose-induced phosphorylation changes in plasma membrane proteins of Arabidopsis. Mol Cell Proteomics 6: 1711-1726

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Okumura M, Inoue S, Takahashi K, Ishizaki K, Kohchi T, Kinoshita T (2012) Characterization of the plasma membrane H⁺-ATPase in the liverwort *Marchantia polymorpha*. Plant Physiol 159: 826-834

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Okumura M, Inoue S, Kuwata K, Kinoshita T (2016) Photosynthesis activates plasma membrane H⁺-ATPase via sugar accumulation. Plant Physiol 171: 580-589

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Osakabe Y, Arinaga N, Umezawa T, Katsura S, Nagamachi K, Tanaka H, Ohiraki H, Yamada K, Seo SU, Abo M, Yoshimura E, Shinozaki K, Yamaguchi-Shinozaki K (2013) Osmotic stress responses and plant growth controlled by potassium transporters in Arabidopsis. Plant Cell 25: 609-624

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Osakabe Y, Yamaguchi-Shinozaki K, Shinozaki K, Tran L-SP (2014) ABA control of plant macroelement membrane transport systems in response to water deficit and high salinity. New Phytol 202: 35-49

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Osakabe Y, Watanabe T, Sugano SS, Ueta R, Ishihara R, Shinozaki K, Osakabe K (2016) Optimization of CRISPR/Cas9 genome editing to modify abiotic stress responses in plants. Sci Rep 6: 26685

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Palmgren MG (2001) Plant plasma membrane H⁺-ATPases: powerhouses for nutrient uptake. Annu Rev Plant Physiol Plant Mol Biol 52: 817-845

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Papanatsiou M, Scuffi D, Blatt MR, Garcia-Mata C (2015) Hydrogen sulfide regulates inward-rectifying K⁺ channels in conjunction with stomatal closure. Plant Physiol 168: 29-35

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Park SY, Fung P, Nishimura N, Jensen DR, Fujii H, Zhao Y, Lumba S, Santiago J, Rodrigues A, Chow TF, Alfred SE, Bonetta D, Finkelstein R, Provart NJ, Desveaux D, Rodriguez PL, McCourt P, Zhu JK, Schroeder JI, Volkman BF, Cutler SR (2009) Abscisic acid inhibits type 2C protein phosphatases via the PYR/PYL family of START proteins. *Science* 324: 1068-1071

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Poulsen LR, López-Marqués RL, Pedas PR, McDowell SC, Brown E, Kunze R, Harper JF, Pomorski TG, Palmgren M (2015) A phospholipid uptake system in the model plant *Arabidopsis thaliana*. *Nat Commun* 6: 7649

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Ren H, Gray WM (2015) SAUR proteins as effectors of hormonal and environmental signals in plant growth. *Mol Plant* 8: 1153-1164

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Rudashevskaya EL, Ye J, Jensen ON, Fuglsang AT, Palmgren MG (2012) Phosphosite mapping of P-type plasma membrane H⁺-ATPase in homologous and heterologous environments. *J Biol Chem* 287: 4904-4913

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Sato A, Sato Y, Fukao Y, Fujiwara M, Umezawa T, Shinozaki K, Hibi T, Taniguchi M, Miyake H, Goto DB, Uozumi N (2009) Threonine at position 306 of the KAT1 potassium channel is essential for channel activity and is a target site for ABA-activated SnRK2/OST1/SnRK2.6 protein kinase. *Biochem J* 424: 439-448

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Schroeder JI, Raschke K, Neher E (1987) Voltage dependence of K⁺ channels in guard-cell protoplasts. *Proc Natl Acad Sci USA* 84: 4108-4112

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Schweighofer A, Hirt H, Meskiene I (2004) Plant PP2C phosphatases: emerging functions in stress signaling. *Trends Plant Sci* 9: 236-243.

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Scuffi D, Álvarez C, Laspina N, Gotor C, Lamattina L, Garcia-Mata C (2014) Hydrogen sulfide generated by L-cysteine desulfhydrase acts upstream of nitric oxide to modulate abscisic acid-dependent stomatal closure. *Plant Physiol* 166: 2065-2076

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Santelia D, Lunn JE (2017) Transitory starch metabolism in guard cells: unique features for a unique function. *Plant Physiol* in press.

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Seo J, Lee HY, Choi H, Choi Y, Lee Y, Kim YW, Ryu SB, Lee Y (2008) Phospholipase A2 γ mediates light-induced stomatal opening in *Arabidopsis*. *J Exp Bot* 59: 3587-3594

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Shimazaki K, Iino M, Zeiger E (1986) Blue light-dependent proton extrusion by guard-cell protoplasts of *Vicia faba*. *Nature* 319: 324-326

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Shimazaki K, Doi M, Assmann SM, Kinoshita T (2007) Light regulation of stomatal movement. *Annu Rev Plant Biol* 58: 219-247

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Spartz AK, Ren H, Park MY, Grandt KN, Lee SH, Murphy AS, Sussman MR, Overvoorde PJ, Gray WM (2014) SAUR inhibition of PP2C-D phosphatases activates plasma membrane H⁺-ATPases to promote cell expansion in *Arabidopsis*. *Plant Cell* 26: 2129-2142

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Spartz AK, Lor VS, Ren H, Olszewski NE, Miller ND, Wu G, Spalding EP, Gray WM (2017) Constitutive expression of *Arabidopsis* SMALL AUXIN UP RNA19 (SAUR19) in *Tomato* confers auxin-independent hypocotyl elongation. *Plant Physiol* 173: 1453-1462

Downloaded from on September 21, 2017 - Published by www.plantphysiol.org

Copyright © 2017 American Society of Plant Biologists. All rights reserved.

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Suetsugu N, Takami T, Ebisu Y, Watanabe H, Iiboshi C, Doi M, Shimazaki K (2014) Guard cell chloroplasts are essential for blue light-dependent stomatal opening in Arabidopsis. PLoS ONE 9: e108374

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Sutter J-U, Sieben C, Hartel A, Eisenach C, Thiel G, Blatt MR (2007) Abscisic acid triggers the endocytosis of the Arabidopsis KAT1 K⁺ channel and its recycling to the plasma membrane. Curr Biol 17: 1396-1402

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Takahashi K, Hayashi K, Kinoshita T (2012) Auxin activates the plasma membrane H⁺-ATPase by phosphorylation during hypocotyl elongation in Arabidopsis. Plant Physiol 159: 632-641

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Takahashi K, Kinoshita T (2016) The regulation of plant cell expansion: Auxin-induced turgor-driven cell elongation. Molecular Cell Biology of the Growth and Differentiation of Plant Cells, 156-173

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Takahashi Y, Ebisu Y, Kinoshita T, Doi M, Okuma E, Murata Y, Shimazaki K (2013) bHLH transcription factors that facilitate K⁺ uptake during stomatal opening are repressed by abscisic acid through phosphorylation. Sci Signal 6: ra48

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Takahashi Y, Kinoshita T, Matsumoto M, Shimazaki K (2016) Inhibition of the Arabidopsis bHLH transcription factor by monomerization through abscisic acid-induced phosphorylation. Plant J 87: 559-5567

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Takemiya A, Kinoshita T, Asanuma M, Shimazaki K (2006) Protein phosphatase 1 positively regulates stomatal opening in response to blue light in Vicia faba. Proc Natl Acad Sci USA 103: 13549-13554

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Takemiya A, Shimazaki K (2010) Phosphatidic acid inhibits blue light-induced stomatal opening via inhibition of protein phosphatase 1. Plant Physiol 153: 1555-1562

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Takemiya A, Sugiyama N, Fujimoto H, Tsutsumi T, Yamauchi S, Hiyama A, Tada Y, Christie JM, Shimazaki K (2013a) Phosphorylation of BLUS1 kinase by phototropins is a primary step in stomatal opening. Nat Commun 4: 2094

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Takemiya A, Yamauchi S, Yano T, Ariyoshi C, Shimazaki K (2013b) Identification of a regulatory subunit of protein phosphatase 1 which mediates blue light signaling for stomatal opening. Plant Cell Physiol 54: 24-35

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Takemiya A, Shimazaki K (2016) Arabidopsis phot1 and phot2 phosphorylate BLUS1 kinase with different efficiencies in stomatal opening. J Plant Res 129: 167-174

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Toda Y, Wang Y, Takahashi A, Kawai Y, Tada Y, Yamaji N, Ma JF, Ashikari M, Kinoshita T (2016) Oryza sativa H⁺-ATPase (OSA) is involved in the regulation of dumbbell-shaped guard cells of rice. Plant Cell Physiol 57: 1220-1230

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Umezawa T, Sugiyama N, Mizoguchi M, Hayashi S, Myouga F, Yamaguchi-Shinozaki K, Ishihama Y, Hirayama T, Shinozaki K (2009) Type 2C protein phosphatases directly regulate abscisic acid-activated protein kinases in Arabidopsis. Proc Natl Acad Sci USA 106: 17588-17593

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Vahisalu T, Kollist H, Wang Y-F, Nishimura N, Chan W-Y, Valerio G, Lamminmäki A, Brosché M, Moldau H, Desikan R, Schroeder JI, Kangasjärvi J (2008) SLAC1 is required for plant guard cell S-type anion channel function in stomatal signaling. Nature 452: 487-491

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Vlad F, Rubio S, Rodrigues A, Sirichandra C, Belin C, Robert N, Leung J, Rodriguez PL, Laurière C, Merlot S (2009) Protein phosphatases 2C regulate the activation of the Snf1-related kinase OST1 by abscisic acid in Arabidopsis. Plant Cell 21: 3170-3184

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Waadt R, Hitomi K, Nishimura N, Hitomi C, Adams SR, Getzoff ED, Schroeder JI (2014) FRET-based reporters for the direct visualization of abscisic acid concentration changes and distribution in Arabidopsis. eLife 3: e01739

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Wang SW, Li Y, Zhang XL, Yang HQ, Han XF, Liu ZH, Shang ZL, Asano T, Yoshioka Y, Zhang CG, Chen YL (2014) Lacking chloroplasts in guard cells of crumpled leaf attenuates stomatal opening: both guard cell chloroplasts and mesophyll contribute to guard cell ATP levels. Plant Cell Environ. 37: 2201-2210

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Wang Y, Papanatsiou M, Eisenach C, Karnik R, Williams M, Hills A, Lew VL, Blatt MR (2012) Systems dynamic modeling of a guard cell Cl⁻ channel mutant uncovers an emergent homeostatic network regulating stomatal transpiration. Plant Physiol 160: 1956-1967

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Wang Y, Noguchi K, Ono N, Inoue S, Terashima I, Kinoshita T (2014a) Overexpression of plasma membrane H⁺-ATPase in guard cells promotes light-induced stomatal opening and enhances plant growth. Proc Natl Acad Sci USA 111: 533-538

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Wang Y, Shimazaki K, Kinoshita T (2014b) Multiple roles of the plasma membrane H⁺-ATPase and its regulation. The Enzymes 35: 191-211

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Yamauchi S, Takemiya A, Sakamoto T, Kurata T, Tsutsumi T, Kinoshita T, Shimazaki K (2016) The plasma membrane H⁺-ATPase AHA1 plays a major role in stomatal opening in response to blue light. Plant Physiol 171: 2731-2743

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Yang Y, Costa A, Leonhardt N, Siegel RS, Schroeder JI (2008) Isolation of a strong Arabidopsis guard cell promoter and its potential as a research tool. Plant Method 4: 6

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Yang Y, Qin Y, Xie C, Zhao F, Zhao J, Liu D, Chen S, Fuglsang AT, Palmgren MG, Schumaker KS, Deng XW, Guo Y (2010) The Arabidopsis chaperone J3 regulates the plasma membrane H⁺-ATPase through interaction with the PKS5 kinase. Plant Cell 22: 1313-1332

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Zhang A, Ren HM, Tan YQ, Qi GN, Yao FY, Wu GL, Yang LW, Hussain J, Sun SJ, Wang YF (2016) S-type anion channels SLAC1 and SLAH3 function as essential negative regulators of inward K⁺ channels and stomatal opening in Arabidopsis. Plant Cell 28: 949-965

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Zhang X, Takemiya A, Kinoshita T, Shimazaki K (2007) Nitric oxide inhibits blue light-specific stomatal opening via abscisic acid signaling pathways in Vicia guard cells. Plant Cell Physiol 48: 715-723

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)